

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
8 July 2004 (08.07.2004)


PCT

(10) International Publication Number
WO 2004/057021 A2


- (51) International Patent Classification⁷: **C12Q 1/68** (72) Inventor; and
(75) Inventor/Applicant (for US only): **BOIVIN, Guy**
(21) International Application Number: **PCT/CA2003/001994** [CA/CA]; 2531 Montbray, Sainte-Foy, Québec G1V 1E7 (CA).
(22) International Filing Date: **19 December 2003 (19.12.2003)** (74) Agent: **OGILVY RENAULT**; Suite 1600, 1981 McGill College Avenue, Montreal, Québec H3A 2Y3 (CA).
(25) Filing Language: **English** (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
(26) Publication Language: **English** (84) Designated States (regional): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
(30) Priority Data:
2,411,264 19 December 2002 (19.12.2002) CA
2,418,004 24 January 2003 (24.01.2003) CA
(71) Applicant (for all designated States except US): **UNIVERSITÉ LAVAL [CA/CA]**; Danielle Allard, Pavillon des sciences de l'éducation, bureau 1474, Québec, Québec G1K 7P4 (CA).

[Continued on next page]

(54) Title: **MOLECULAR METHODS AND COMPOSITIONS FOR DETECTING AND QUANTIFYING RESPIRATORY VIRUSES**



(57) Abstract: There is provided compositions and methods for the detection and/or quantification of respiratory viruses using specific nucleic acid sequences. In particular methods and composition are provided to detect human Metapneumovirus (hMPV). The invention also provides with multiplex assays for detecting respiratory viruses in test samples.





ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE,
SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

- *without international search report and to be republished upon receipt of that report*

- 1 -

**MOLECULAR METHODS AND COMPOSITIONS FOR DETECTING AND
QUANTIFYING RESPIRATORY VIRUSES**

CROSS-REFERENCE TO RELATED APPLICATIONS

- 5 This application claims priority on Canadian applications serial number 2,411,264 filed on December 19, 2002 entitled "Sequences for detection and identification of the human metapneumovirus" and serial number 2,418,004 filed on January 24, 2003 entitled "Multiplex assays for detection and identification of viral respiratory pathogens.

10

TECHNICAL FIELD

This application relates to the field of microorganisms detection and more particularly to the detection of respiratory tract viruses using virus specific nucleic acids.

15

BACKGROUND OF THE INVENTION

- Respiratory tract infections are a significant cause of morbidity and mortality in all age groups but especially in young children, elderly subjects, and immunocompromised patients. Most of these infections are caused by viruses with influenza A and B and the human respiratory syncytial virus (hRSV) being associated with the most frequent and severe complications i.e. pneumonitis and death (Dowell et al., 1996, J. Infect. Dis. 174:456-462; Simonsen et al., 2000, J. Infect. Dis. 181:831-837; Han et al., 1999, J. Infect. Dis. 179:25-30; Neuzil et al., 2000, N. Engl. J. Med. 342:225-231; Simoes, 1999, Lancet 354:847-852.). Other viruses such as adenoviruses (> 50 serotypes), parainfluenza viruses (PIV) (4 serotypes), rhinoviruses (> 100 serotypes), and coronaviruses also contribute to acute respiratory tract infections (Trecanor, 2002, Clinical Virology, 2nd ed., ASM Press, Washington, D.C., 7-26). Also, recent studies by the inventor and others have suggested that the new human metapneumovirus (hMPV) should be added to the list of human respiratory viral pathogens in all age groups (van den Hoogen et al., 2001, Nat. Med. 7:719-724;

30

Peret et al., 2002, J. Infect. Dis. 185:1660-1663; Boivin et al., 2002, J. Infect. Dis. 186:1330-1334).

The hMPV was initially isolated from nasopharyngeal aspirates recovered from 28 children suffering from respiratory tract infections. Based on partial nucleic acid sequences, gene organization, and electron microscopy (EM) findings, this virus was preliminary assigned to the *Paramyxoviridae* family, subfamily *Pneumovirinae*, genus *Metapneumovirus*. Since then, hMPV has been isolated in North America by our group (Peret et al., 2002, J. Infect. Dis. 185:1660-1663), in Australia (Nissen et al., 2002, Med. J. Aust. 176:188), and in England (Stockton et al., 2002, Emerg. Infect. Dis. 8:897-901).

Preliminary virological characteristics of hMPV.

The initial report by Van Den Hoogen *et al.* (van den Hoogen et al., 2001, Nat. Med. 7:719-724) indicated that the new virus replicated slowly in tertiary monkey kidney (TMK) cells with cytopathic effects (CPE) consisting of syncytium formation appearing 10-14 days post inoculation. Notably, the virus grew very poorly in Vero and A549 cells and could not be propagated in Madin Darby canine kidney (MDCK) cells. Virus-infected TMK cell-culture supernatants also did not display any hemagglutinating activity. The supernatants of infected TMK cells were used for EM studies, which revealed the presence of paramyxovirus-like pleomorphic particles in the range of 150-600 nm with short envelope projections in the range of 13-17 nm. However, RT-PCR studies with primer sets specific for known paramyxoviruses were initially all negative. Using a random PCR amplification procedure known as RAP-PCR (Welsh et al., 1990, Nucleic Acids Res. 18:7213-7218; Ralph et al., 1993, Proc. Natl. Acad. Sci. U.S.A. 90:10710-10714), the Dutch investigators identified differentially displayed bands (compared to human parainfluenza-1 virus) which, upon sequencing, were found to be most closely related to the avian pneumovirus (APV).

APV, the agent of turkey rhinotracheitis, has been the sole member of the recently assigned *Metapneumovirus* genus within the *Pneumovirinae* subfamily (Taxonomy Virus, 2000, Seventh Report of the International Committee on Taxonomy of Viruses. San Diego. 551, 557, 559-560). The latter virus is a major respiratory pathogen of turkeys and other domestic fowl worldwide (Yu et al., 1992, Virology. 186:426-434; Jones et al., 1987, Vet. Rec. 120:301-302; Jones, 1996, Avian Pathol. 25:639-648; Seal et al., 2000, Virus Res. 66:139-147). There are four serotypes of APV with serotype C (present in US) being the most closely related to hMPV (van den Hoogen et al., 2001, Nat. Med. 7:719-724). The metapneumoviruses (including APV and hMPV) are structurally similar to the human respiratory syncytial virus (hRSV), which also belongs to the *Pneumovirinae* subfamily, except for two features: they lack the two non-structural proteins (NS1 and NS2) and they have a different genomic organization (Yu et al., 1992, Virology. 186:426-434). Recently, an almost complete hMPV sequence (13.4 kb) was reported in GenBank (accession number: AF371337) (van den Hoogen et al., 2002, Virology. 295:119-132). The overall percentage of amino acid sequence identity for the N (nucleoprotein), P (phosphoprotein), M (matrix), F (fusion), M2 (membrane), and L (polymerase) proteins is in the range of 56-88% between hMPV and APV whereas it is 18-44% between hMPV and hRSV. However, the G (glycoprotein) and SH (small hydrophobic) proteins of hMPV do not possess significant amino acid sequence identity with those of any other viruses. In contrast to APV, hMPV was shown to be unable to replicate in turkeys and chickens but the human virus replicated efficiently in the respiratory tract of cynomolgous macaques inducing upper respiratory tract symptoms that correlated with peak viral replication in throat samples between 2 to 8 days post infection (van den Hoogen et al., 2001, Nat. Med. 7:719-724). Thus, these preliminary data indicated that hMPV was a primate pathogen associated with respiratory diseases.

Clinical importance of hMPV.

There is little information about the clinical presentation of hMPV infection. In the original report by Van Den Hoogen *et al.*, 27 of the 28 infected children from the Netherlands were below the age of five (van den Hoogen *et al.*, 2001, Nat. Med. 7:719-724). It was mentioned that those infected patients were diagnosed with a range of acute respiratory tract infections (ARTI) from mild to severe including bronchiolitis and pneumonia although no detailed clinical data were provided. In a limited study conducted in the winter of 2000, the same authors found that 7 ($\approx 10\%$) of 68 pediatric patients with unexplained ARTI had evidence of acute hMPV infection by RT-PCR (van den Hoogen *et al.*, 2001, Nat. Med. 7:719-724). In addition, Australian investigators reported the detection of hMPV by RT-PCR in 3/200 (1.5%) nasopharyngeal aspirates from children whose specimens were negative for the common respiratory viruses during the winter of 2001 (Nissen *et al.*, 2002, Med. J. Aust. 176:188). Recently, hMPV RNA was detected by RT-PCR in 9/405 (2.2%) of nasopharyngeal swabs from English patients who consulted for influenza-like illnesses and had negative tests for other pathogens (Stockton *et al.*, 2002, Emerg. Infect. Dis. 8:897-901). These studies, however, suffer from major limitations such as their small size (in most cases), the superficial description of clinical features, the lack of morbidity/mortality data, their retrospective nature, the exclusion of patients who tested positive for other viruses, and the absence of a control group.

Thus far, seroprevalence studies have been limited to the Dutch population. An indirect immunofluorescence assay (IFA) using infected TMK cells has revealed that 25% of the children aged between 6 and 12 months had hMPV antibodies and 70% had become seropositive by the age of 5 years ($n=60$) (van den Hoogen *et al.*, 2001, Nat. Med. 7:719-724). Retrospective serological testing also indicates that hMPV has been circulating in the Dutch population for at least half a century.

Diagnosis of hMPV infections

Our laboratory was the first to report the presence of hMPV in America and the second in the world following the initial report from the Netherlands (Peret et al., 2002, *J. Infect. Dis.* 185:1660-1663). Our first 11 hMPV strains were isolated on LLC-MK2 cells, a continuous cell line from rhesus monkey kidneys that is used in our center to recover parainfluenza viruses (PIV). Electron microscopy studies of these isolates revealed pleomorphic, spherical and filamentous particles with characteristics (nucleocapsid diameter and pitch spacing, length of projections) that were consistent with paramyxoviruses. RT-PCR and immunofluorescent assays for paramyxoviruses (PIV 1-3, hRSV) and other respiratory viruses (adenoviruses, coronaviruses, influenza viruses A and B, rhinoviruses) were negative for all 11 isolates. In contrast, amplification of the hMPV F (fusion) and M (matrix) genes based on published partial sequences (van den Hoogen et al., 2001, *Nat. Med.* 7:719-724; van den Hoogen et al., 2002, *Virology*. 295:119-132) was positive for all isolates (Peret et al., 2002, *J. Infect. Dis.* 185:1660-1663).

Subsequently, the virological features and clinical findings associated with a larger set of 38 hMPV strains from subjects hospitalized at our institution (CHUL, Québec City) since 1993 were examined in our laboratory (Boivin et al., 2002, *J. Infect. Dis.* 186:1330-1334). With all cell lines used in our laboratory, we were able to determine that the previously unidentified hMPV strains only consistently grew on LLC-MK2 cells after a mean incubation time of 17.3 days (range 3-23 days). Interestingly, most isolates recovered before 2002 produced only limited syncytia on LLC-MK2 cells, whereas some of the recent strains produced large syncytia typical of those induced by hRSV. Notably, we also observed a seroconversion for hMPV in two adult patients for whom acute and convalescent sera were available using an IFA test designed in our laboratory. Such seroconversion episodes (titers from $< 1/10$ to $1/80$ and from $< 1/10$ to $1/160$ over 2 and 4 weeks, respectively) occurred during the course of the patients' illness suggesting that hMPV was a true respiratory pathogen. A phylogenetic analysis of hMPV Canadian strains based on F gene sequences identified two major phylogenetic groups. At the nucleotide level, we found a similarity of 94.2-100% and 92.6-100% for isolates of groups 1 and 2, respectively, whereas the similarity at the amino acid level was higher i.e. 96.8-100% (group 1) and

97.2-100% (group 2) suggesting a functional/immunological conservation of the F protein. Interestingly, co-circulation of the two hMPV groups during some years was observed.

5 Over the past 9 years, most (86.7%) hMPV strains from our center were isolated during a 5-month period, from December to May. As an indication of the relative frequency of detection of hMPV compared to other respiratory viruses, we reviewed all respiratory samples (n = 862) sent to our virology laboratory for viral culture during the winter season (Dec. to May) of 2000-01. During this period, hMPV was
10 recovered in 2.3% of all respiratory samples tested by viral culture compared to 2.3% for PIV, 3.4% for adenoviruses, 4.8% for hRSV, and 19.8% for influenza A or B viruses. Overall, hMPV represented 7.1% of all positive cultures from respiratory samples during that winter season with percentages of 4.1%, 2.8%, and 29.3% for patients aged 0-5, 5-65, and > 65 years, respectively. A retrospective review of the
15 clinical charts from hMPV-infected subjects identified for the first time the role of this virus as a cause of severe lower respiratory tract infections in all age groups but especially in young children, elderly subjects, and immunocompromised patients. More specifically, the most frequent diagnoses associated with hMPV infection in hospitalized children were pneumonitis (66%) and bronchiolitis (58.3%) whereas
20 bronchitis/bronchospasm (60%) and pneumonitis (40%) were most commonly seen in elderly subjects (Boivin et al., 2002, *J. Infect. Dis.* 186:1330-1334). Although no histopathological studies were available, hMPV was the only pathogen identified in respiratory tract secretions of 3 subjects (including 2 with immunosuppressive conditions) who died of respiratory insufficiency.

25

We recently studied in greater details one of the cases associated with a fatal hMPV infection (Pelletier et al., 2002, *Emerg. Infect. Dis.* 8:976-978). Interestingly, this child with acute lymphoblastic leukemia was successively infected by two genetically-distinct hMPV strains on two consecutive winter seasons. On each
30 occasion, hMPV was the only pathogen isolated from a nasopharyngeal aspirate obtained during the course of lower respiratory tract infections (bronchiolitis in the

first year, bronchiolitis followed by fatal pneumonitis in the second year). Comparative sequence analysis of the amplified F gene products (759 bp) revealed that the two viral isolates differed by 114 nucleotides (10 amino acids) and belonged to two different hMPV groups. Overall, our data suggest that many hMPV strains may co-circulate in a specific population during a given period and such viral diversity coupled with a waning immunity (as found in elderly subjects and immunocompromised patients) may lead to multiple and sometimes fatal reinfections similar to those reported with hRSV (Hall, 2001, N. Engl. J. Med. 344:1917-1928; Couch et al., 1997, Am. J. Med. 102:2-9).

10

The role and relative contribution of hMPV were recently further expanded in a prospective case-control study performed at our institution during the winter of 2001-02 (Dec. 15-April 20). All children ≤ 3 years were eligible as cases if they were hospitalized for a lower respiratory tract infection (bronchiolitis, croup, pneumonitis) whereas controls consisted of children matched for age and hospitalized for an elective surgery during the same period with no respiratory symptoms. Nasopharyngeal aspirates were obtained from all participants for real-time PCR studies performed on a LightCycler (Roche). The latter included a multiplex RT-PCR for common respiratory viruses (influenza A, influenza B, hRSV) and a real-time RT-PCR assay for hMPV (analytical sensitivity of 100 copies per PCR reaction). No viruses were detected in samples from the 51 control patients. In contrast, at least one virus was identified in samples from 163 (78.4%) of the 208 cases (51% hRSV, 21.6% influenza A, 0% influenza B, 5.8% hMPV, and 5.9% co-infections). hRSV and influenza A infections occurred predominantly in the period between January and March 2002, whereas hMPV infections increased after February. In fact, 11/12 hMPV cases occurred within a 4-week period from mid-March to mid-April accounting for 18.9% of all ARTI in children during this period. Data collected from our virology laboratory indicated that the hMPV outbreak extended after the end of the study protocol (April 20) which suggests that hMPV seasonal pattern may differ from that of other respiratory viruses. A bronchiolitis was diagnosed in 67% of hMPV-infected children whereas 16.7% developed a pneumonia. In brief, there was no statistical

30

difference between the different viruses with regard to the symptomatology and reported diagnoses. Similarly, the median duration of hospitalization for the 3 viruses was identical (4-5 days).

- 5 In view of the above, it will be appreciated that diagnosis assay that can rapidly detect the presence of many respiratory viruses is needed.

Respiratory viruses have been classically identified by viral culture using a variety of permissive cell lines. However, viral culture is not convenient for clinical
10 management due to the need of specialized techniques (cell culture, immunofluorescence methods), the long turnaround time before appearance of typical cytopathic effects for many viruses (up to a few weeks), and the need for rapid inoculation of an infectious virus into multiple cell lines for optimal sensitivity (Specter et al., 2002, Clinical Virology, 2nd ed., ASM Press, Washington, D.C., 243-
15 272).

Antigenic detection kits for some respiratory viruses (influenza viruses and hRSV) have been developed by many companies. Most are immunoenzymatic assays (ELISA) based on the capture of a viral antigen by a specific monoclonal antibody.
20 Although such assays eliminate many of the problems inherent to viral cultures (no need for specialized equipment, rapid turnaround time in < 30 min, no need to recover an infectious virus), they have not been widely adopted due to poor sensitivity and/or specificity compared to viral culture. For instance, the sensitivity and specificity of some rapid antigenic tests for influenza has varied from 46 to 96 % and from 52 to 99
25 %, respectively, compared to viral culture. The performance of these tests is dependent on many variables such as the patient's age, the type of specimen and the duration of the disease (Hayden et al., 2002, Clinical Virology, 2nd ed., ASM Press, Washington, D.C., 891-920). The performance of the rapid antigenic tests for hRSV has been considered quite good in children with sensitivity and specificity varying
30 from 80 to 95 % (Piedra et al., 2002, Clinical Virology, 2nd ed., ASM Press, Washington, D.C., 763-790) but poor in adults (Englund et al., 1996, J. Clin.

Microbiol. 34:1649-1653). Detection of antibodies against viral pathogens (serology) has a limited value in the diagnosis of acute viral infections since two serums, collected at a 2 to 3 week interval, are required in order to document a significant rise in viral antibody titers.

5

More recently, PCR or RT-PCR assays have been developed for many respiratory viruses allowing detection of small amounts of viral DNA or RNA in a clinical sample. In the so-called "multiplex" format, PCR assays have been designed to amplify more than one viral target in the same PCR reaction (Ellis et al., 1997, J. Clin. Microbiol. 35:2076-2082; Fan et al., 1998, Clin. Infect. Dis. 26:1397-1402; Liolios et al., 2001, J. Clin. Microbiol. 39:2779-2783; Grondahl et al., 1999, J. Clin. Microbiol. 37:1-7). However, most multiplex PCR assays reported to date do not allow the simultaneous detection of all viruses involved in acute respiratory tract infections of humans and they require separate steps for the amplification and detection of viral genes, greatly increasing the assay's turnaround time and preventing reliable quantification of the amplicons. At present, only one multiplex PCR assay (the Hexaplex test from Prodesse, Waukesha, WI) is commercially-available. The Hexaplex detects 6 respiratory viruses (influenza A and B, PIV 1-3 and hRSV) using specific viral probes in a microplate format (Fan et al., 1998, Clin. Infect. Dis. 26:1397-1402; Kehl et al., 2001, J. Clin. Microbiol. 39:1696-1701). This assay does not use real-time amplification methodologies and has a turnaround time of about one full working day. Thus, there is a need for developing more rapid molecular assays that permit detection of a larger panel of respiratory viral pathogens.

25 Rapid and sensitive detection of respiratory viruses is an important clinical goal to help physicians select the most appropriate antiviral, reduce nosocomial transmission of pathogens, decrease the length of hospitalization, and reduce antibiotic use and bacterial resistance. The present invention relates to the development of multiplex assays based on nucleic acids detection for sensitive and rapid diagnosis of most viral
30 respiratory pathogens of humans.

SUMMARY OF THE INVENTION

It is an object of the present invention to provide a rapid and sensitive method using amplification primers and/or probes for determining the presence and/or the amount in a test sample of nucleic acids from clinically important respiratory viral pathogens including human metapneumovirus.

In another aspect there is provided multiplex assays aimed at detecting conserved genes of common respiratory viral pathogens such as influenza A and B, human respiratory syncytial virus (hRSV) A and B, human metapneumovirus (hMPV), parainfluenza viruses (PIV) types 1-4, adenoviruses, rhinoviruses, enteroviruses, and coronaviruses.

The nucleic acid amplification method is preferably but not exclusively PCR. In a preferred embodiment, identification of the amplicons produced by the amplification of nucleic acids from respiratory viruses is achieved by verifying the specific melting temperature of amplified viral products or by hybridization with specific viral probes. Other amplification technologies including target and probe amplification techniques as well as signal amplification techniques performed in liquid phase or onto solid supports may also be used.

20

It is an object of the present invention to provide a method using nucleic acid amplification and/or detection for determining the presence and/or the amount of nucleic acids of the human metapneumovirus (hMPV).

In a preferred embodiment, the nucleic acid amplification method is PCR. However, other amplification technologies including target and probe amplification techniques as well as signal amplification performed in liquid phase or onto solid support may also be used.

11

This invention provides a method to detect the presence of hMPV in a test sample based on the detection of a nucleotide sequence from either the nucleocapsid (N), phosphoprotein (P), matrix (M), fusion (F), membrane (M2), polymerase (L), glycoprotein (G), and small hydrophobic (SH) genes. In a particularly preferred
5 embodiment, the method to detect the presence of hMPV in a test sample is based on the detection of a nucleotide sequence from the nucleoprotein (N) or polymerase (L) genes.

In another embodiment, identification of amplified hMPV products is achieved by
10 verifying the specific melting temperature of amplified viral products or by hybridization with hMPV-specific probes.

It is another object of this invention to provide a method for typing hMPV strains which comprises the steps of reproducing the above amplification method followed
15 by either hybridization using type-specific probes, DNA sequencing or restriction enzyme analysis. Typing is done preferentially, but not exclusively, on the most variable hMPV genes i.e. the fusion (F) and glycoprotein (G) genes.

All the proprietary DNA fragments and oligonucleolides useful in the detection or
20 identification of a hMPV strain are further objects of this invention as well as any composition of matters that include them (recombinant vectors, hosts, diagnostic kits...).

25 BRIEF DESCRIPTION OF THE DRAWINGS

Further features and advantages of the present invention will become apparent from the following detailed description, taken in combination with the appended drawings, in which:

12

Figure 1 represents a melting curve analysis of amplified hMPV strains using specific nucleoprotein primers (Sequence ID Nos. 92 and 94);

Figure 2 shows an ethidium bromide-stained agarose gel showing restriction fragment length polymorphism of amplified hMPV fusion (F) gene sequences digested with enzymes ApaI and Avr II. PCR products of the hMPV F gene (amplified with primers sequence ID Nos. 97 and 98) were digested with ApaI and Avr II. Strains belonging to hMPV genotype F-1 were digested only by ApaI resulting in two fragments of 532 bp and 227 bp whereas strains belonging to hMPV genotype F-2 were digested only by Avr II resulting in fragments of 447 and 312 bp;

Figure 3 shows the age at admission of children hospitalized for acute respiratory tract infections (ARTI) caused by hMPV (A), hRSV (B), and influenza A (C) as well as for the whole study population (D) described in example 4;

15

Figure 4 represents the biweekly distribution of virologically cases with acute respiratory tract infections and their controls in the study reported in example 4;

Figure 5 represents the biweekly distribution of hMPV isolates in the study group (hospitalized children) described in example 4 and in respiratory specimens tested at the regional virology laboratory (RVL);

Figure 6 illustrates a phylogenetic tree of hMPV strains recovered as part of the study described in example 4 also including the prototype strain from the Netherlands (GenBank af371367) and a Canadian isolate from season 2000-01 (hMPV35-2001). The dates of sample collection are in parentheses;

Figure 7 is a melting curve analysis of viral strains amplified using the respiratory-1 multiplex real-time PCR assay. A- Multiplex real-time PCR assay using hMPV N primers (SEQ ID NOS. 95 and 96). B- Multiplex real-time PCR assay using hMPV L

30

13

primers (SEQ ID NOS. 112 and 114). Note: the internal control is apparent (T_m : 91°C) in the absence of amplification of specific viral products.;

5 Figure 8 is a melting curve analysis of viral strains amplified using the respiratory-2 multiplex real-time PCR assay. Note: The internal control is apparent (T_m : 91°C) in the absence of amplification of specific viral products;

10 Figure 9A is a fluorescence signal as a function of elongation step for the determination of the hMPV load in clinical samples using quantitative real-time PCR;

Figure 9B is a standard curve for determination of the hMPV using quantitative real-time PCR;

15 Figure 10A is a phylogenetic tree for the F gene of hMPV;

Figure 10B is a phylogenetic tree for the G gene of hMPV;

20 Figure 11 is a melting curve analysis of amplified viral genes and of an internal control as determined by the LightCycler instrument. Note that high amounts of amplified viral products may preclude detection of the internal control in some PCR runs; and

Figure 12 is a melting curve analysis SARS-coronavirus using a real time PCR assay.

25 It will be noted that throughout the appended drawings, like features are identified by like reference numerals.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

The invention provides nucleic acid based diagnostic methods and compositions to detect and quantify respiratory viruses.

5

In a preferred embodiment respiratory viral pathogens targeted by the diagnostic assays include all genotypes of influenza A and B viruses, hRSV, hMPV, PIV, adenoviruses, rhinoviruses, and enteroviruses. Other less common (coronaviruses, reoviruses) and yet to be discovered respiratory viruses are also targets under the scope of the present invention.

10

In the description of this invention, the terms «nucleic acids» and «sequences» might be used interchangeably. However, «nucleic acids» are chemical entities while «sequences» are the pieces of information encoded by these «nucleic acids». Both nucleic acids and sequences are equivalently valuable sources of information for the matter pertaining to this invention.

15

In one embodiment the method targets conserved hMPV nucleotide sequences for diagnostic purposes and variable hMPV sequences for epidemiological typing purposes. Nucleotide sequences were selected either from a number of hMPV viral sequences obtained in the inventor's laboratory or from public databases (GenBank accession number AF371337).

20

Our proprietary DNA fragments and oligonucleotides (a non-restrictive list of primers and probes) are listed in the following tables.

25

It will be understood by those skilled in the art that sequences that can selectively hybridize to the sequences of the present invention are also comprised in the scope of the invention. By selectively hybridizing it is meant that a nucleic acid molecule binds to a given target in a manner that is detectable in a different manner from non-target

15

sequence under moderate, or more preferably under high, stringency conditions of *hybridization*. "Complementary" or "target" nucleic acid sequences refer to those nucleic acid sequences which selectively hybridize to a nucleic acid molecule. Proper annealing conditions depend, for example, upon a nucleic acid molecule's length, base composition, and the number of mismatches and their position on the molecule, and must often be determined empirically. For discussions of nucleic acid molecule (probe) design and annealing conditions, see, for example, Sambrook et al. 1989 (Sambrook, et al., *Molecular Cloning: A Laboratory Manual*, 2d Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)) or Ausubel et al. 1987 (Ausubel, F., et al., *Current Protocols in Molecular Biology*, New York (1987)).

The detection of the viruses including the hMPV is conducted through compositions of matters such as diagnostic kits comprising the amplification primers or probes of this invention.

15

In the methods and kits of the present invention, probes and primers are not limited to nucleic acids and may include, but are not restricted to, analogs of nucleotides. The diagnostic reagents constituted by the probes and the primers may be present in any suitable form (bound to a solid support, liquid, lyophilized, etc.) for detection and/or amplification of nucleic acid sequences of the viruses.

In the methods and kits of the present invention, amplification reactions may include but are not restricted to: a) polymerase chain reaction (PCR), b) ligase chain reaction (LCR), c) nucleic acid sequence-based amplification (NASBA), d) self-sustained sequence replication (3SR), e) strand displacement amplification (SDA), f) branched DNA signal amplification (bDNA), g) transcription-mediated amplification (TMA), h) cycling probe technology (CPT), i) nested PCR, j) multiplex PCR, k) solid phase amplification (SPA), l) nuclease dependent signal amplification (NDSA), m) rolling circle amplification technology (RCA), n) anchored strand displacement amplification, o) solid-phase (immobilized) rolling circle amplification.

In the methods and kits of the present invention, detection of the nucleic acids of target genes may include real-time or post-amplification technologies. These detection technologies can include, but are not limited to, the use of intercalating agents such as SYBR green, fluorescence resonance energy transfer (FRET)-based methods such as adjacent hybridization of probes (including probe-probe and probe-primer methods), *TaqMan* probe, molecular beacon probe, Scorpion probe, nanoparticle probes and Amplifluor probe. Other detection methods include target gene nucleic acids detection via immunological methods, solid phase hybridization methods on filters, chips or any other solid support. In these systems, the hybridization can be monitored by fluorescence, chemiluminescence, potentiometry, mass spectrometry, plasmon resonance, polarimetry, colorimetry, flow cytometry or scanometry. Nucleotide sequencing, including sequencing by dideoxy termination or sequencing by hybridization (e.g. sequencing using a DNA chip) represents another method to detect and characterize the nucleic acids of target genes.

Amplification of hMPV sequences

In one aspect of the invention there is provided methods and compositions for detecting hMPV in a sample.

In a preferred embodiment, a reverse-transcription (RT)-PCR protocol is used for nucleic acid amplification. The single-stranded viral RNA is converted to complementary DNA using an enzyme possessing a reverse transcriptase activity and a specific hMPV primer. Alternatively, random hexamer primers can be substituted to the specific hMPV primer. In a preferred embodiment, viral RNA was preably extracted from 200 μ l of naso-pharyngeal aspirates using the QIAamp Viral RNA Mini Kit (QIAGEN). Complementary DNA was synthesized using 10 μ l of eluted RNA, 0.75 μ M of a specific hMPV primer (Table 2, sequences ID Nos. 91, 97, 100, 102, 109, 112), and the Omniscript Reverse Transcriptase Kit (QIAGEN, Mississauga, ON, Canada) following the manufacturer's recommendations.

- For DNA amplification by the widely used PCR method, primer pairs were derived from our proprietary DNA fragments or from public database sequences (Table 1). During DNA amplification by PCR, two oligonucleotide primers binding respectively to each strand of the heat-denatured target DNA from the microbial genome are used
- 5 to amplify exponentially *in vitro* the target DNA by successive thermal cycles allowing denaturation of the DNA, annealing of the primers and synthesis of new targets at each cycle (Persing, 1993, Diagnostic Molecular Microbiology: Principles and Applications, American Society for Microbiology, Washington, DC. 88-109).
- 10 In a preferred embodiment, complementary DNA was amplified using a real-time PCR procedure with the LC Faststart DNA Master SYBR Green 1 Kit (Roche Diagnostics, Laval, QC, Canada) or the SYBR Green TAQ ReadyMix for quantitative PCR capillary formulation (Sigma-Aldrich, Oakville, ON) in a LightCycler instrument (Roche Diagnostics). Each reaction had a total volume of 20 μ l including 2
- 15 μ l of RT mixture and 18 μ l of a reaction mixture containing 4 mM $MgCl_2$, 2 μ l of Faststart DNA SYBR Green Master Mix or the SYBR Green TAQ ReadyMix for quantitative PCR capillary formulation, 3% DMSO, and 0.5 μ M of each hMPV primer (see Tables 2 and 4 for sequence ID Nos. and specific combinations of primers for amplification, respectively). Cycling conditions typically included an initial
- 20 denaturation step of 10 min. at 94°C, followed by 50 cycles of 15 s at 94°C, 5 s at 54°C, and 30 s at 72°C. However, the cycling conditions could slightly vary according to the hMPV gene to amplify.
- The person skilled in the art of nucleic acid and/or signal amplification knows the
- 25 existence of other rapid amplification procedures such as ligase chain reaction (LCR), reverse transcriptase PCR (RT-PCR), transcription-mediated amplification (TMA), self-sustained sequence replication (3SR), nucleic acid sequence-based amplification (NASBA), strand displacement amplification (SDA), branched DNA (bDNA), cycling probe technology (CPT), solid phase amplification (SPA), rolling circle
- 30 amplification technology (RCA), solid phase RCA, anchored SDA and nuclease dependent signal amplification (NDSA) (Persing, 1993, Diagnostic Molecular

Microbiology: Principles and Applications, American Society for Microbiology, Washington, DC. 88-109; Lee, 1997, Nucleic Acid Amplification Technologies: Applications to Disease Diagnosis, Eaton Publishing, Boston, MA. 61-125; Westin et al., 2000, Nat. Biotechnol. 18:199-204). The scope of this invention is not limited to the use of amplification by PCR, but rather includes the use of any nucleic acid amplification method or any other procedure which may be used to increase the sensitivity and/or the rapidity of nucleic acid-based diagnostic tests. The scope of the present invention also covers the use of any nucleic acid and/or signal amplification and detection technology including real-time or post-amplification detection technologies, any amplification technology combined with detection, any hybridization nucleic acid chips or array technologies, any amplification chips or combination of amplification and hybridization chip technologies. Detection and identification by any nucleotide sequencing method is also under the scope of the present invention.

15

Any oligonucleotide, useful for diagnosis, and which are derived from hMPV sequences and used with any nucleic acid amplification and/or hybridization technologies are also under the scope of this invention.

20 Detection of amplification products

Classically, the detection of PCR amplification products is performed by standard ethidium bromide-stained agarose gel electrophoresis. Using this approach, primers specific for a nucleotide sequence from either the nucleocapsid (N), phosphoprotein (P), matrix (M), fusion (F), polymerase (L), glycoprotein (G), genes generated amplicons of various lengths depending on the selected primers pairs (Table 4). It is however clear that other detection methods of amplified products may be faster, more sensitive, and more practical. Amplicon detection may also be performed by solid support or liquid hybridization using strain-specific internal DNA probes hybridizing to an amplification product. Such probes may be generated from sequences from our repertory (Table 1) and designed to specifically hybridize to DNA amplification

products which are objects of the present invention. Amplicons can also be characterized by DNA sequencing. Alternatively, a rapid detection method developed with real-time PCR assays consists of determining the specific melting temperature of amplicons at the end of the amplification reaction.

5

In a preferred embodiment, detection of hMPV amplicons was characterized using the melting curve analysis feature of a real-time PCR instrument. Briefly, following the last amplification cycle on a LightCycler instrument, the internal temperature was rapidly increased to 94°C then decreased to 60°C for 30 seconds, followed by slowly
10 increasing to 94°C at a rate of 0.1°C per second, with continuous fluorescence lecture. Using this procedure, the specific melting temperature obtained with the specific amplicons produced by the pairs of primers specific for the N (SEQ ID No 92+94), M (SEQ ID No 100+101), F (SEQ ID 97+98), P (SEQ ID No 102 +103) and L (SEQ ID No 112 + 113) gene sequences was found to be 82.63°C \pm 0.87°C, 82.75°C \pm 0.80°C,
15 83.29°C \pm 0.45°C, 80.48°C \pm 0.46°C, and 77.99 \pm 0.33°C, respectively (Table 5 and see also Figure 1 for example of the melting temperature obtained for different hMPV strains with the N primers). Alternatively, we also designed specific adjacent and TaqMan fluorescent probes for the various hMPV gene products (N, M, F, G, P and L) that can be used for real-time detection and quantification of amplicons (Table 3).
20 Here, quantification of hMPV gene products is achieved through fluorescence resonance energy transfer (FRET) when the two probes hybridize nearby on a complementary DNA sequence. Other types of detection methods including the use of TaqMan or molecular beacon probes specific for the hMPV sequences described above are also under the scope of this invention.

25

Typing of hMPV

Previous studies have shown that two major genotypes of hMPV circulate in North America and Europe (van den Hoogen et al., 2001, Nat. Med. 7:719-724; Boivin et
30 al., 2002, J. Infect. Dis., 186:1330-1334). We also designed a rapid molecular typing method for epidemiologic studies based on amplification of a portion of the F (fusion)

glycoprotein (primer sequence ID Nos. 97 and 98, Tables 2 and 4) followed by DNA sequencing or digestion with restriction enzymes (ApaI I and Avr II). Table 6 summarizes the size of the F fragments obtained for different hMPV isolates following enzymatic digestion and the corresponding genotype (see also Figure 2 for an example of the band pattern following electrophoresis on an agarose gel). Alternative typing methods based on the use of type-specific hMPV F primers and/or probes as well as DNA sequencing of amplified products are also within the scope of this invention. hMPV subtyping using similar methodologies for the other surface glycoprotein (i.e. the G gene) may also be used.

10

A method for detection of a plurality of potential hMPV strains having different genotypes may be conducted in separate reactions and physical enclosures, one type at the time. Alternatively, it could be conducted simultaneously for different types in separate physical enclosures, or in the same physical enclosures. In the latter scenario a multiplex PCR reaction could be conducted which would require that the oligonucleotides are all capable of annealing with a target region under common conditions. Since many probes or primers are specific for a determined genotype, typing an hMPV strain is under the scope of the present invention. When a mixture of oligonucleotides annealing together with more than one type of hMPV is used in a single physical enclosure or container, different detection labels would be used to distinguish one type from another.

15

The diagnostic kits, primers and/or probes mentioned above can be used to detect and/or identify hMPV, whether said diagnostic kits, primers and probes are used for *in vitro* or *in situ* applications. The said samples may include but are not limited to: any clinical sample, any environmental sample, any viral culture, any tissue, or any cell line.

20

It is also an object of the present invention that said diagnostic kits, primers and/or probes can be used alone or in combination with any other assay suitable to detect and/or identify other microorganisms (i.e. viruses, bacteria, fungi, and parasites) including but not limited to: any assay based on nucleic acids detection, any

25

30

immunoassay, any enzymatic assay, any biochemical assay, any lysotypic assay, any serological assay, any culture on specific cell lines, and any infectivity assay on animals.

- 5 In the methods and kits described herein below, the oligonucleotide probes and amplification primers have been derived from larger sequences (i.e. DNA fragments of at least 100 base pairs). All DNA sequences have been obtained either from our proprietary sequences (Table 1) or from SEQ ID No. 1 from public databases.
- 10 It is clear to the individual skilled in the art that oligonucleotide sequences other than those described in the present invention and which are appropriate for detection and/or identification of hMPV may also be derived from the proprietary fragment sequences or selected public database sequences. For example, the oligonucleotide primers or probes may be shorter but of a length of at least 10 nucleotides or longer
- 15 than the ones chosen; they may also be selected anywhere else in the proprietary DNA fragments or in the sequences selected from public databases; they may also be variants of the same oligonucleotide. If the target DNA or a variant thereof hybridizes to a given oligonucleotide, or if the target DNA or a variant thereof can be amplified by a given oligonucleotide PCR primer pair, the converse is also true : a given target
- 20 DNA may hybridize to a variant oligonucleotide probe or be amplified by a variant oligonucleotide PCR primer. Alternatively, the oligonucleotides may be designed from said DNA fragment sequences for use in amplification methods other than PCR. Consequently, the core of this invention is the detection and/or identification of hMPV by targeting DNA sequences which are used as a source of specific and
- 25 ubiquitous oligonucleotide probes and/or amplification primers. Although the selection and evaluation of oligonucleotides suitable for diagnostic purposes require much effort, it is quite possible for the individual skilled in the art to derive, from the selected DNA fragments, oligonucleotides other than the ones listed in Tables 2 and 3 which are suitable for diagnostic purposes. When a proprietary fragment or a public
- 30 database sequence is selected for its specificity and ubiquity (i.e. ability to detect all

hMPV strains), it increases the probability that subsets thereof will also be specific and ubiquitous.

The proprietary DNA fragments have been obtained as a repertory of sequences created by amplifying hMPV nucleic acids with proprietary primers. These primers and the repertory of nucleic acids as well as the repertory of nucleotide sequences are further objects of this invention (Tables 1-3).

As part of the design rules, all oligonucleotides were evaluated for their suitability for hybridization or PCR amplification by computer analysis using standard programs (i.e. the GCG Wisconsin package programs, the primer analysis software Oligo™ 6 and MFOLD 3.0). The potential suitability of the PCR primer pairs was also evaluated prior to their synthesis by verifying the absence of unwanted features such as long stretches of the same nucleotide and a high proportion of G or C residues at the 3' end (Persing, 1993, Diagnostic Molecular Microbiology: Principles and Applications, American Society for Microbiology, Washington, DC. 88-109). Oligonucleotide amplification primers were synthesized using an automated DNA synthesizer (Applied Biosystems).

The oligonucleotide sequence of primers or probes may be derived from either strand of the duplex DNA. The primers or probes may consist of the bases A, G, C, or T or analogs and they may be degenerated at one or more chosen nucleotide position(s) (Nichols et al., 1994, Nature. 369:492-493). Primers and probes may also consist of nucleotide analogs such as Locked Nucleic Acids (LNA) (Koskhin et al., 1998, Tetrahedron Lett. 54:3607-3630), and Peptide Nucleic Acids (PNA) (Egholm et al., 1993, Nature. 365:566-568). The primers or probes may be of any suitable length and may be selected anywhere within the DNA sequences from proprietary fragments, or from selected database sequences which are suitable for the detection of hMPV.

Variants for a given target microbial gene are naturally occurring and are attributable to sequence variation within that gene. For example, different strains of the same viral

23

species may have a single or more nucleotide variation(s) at the oligonucleotide hybridization site. The person skilled in the art is well aware of the existence of variant nucleic acids and/or sequences for a specific gene and that the frequency of sequence variations depends on the selective pressure during evolution on a given gene product. The detection of a variant sequence for a region between two PCR primers may be demonstrated by sequencing the amplification product. In order to show the presence of sequence variations at the primer hybridization site, one has to amplify a larger DNA target with PCR primers outside that hybridization site. Sequencing of this larger fragment will allow the detection of sequence variation at this primer hybridization site. A similar strategy may be applied to show variations at the hybridization site of a probe. Insofar as the divergence of the target nucleic acids and/or sequences or a part thereof does not affect significantly the sensitivity and/or specificity and/or ubiquity of the amplification primers or probes, variant microbial DNA is under the scope of this invention. Variants of the selected primers or probes may also be used to amplify or hybridize to a variant target DNA.

In an other aspect of the invention there is also provided newly isolated human metapneumoviruses deposited with International Depository Authority of Canada on December 11 2003 and bearing IDAC numbers 111203-01 and 111203-02 (also referred herein as CAN 97-83 and CAN 98-75 respectively).

In a further aspect of the invention the diagnostic method is designed at performing simultaneous detection of nucleic acids from a variety of respiratory viruses using specific primers and/or probes. The method is aimed at detecting and/or quantifying conserved viral sequences from respiratory viruses for diagnostic purposes. Target nucleotide sequences were selected public databases and from viral strain sequences obtained in the inventor's laboratory.

The specific respiratory viral pathogens targeted by the diagnostic assays include all genotypes of influenza A and B viruses, hRSV, hMPV, PIV, adenoviruses, rhinoviruses, and enteroviruses. Other less common (coronaviruses, reoviruses) and

yet to discover respiratory viruses are also targets under the scope of the present invention.

The specific combination of oligonucleotides (primers and/or probes) in a multiplex
5 format is also another object of this invention.

Composition of matters such as diagnostic kits comprising amplification primers and/or probes for the detection of respiratory viruses are also objects of the present invention.

10

Amplification of viral sequences

In a preferred embodiment, a PCR protocol is used for nucleic acid amplification. Multiplex assay, aimed at amplifying a plurality of respiratory viruses in the same
15 physical enclosure, is an object of the present invention.

We aim at developing DNA-based tests or diagnostic kits to detect and identify most respiratory viruses in a limited number of amplification reactions using specific primers (Table 10) and/or probes (Table 11). Although some of the primer sequences
20 used for amplification of the various respiratory viruses, with the exception of hMPV (SEQ. ID. Nos. 95 and 96 or 112 and 114), PIV-1 (SEQ. ID. Nos. 147 and 148), PIV-3 (SEQ. ID. Nos. 151 and 152), and adenovirus (SEQ. ID. Nos. 153), have been described previously by others (Fouchier et al., 2000, J. Clin. Microbiol. 38:4096-4101 for influenza A virus / SEQ. ID. Nos. 139 and 140; Li et al., 2001, J. Clin.
25 Microbiol. 39:696-704 for influenza B virus / SEQ. ID. Nos. 141 and 142; Mazulli et al., 1999, J. Infect. Dis. 180:1686-1689 for hRSV / SEQ. ID. Nos. 143 and 144; Osiowy et al., 1998, J. Clin. Microbiol. 36: 3149-3154 for PIV-1, PIV-3, and adenovirus / SEQ. ID. Nos. 145 and 146, 149 and 150, and 154 and 155, respectively; Atmar et al., 1993, J. Clin. Microbiol. 31: 2544-2546 for rhinovirus and enterovirus /
30 SEQ. ID. Nos. 160 and 161) and that one influenza A primer (SEQ. ID. No. 139) is

25

almost identical to the one described in another patent (U.S. patent number 6,015,664), the object of the present invention is to combine many sets of primers and/or probes in the same physical enclosure (multiplex format).

5 Briefly, viral RNA was extracted from 200 µl of naso-pharyngeal aspirates or other respiratory tract samples using the QIAamp Viral RNA Mini kit (QIAGEN, Mississauga, ON, Canada). Complementary DNA (cDNA) was then synthesized using 10 µl of the RNA preparation, 0.75 µM of random hexamer primers (Amersham Pharmacia Biotech, Baie d'Urfé, QC, Canada), 300 copies of an internal control
10 template, and the Omniscript Reverse Transcriptase Kit (QIAGEN, Mississauga, ON, Canada) following the manufacturer's instructions. The internal control template consisted of a 554 or 558 bp (depending on the multiplex assay) transcribed region of an herpes simplex virus type 2 DNA polymerase region flanked by either influenza B or rhinovirus/enterovirus complementary primer sequences (depending on the
15 multiplex assay) cloned in the pDrive plasmid (QIAGEN).

For DNA amplification by the widely used PCR method, primer pairs were mostly derived from public database sequences (Table 10). During DNA amplification by PCR, two oligonucleotide primers binding respectively to each strand of the heat-
20 denatured target DNA from the viral genome are used to amplify exponentially *in vitro* the target DNA by successive thermal cycles allowing denaturation of the DNA, annealing of the primers and synthesis of new targets at each cycle (Persing, 1993, Diagnosis Molecular Microbiology: Principles and Applications, American Society for Microbiology, Washington, D.C., 88-109).

25

In a preferred embodiment, cDNA was amplified using a real-time PCR procedure with the LC Faststart DNA Master SYBR Green 1 Kit (Roche Diagnostics, Laval, QC, Canada) or the SYBR Green TAQ ReadyMix for quantitative PCR capillary formulation (Sigma-Aldrich, Oakville, ON) in a LightCycler instrument (Roche
30 Diagnostics). Each reaction had a total volume of 20 µl including 2 µl of cDNA and 18 µl of a reaction mixture containing 4 mM MgCl₂, 2 µl of LC Faststart DNA SYBR

Green 1 Master Mix or the SYBR Green TAQ ReadyMix for quantitative PCR capillary formulation, 3% DMSO, and 0.3-1.0 μ M of each viral primer (see Tables 10, 12, and 13 for primer sequence ID NOs., length of amplicons, and specific combination of primers, respectively). Cycling conditions typically included an initial
5 denaturation step of 10 min at 94°C, followed by 50 cycles of 15 s at 94°C, 5 s at 58°C, and 25 s at 72°C. However, the cycling conditions could vary slightly according to the multiplex PCR assay.

In a preferred embodiment, two real-time multiplex PCR assays were used to amplify
10 and detect the panel of common respiratory viruses (Table 13). The respiratory-1 multiplex assay combines 4 sets of primers for amplification of influenza A and B viruses, hRSV, hMPV, and an internal control template. The respiratory-2 multiplex assay comprises 4 other sets of primers for amplification of PIV-1, PIV-3, adenoviruses, rhinoviruses/enteroviruses, and an internal control template. The use of
15 any other combinations of the same or similar primer sequences in two or more than two multiplex assays for detection of common respiratory viral pathogens is also under the scope of this invention.

The person skilled in the art of nucleic acid amplification knows the existence of other
20 rapid amplification procedures such as ligase chain reaction (LCR), reverse transcriptase PCR (RT-PCR), solid phase PCR, transcription-mediated amplification (TMA), self-sustained sequence replication (3SR), nucleic acid sequence-based amplification (NASBA), strand displacement amplification (SDA), branched DNA (bDNA), cycling probe technology (CPT), solid phase amplification (SPA), rolling
25 circle amplification technology (RCA), solid phase RCA, anchored SDA, and nuclease dependent signal amplification (NDSA) (Persing, 1993, Diagnosis Molecular Microbiology: Principles and Applications, American Society for Microbiology, Washington, D.C., 88-109; Lee, 1997, Nucleic Acid Amplification Technologies: Application to Disease Diagnosis, Eaton Publishing, Boston, MA., 61-125; Westin et
30 al., 2000, Nat. Biotechnol.18:199-204). The scope of this invention is not limited to the use of amplification by PCR, but rather includes the use of any nucleic acid

amplification method or any other procedure which may be used to increase the sensitivity and/or the rapidity of nucleic acid-based diagnostic tests. The scope of the present invention also covers the use of any nucleic acids amplification and detection technology including real-time or post-amplification detection technologies, any
5 amplification technology combined with detection, any hybridization nucleic acid chips or array technologies, any amplification chips or combination of amplification and hybridization chip technologies. Detection and identification by any nucleotide sequencing method is also under the scope of the present invention.

10 Detection of amplification products

Classically, the detection of PCR amplification products is performed by standard ethidium bromide-stained agarose gel electrophoresis. Using this approach, the amplicons may be discriminated based on their size. In the present invention, the
15 length of the different amplicons varies from 84 to 929 bp (Table 12). It is however clear that other detection methods of amplified products may be faster, more sensitive, and more practical. Amplicon detection may also be performed by solid support or liquid hybridization using strain-specific, internal DNA probes hybridizing to an amplification product. Such probes may be generated from sequences designed to
20 specifically hybridize to DNA amplification products which are objects of the present invention. Amplicons can be also characterized by DNA sequencing. Alternatively, a rapid detection method developed with real-time PCR assays consist of determining the specific melting temperature of amplicons at the end of the amplification reaction.

25 The detection of viral amplicons was first characterized using the melting curve analysis feature of a real-time PCR instrument. Using this procedure, the different viral targets amplified in each of the multiplex PCR assays could be differentiated by their specific melting temperature (Table 13 and Fig. 7 and 8). Alternatively, specific fluorescent probes (Table 11) for each viral target can be used for real-time detection
30 and quantification of amplicons in a single amplification/detection assay. Real-time quantification of viral products can be performed through a variety of fluorescence-

based technologies such as adjacent probes, Taqman probes, and molecular beacons. Alternatively, these probes may be used in a separate post-amplification step.

5 The diagnostic kits, primers and probes mentioned above can be used to detect and/or identify respiratory viruses, whether said diagnostic kits, primers and/or probes are used for *in vitro* or *in situ* applications. The said samples may include but are not limited to: any clinical sample, any environmental sample, any viral culture, any tissue, and any cell line.

10 It is also an object of the present invention that said diagnostic kits, primers and/or probes can be used alone or in combination with any other assay suitable to detect and/or identify and /or quantify microorganisms, including but not limited to: any assay based on nucleic acids detection, any immunoassay, any enzymatic assay, any biochemical assay, any serological assay, any culture on specific cell lines, and any
15 infectivity assay on animals.

In the methods and kits described herein, the oligonucleotide probes and amplification primers have been derived from larger sequences (i.e. DNA fragments of at least 100 base pairs). It is clear to the individual skilled in the art that oligonucleotide sequences
20 other than those described in the present invention and which are appropriate for detection and/or identification of any respiratory viruses may also be derived from those selected in the present invention. For example, the oligonucleotide primers or probes may be shorter but of a length of at least 10 nucleotides or longer than the ones chosen; they may also be variants of the same oligonucleotide. If the target DNA or a
25 variant thereof hybridizes to a given oligonucleotide, or if the target DNA or a variant thereof can be amplified by a given oligonucleotide PCR primer pair, the converse is also true; a given target DNA may hybridize to a variant oligonucleotide probe or be amplified by variant oligonucleotide PCR primers. Alternatively, the oligonucleotides may be designed for use in amplification methods other than PCR. Consequently, one
30 aspect of the present invention is the simultaneous detection and/or identification of various respiratory viruses by targeting nucleic acid sequences which are used as a

source of specific and ubiquitous oligonucleotide probes and/or amplification primers. Although the selection and evaluation of oligonucleotides suitable for diagnostic purposes require much effort, it is quite possible for the individual skilled in the art to derive, from the selected oligonucleotides (listed in Tables 10 and 11), other
5 oligonucleotides which are suitable for diagnostic purposes.

Thus, the specific primers and/or probes selected for amplification and detection of respiratory viruses and their specific combination in multiplex assays are objects of this invention (Tables 10-13).

10

EXAMPLES:

Example 1: Evaluation of different primers for hMPV detection

15

An evaluation of the sensitivity of different hMPV primer sets in a real-time PCR assay was performed using 20 positive hMPV cultures grown in LLC-MK2 cells and 10 naso-pharyngeal aspirate specimens. These samples were collected in the Québec City area (QC, Canada) over a period of 3 years (2000-2002). Viral RNA was
20 prealably extracted from 200 µl of viral culture supernatants or naso-pharyngeal aspirates using the QIAamp Viral RNA Mini Kit (QIAGEN). Complementary DNA was synthesized using 10 µl of eluted RNA, 0.75 µM of a specific hMPV primer (Table 2, sequences ID Nos. 91, 97, 100, 102, 112), and the Omniscript Reverse Transcriptase Kit (QIAGEN, Mississauga, ON, Canada) following the manufacturer's
25 recommendations. Complementary DNA was amplified using a real-time PCR procedure with the LC Faststart DNA Master SYBR Green 1 Kit (Roche Diagnostics, Laval, QC, Canada) in a LightCycler instrument (Roche Diagnostics). Each reaction had a total volume of 20 µl including 2 µl of RT mixture and 18 µl of a reaction mixture containing 4 mM MgCl₂, 2 µl of Faststart DNA SYBR Green Master Mix,
30 3% DMSO, and 0.5 µM of each hMPV primer (Table 4: sequence ID Nos.: 92 and 94; 100 and 101; 97 and 98; 102 and 103; and 112 and 113, for the nucleoprotein (N),

30

matrix (M), fusion (F), phosphoprotein (P), and polymerase (L) genes respectively). Cycling conditions typically included an initial denaturation step of 10 min. at 94°C, followed by 50 cycles of 15 s at 94°C, 5 s at 54°C, and 30 s at 72°C. However, the cycling conditions could slightly vary according to the hMPV gene to amplify.

- 5 Detection of hMPV amplicons was characterized using the melting curve analysis feature of a real-time PCR instrument. Briefly, following the last amplification cycle on a LightCycler instrument, the internal temperature was rapidly increased to 94°C then decreased to 60°C for 30 seconds, followed by slowly increasing to 94°C at a rate of 0.1°C per second, with continuous fluorescence lecture.

10

Using this procedure, the specific melting temperature obtained with the hMPV amplicons produced by the primers specific for target genes coding for N, M, F, P and L proteins was found to be 82.63°C ± 0.87°C, 82.75°C ± 0.80°C, 83.29°C ± 0.45°C, 80.48°C ± 0.46°C, and 77.99 ± 0.33°C, respectively. As shown in Table 5, a real-time

- 15 PCR assay using primers complementary to the N gene successfully amplified RNA from all 20 cultures and 10 clinical specimens. In contrast, hMPV primers specific for the M, F, P, and L genes had sensitivities of only 70%, 63.3%, 50%, and 89.7% respectively. Thus, primers aimed at amplifying the hMPV N gene appear to target more conserved regions of the hMPV genome and, consequently, were selected for
20 further detailed evaluation (see example 2).

Example 2: Validation of a real-time PCR assay for the hMPV nucleoprotein (N) gene in a pediatric population

- 25 An hMPV N plasmid was constructed by subcloning the amplified hMPV N region in the plasmid pDrive (QIAGEN, Mississauga, ON, Canada). The new plasmid was transcribed using the RNA Transcription Kit (Stratagene, Vancouver, BC, Canada) for sensitivity analysis. Using the previously described RT procedure (N primer, sequence ID No. 91) and real-time PCR protocol (N primers, sequence ID Nos. 92
30 and 94) reported in example 1, the sensitivity of the assay was estimated at 100 copies per reaction with a specific melting temperature of 82 °C. The assay was found to be

specific with no amplification signal observed when viral cultures positive for the human respiratory syncytial virus, the parainfluenza viruses, the influenza viruses A and B, and the adenoviruses were tested. The real-time PCR assay for the hMPV N gene was subsequently validated in a prospective case-control study in children below
5 the age of 3 years hospitalized at Laval University Hospital Center (CHUQ-CHUL) in Québec City, Canada, during the winter season of 2001-02. Cases consisted of children hospitalized for a respiratory syndrome whereas controls consisted of children hospitalized during the same period for a surgery with no respiratory symptoms. At admission, a nasopharyngeal aspirate was collected for viral antigen
10 tests, viral cultures, and RT-PCR studies for influenza A and B viruses, the human respiratory syncytial virus (hRSV), and the hMPV N gene.

No virus was detected in any of the samples from the 51 control children. Among the 208 cases tested by real-time PCR for hMPV, hRSV, influenza A and B viruses, the
15 positivity rates were 5.8%, 51.0%, 21.6%, and 0%, respectively. Viral culture was positive for hMPV in only 2 cases whereas the real-time hMPV PCR assay for the N gene was positive for 12 cases including the 2 positive by viral culture. All positive hMPV PCR results for the N gene were confirmed by a second PCR assay targeting the hMPV fusion (F) gene and by the use of DNA sequencing which indicated the
20 presence of multiple hMPV strains clustering in the two previously-reported F genotypes. Thus, the real-time PCR assay described in this invention appears to be much more sensitive than conventional viral culture for the detection of hMPV in clinical samples.

Example 3: Typing of hMPV clinical strains using restriction enzyme analysis

We designed a rapid molecular typing method for hMPV epidemiological studies based on amplification of a portion of the fusion (F) gene which is then digested with specific restriction enzymes. Briefly, a 759-bp fragment of the hMPV F gene was
5 amplified by a real-time PCR procedure as described in example 1 using specific primers (sequence ID Nos. 97 and 98). After PCR amplification, two aliquots (10 uL) of the PCR product were digested during 2 hours at 37°C with 10 units of ApaI, I and 4 units of Avr II (New England Biolabs, Mississauga, ON, Canada) according to the manufacturer's recommendations. Digested products were then loaded on an 1%
10 agarose gel for electrophoresis.

Table 6 summarizes the size of the F fragments obtained for different hMPV isolates following enzymatic digestion and the corresponding genotype as defined by DNA sequencing and phylogenetic studies. In brief, all hMPV isolates belonging to the
15 genotype F-I were digested by ApaI, I (fragments of 532 and 227 bp) but not by Avr II whereas those belonging to the genotype F-II were digested by Avr II (fragments of 447 and 312 bp) but not by ApaI, I (see Figure 2 for an example of the band pattern following electrophoresis on an agarose gel).

20 **Example 4 : A prospective case-control study for defining the role of respiratory viruses in children hospitalized for acute respiratory tract infections**

Summary

The human metapneumovirus (hMPV) is a new paramyxovirus recently isolated from
25 young children with respiratory syndromes. The objectives of this study were to estimate the relative contribution of hMPV in children's hospitalization for acute respiratory tract infections (ARTI) and to define its clinical presentation and seasonal pattern.

We evaluated the percentage of hospitalizations for ARTI in children aged 0-3 years attributable to hMPV and other respiratory viruses in a prospective case-control study during the winter/spring seasons of 2002. We used multiplex real-time PCR assays and other conventional diagnostic methods to detect hMPV, the human respiratory syncytial virus (hRSV), and the influenza viruses in nasopharyngeal aspirates of children.

HMPV was detected in 12 (5.8%) of the 208 children hospitalized for ARTI as compared to 118 (56.7%) for hRSV and 49 (23.6%) for influenza A. In contrast, none of the 51 controls harbored any of the respiratory viruses. Most hRSV and influenza infections occurred in January and February whereas the peak hMPV activity was in March and April. The peak age for hospitalizations for hRSV and hMPV infections was < 3 months and 3-5 months, respectively. Diagnoses of bronchiolitis and pneumonitis were made in respectively 67% and 17% of hMPV-infected children compared to 84% and 25% for those with hRSV infection. None of the hMPV-infected children was admitted to the intensive care unit compared to 15% for those with hRSV or influenza A infections.

HMPV causes significant morbidity in young children and its clinical presentation is similar although less severe than that of hRSV.

Introduction

The human metapneumovirus (hMPV) is the first human member of the new *Metapneumovirus* genus within the *Paramyxoviridae* family (van den Hoogen BG, de Jong JC, Groen J, et al. Nat Med 2001;7:719-24; van den Hoogen BG et al. Virology 2002;295:119-32.) The human respiratory syncytial virus (hRSV) belongs to a separate genus within the same family (Taxonomy V. Seventh Report of the International Committee on Taxonomy of Viruses. Academic, San Diego, 2000.).

HMPV has been recently identified in nasopharyngeal aspirates of children and adults suffering from a range of respiratory tract infections in various parts of the world (van

den Hoogen BG, de Jong JC, Groen J, et al. Nat Med 2001;7:719-24; Nissen MD, et al. Med J Aust 2002;176:188; Peret TC, Boivin G, Li Y, et al. J Infect Dis 2002;185:1660-3; Stockton J, et al. Emerg Infect Dis 2002;8:897-901). The clinical syndrome of the infected children encompasses mild respiratory problems to
5 bronchiolitis and pneumonitis (van den Hoogen BG, de Jong JC, Groen J, et al. Nat Med 2001;7:719-24; Pelletier G, et al. Emerg Infect Dis 2002;8:976-8; Boivin G, Abed Y, Pelletier G, et al. J Infect Dis 2002;186:1330-4).

Using reverse-transcription polymerase chain reaction (RT-PCR), the rate of hMPV
10 detection in nasopharyngeal aspirates (NPA) of pediatric patients with unexplained ARTI varied between 1.5 and 10% (van den Hoogen BG, de Jong JC, Groen J, et al. Nat Med 2001;7:719-24; Nissen MD, et al. Med J Aust 2002;176:188). These retrospective studies are small, excluded patients who tested positive for other viruses, only superficially describe the clinical features of the disease and lack
15 morbidity/mortality data. In absence of a control group, these studies cannot differentiate if hMPV is a colonizing or a pathogenic virus. More recently, Stockton et al. identified hMPV RNA in 2.2% of 405 specimens from patients with influenza-like illnesses who consulted general practitioners in England although few swabs were
20 collected from children < 5 years (Stockton J, et al. Emerg Infect Dis 2002;8:897-901).

The objectives of this study were to estimate the relative contribution of hMPV in children's hospitalization for acute respiratory tract infections (ARTI) and to define
its clinical presentation and seasonal pattern.

Materials and Methods

Study design. Participants were children aged ≤ 3 years hospitalized between December 15, 2001 and April 20, 2002 at Laval University Hospital Center (CHUQ-CHUL) in Québec City, Canada. Cases were admitted for an ARTI (bronchiolitis, pneumonia, laryngotracheobronchitis, etc...) whereas controls were hospitalized for an elective surgery and had no respiratory symptoms or fever. Children hospitalized twice were counted as two cases. At admission, after signed consent was obtained from their parents, all children had an NPA (1-2 ml) collected and tested for hRSV antigen (TestPack, Abbott Laboratories, Abbott Park, IL). The specimen was frozen at -80°C until subsequent RT-PCR studies. Additionally, viral cultures, direct immunofluorescence staining, and microbiological testing were performed according to the treating physician. A standardized questionnaire was completed at admission by a single research nurse with the parents. At the end of the hospitalization, the children's charts were reviewed to collect clinical and laboratory data. The study was approved by the CHUQ-CHUL research ethics board.

RNA extraction and cDNA synthesis. Viral RNA was extracted from 200 µl of NPA specimens using the QIAamp viral RNA Mini Kit (QIAGEN, Mississauga, ON, Canada). Complementary cDNA was synthesized using 10 µl of eluted RNA and the Omniscript Reverse Transcriptase kit (QIAGEN). Random hexamer primers (Amersham Pharmacia Biotech, Baie d'Urfé, QC, Canada) were used in the RT step of the multiplex respiratory PCR assay whereas a specific primer (5'-TGGGACAAGTGAAAATGTC-3') serves to synthesize hMPV cDNA in a separate PCR assay. An internal control (300 copies) consisting of a 534-bp transcribed region of the pDrive plasmid (QIAGEN) flanked by influenza B complementary primer sequences (see below) was spiked in the RT reaction of the multiplex respiratory PCR assay to verify the presence of PCR inhibitors.

RT-PCR assays and phylogenetic studies. The multiplex respiratory PCR assay was designed to amplify conserved regions of the influenza A (SEQ. ID Nos. 139-140) (Fouchier RA, et al. J Clin Microbiol 2000;38:4096-101) and B (SEQ. ID Nos. 141-

142) (Li J, Chen S, Evans DH. J Clin Microbiol 2001;39:696-704) matrix genes as well as the fusion gene of hRSV (SEQ. ID Nos. 143-144) (Mazzulli T, Peret TC, McGeer A, et al. J Infect Dis 1999;180:1686-9).

- 5 Complementary DNA was amplified using a real-time PCR procedure with the LC Faststart DNA Master SYBR Green 1 Kit (Roche Diagnostics, Laval, Québec, Canada) in a LightCycler instrument (Roche Diagnostics). Each reaction had a total volume of 20 µl including 2 µl of cDNA and 18 µl of a reaction mixture containing 2.4 mM MgCl₂, 2 µl of Faststart DNA SYBR Green Master Mix, 3% DMSO, 0.3 mM
- 10 each of influenza A and B primers, and 0.8 mM of hRSV primers. The thermal cycling incubations consisted of an initial denaturation step at 94°C for 10 min, followed by 50 cycles of 15 s at 94°, 5 s at 58°C, and 25 s at 72°C. At the end of each cycle, the fluorescent signal was measured at a wavelength of 530 nm using the LightCycler Fluorimeter. The melting curve analysis program of the LightCycler was
- 15 used to identify specific PCR products of influenza A (T_m: 85°C), influenza B (T_m: 83°C), hRSV (T_m: 79°C), and the internal control (T_m: 92°C). The sensitivity of the multiplex respiratory PCR assay was estimated at 50 copies for each viral target when mixed with 300 copies of the internal control.
- 20 A specific PCR assay was designed for amplification of the hMPV Nucleoprotein (N) gene in the LightCycler instrument. The sequences of the forward (MPV-NCS) and reverse (MPV-NCA) primers were respectively 5'-GAGTCTCAGTACACAATTAA-3' (SEQ ID NO 92) and 5'-GCATTTCCGAGAACAACAC-3' (SEQ ID NO 94). The specific cDNA was amplified using conditions similar to those of the multiplex
- 25 respiratory PCR assay except that 0.5 mM of each primer was added to the master mix. Cycling conditions included an initial denaturation step of 10 min at 94°C, followed by 50 cycles of 15 s at 94°C, 5 s at 54°C, and 30 s at 72°C. The melting temperature of the amplified hMPV N fragment was 82°C and the sensitivity of the PCR assay was 100 copies. For phylogenetic studies, nucleotide sequences were
- 30 determined from amplified hMPV F (fusion) gene products then analyzed using the

neighbor joining algorithm and Kimura-2 parameters. (Boivin G, Abed Y, Pelletier G, et al. J Infect Dis 2002;186:1330-4).

hMPV in the general population. To further assess the seasonal distribution, affected age groups and frequency of hMPV, we compared data from the study to those of the general population derived from the viral cultures found positive for hMPV in our regional virology diagnostic laboratory, the only one performing viral cultures for the Québec City area (population approximately 600 000). Respiratory specimens were inoculated in 96-well plates containing ten cell lines (MDCK, LLC-MK2, Hep-2, human foreskin fibroblast, Vero, mink lung, A-549, rhabdomyosarcoma, 293, and HT-29) then incubated for a period of 21 days. A positive cytopathic effect was confirmed by immunofluorescence testing using monoclonal antibodies or by RT-PCR (hMPV).

Statistical analyses. Proportions were compared by the χ^2 test.

Results

The study population included a total of 208 hospitalized cases with ARTI (including 8 children who were admitted twice) and 51 children who served as controls.

20

Description of the study population and viral etiologies. The age distribution of children with ARTI is presented in Fig. 3: 56% (116) were male and 44% (92) were female. Hospitalization was the highest in infants ≤ 3 months of age and steadily decreased thereafter. Most (90%) children had no underlying medical conditions at admission.

25

All 208 cases had an NPA. The mean delay between the onset of symptoms and collection of NPA was 6 days (median 4 days). This delay did not differ for the different viruses detected. All 208 cases were tested by PCR, 204 had an hRSV

antigen detection test, 100 had a direct examination procedure for other viruses, and 100 had a viral culture. At least one respiratory virus was detected in 164 (79%) cases whereas none was detected in 44. Twelve children (5.8%) were positive for hMPV, 118 (56.7%) for hRSV, 49 (23.6%) for influenza A, and none for influenza B (Table 7). There was no PCR testing for adenoviruses and PIV type 2 but these two viruses were found in 6% and 2%, respectively, of the 100 viral cultures or antigenic assays. Single virus infections occurred in 141 (86%) of the 164 positive cases and mixed infection was present in 23 (14%). Two of the 12 hMPV infections were mixed (hMPV-influenza and hMPV-hRSV). The other combinations were hRSV-influenza A (18 cases), hRSV-adenovirus (2), and influenza A-adenovirus (1).

Among the 208 cases tested by PCR for hMPV, hRSV, influenza A and B viruses, the positivity rates were 5.8%, 51.0%, 21.6%, and 0%, respectively (Table 7). In addition, 16 other cases had one of the four latter respiratory viruses identified only by culture (3 influenza A and one hRSV), only by an antigen detection test (10 hRSV and one influenza A) or by both culture and antigen detection test (1 hRSV). Among the 8 children hospitalized twice, none had the same viral infection at both admissions. The combinations observed were hMPV - hRSV (2), hMPV - no virus (1), hRSV - no virus (2), hRSV - influenza (2), no virus - no virus (1).

The biweekly distribution of cases with respiratory tract viruses is shown in Fig. 4. hRSV and influenza A infections occurred predominantly between January and March, whereas hMPV infections occurred mostly in March and April. The proportion of children with virologically-confirmed respiratory tract infection decreased after February.

Clinical features of cases. The distribution of the age distribution by different viral infections is shown in Fig. 3. While the peak age for hMPV infection was 3-5 months, it was before 3 months for hRSV infection. For influenza A, it spread evenly during the first year of life. The peak age for mixed infection was 6-11 months and decreased thereafter. Gender was distributed evenly with each virus group but more males

(70%) had mixed viral infections. The majority (75% with hMPV, 93% with hRSV, 90% with influenza A virus infection) of the children in different etiological groups had no medical underlying conditions. Three (25%) children with hMPV infection had a cardiac disorder, one having concomitant cardiac, pulmonary, and neurological problems.

Signs and symptoms recorded with the different respiratory viruses were generally similar except for the vomiting which occurs more frequently with hMPV infections ($p = 0.05$) (Table 8). The mean and median duration of hospitalization was similar for hMPV, hRSV and influenza A viruses being respectively 5.5 and 4.5, 5.8 and 5.0, and 4.9 and 4.0 days. Of note, one third of hMPV-infected children were hospitalized more than 7 days. None of the children with hMPV infection was admitted to the intensive care unit (ICU) contrasting with 15% and 16% of children with hRSV and influenza A, respectively. There was no death as part of this study. The duration of the hospitalization in children with no detectable virus was shorter than that in children with single or mixed infection (Kruskal-Wallis Test, $p < .0001$). Two-thirds of the children were given antibiotics during their hospitalization, independently of the etiology of their respiratory disease.

At hospital discharge, a final diagnosis of bronchiolitis was given to 67% of children with hMPV, 84% with hRSV, and 51% with influenza A ($p < 0.001$) (Table 9). Otitis occurred in about half of children with hMPV, hRSV, and influenza A infections. Pneumonitis was less frequently diagnosed in children with hMPV compared to those with hRSV or influenza A (17%, 25%, and 37%, $p = 0.22$). Final diagnoses were similar with single and mixed infections.

Controls. There were 51 children in the control group, 29 (56.9%) males and 22 (43.1%) females. The age distribution of the controls was similar to that of the cases (Fig. 3) as was the period of hospital admission (Fig. 4). None of the controls had a virus detected by PCR.

hMPV in the general population. The regional virology laboratory received 1505 respiratory specimens for viral culture between January 1 and June 30, 2002. In total, 36 (including two study participants) or 2.9% were positive for hMPV: 24 (66.6%) in children < 2 years of age, 5 in those 2 to 4 years, 4 in adults 30-49 years, and 3 in those 70 years or older. Most cases (81%) were isolated over a two-month period (from March 23 to May 18) (Fig. 5). When comparing the seasonal distribution of hMPV in hospitalized children (study population) and in the general population, it appears that the study in hospitalized children did not cover the entire hMPV season as it was stopped just after the peak time of hMPV transmission (April 6-20).

10

Phylogenetic analyses of hMPV strains. The 12 hMPV strains isolated in the prospective study clearly clustered into two F lineages with nine strains belonging to group 1 and three to group 2 (Fig. 6). Seven of the 9 group 1 strains had identical F gene sequences. At the nucleotide level, there was 84-85% similarity between groups versus 98-100% (group 1) and 93-99% (group 2) similarity within groups. There were fewer differences at the amino acid level i.e. 94-96% similarity between groups versus 99-100% (group 1) and 98-99% (group 2) among strains within the same group. Although there was a trend for more recent strains to cluster in group 2, the correlation between genotypes and date of hospitalization was not perfect as two strains detected on the same day (No. 216 and 217) belonged to two different F lineages.

20

Discussion

Our prospective study has revealed important clinical and epidemiological features related to hMPV. First, our data indicate that hMPV is really a respiratory pathogen. Second, we found that it contributes to ARTI leading to children's hospitalization in proportion similar to adenoviruses or PIV but significantly less than hRSV and influenza viruses. Third, the symptoms and complications associated with hMPV were indistinguishable from those caused by hRSV. Finally, our results suggest that the epidemiology of hMPV infection in children may differ from that of hRSV and

25

30

influenza viruses both in terms of the age of first acquisition and in the timing of the seasonal epidemics.

Recent studies by our group (Pelletier G, et al. *Emerg Infect Dis* 2002;8:976-8; Boivin
5 G, Abed Y, Pelletier G, et al. *J Infect Dis* 2002;186:1330-4) and others (van den
Hoogen BG, de Jong JC, Groen J, et al *Nat Med* 2001;7:719-24.; Nissen MD et al.
Med J Aust 2002;176:188; Stockton J, et al. *Emerg Infect Dis* 2002;8:897-901) had
suggested that hMPV should be added to the list of human respiratory viral pathogens
(Dowell SF, Anderson LJ, Gary HE, Jr., et al. *J Infect Dis* 1996;174:456-62; Treanor
10 JJ. *Respiratory Infections*. In: Richman DD, Whitley, R.J., Hayden, F.G., ed. *Clinical
Virology*. Vol. 1. Washington: ASM Press, 2002:7-26).

The absence of hMPV in respiratory tract of our 51 control children, the absence of
other respiratory viruses in 83% of the 12 hMPV-infected children and the severity of
15 their symptoms (bronchiolitis and/or pneumonitis) provide further evidence that
hMPV is really a pathogenic respiratory virus. The absence of underlying medical
conditions in 75% of the hMPV-infected children further demonstrates its
pathogenicity. The use of PCR assays was particularly advantageous for hMPV
because its growth in culture is fastidious and limited to a few permissive cell lines
20 (Boivin G, Abed Y, Pelletier G, et al. *J Infect Dis* 2002;186:1330-4) and because
rapid antigenic detection tests are currently unavailable.

Our study provides for the first an estimate of the proportion of ARTI hospitalizations
attributable to hMPV in a pediatric population. During the period ranging from
25 December 15 to April 20 2002, hMPV accounted for 5.8% of all hospitalizations for
respiratory tract infections in children aged 0-3 years. This underestimates the real
proportion as our hospitalization study stopped before the end of the hMPV season
prevailing in the community (Fig. 5). Although we did not perform PCR testing for
adenoviruses and PIV, the percentage of hospitalizations caused by hMPV seems
30 similar to that attributable to these two viruses and much smaller than that of hRSV or
influenza A.

One of the most important conclusion of our study is that hMPV disease cannot be distinguished on clinical findings from that caused by hRSV and influenza A (Tables 8 and 9) but tended to be somewhat less severe with fewer pneumonia and no admission in the ICU. Nevertheless, hMPV infection was associated with a significant clinical and economical burden as shown by a mean hospital stay of 5.5 days and one-third of hMPV-infected cases being hospitalized for > 7 days.

A small serological study from the Netherlands showed that all children over the age of 5 years had hMPV antibodies suggesting a high level of transmission.¹ Data from our study suggest that morbidity caused by hMPV is greatest in children < 2 years of age as they represented 83% (10/12) of our hospitalized cases and two-thirds of the hMPV isolates recovered in our diagnostic virology laboratory. This suggests that similar to other paramyxoviruses such as hRSV, the morbidity of hMPV in children occurs through primo-infection. In contrast to hRSV which peaked during the first three months of life, hMPV infections peaked at a slightly older age i.e. between the third and fifth month of life. Whether this difference depends on a longer persistence of maternal antibodies or a less efficient transmission mode in the case of hMPV would require additional studies.

During the 4-week period from mid-March to mid-April, hMPV infections clustered (11/12 detected) and accounted for 18.9% of all hospitalizations for ARTI in children. This contrasts with hRSV and influenza A which occurred mostly in January and February (Figure 4). Based on passive surveillance data from our regional virology laboratory, the peak time of hMPV transmission in the community occurred between April 6-20, 2002, and continued beyond the end of our study in hospitalized children (Fig. 5). Year-long active surveillance studies will be required to better assess the seasonal patterns of hMPV transmission.

As described for hRSV (Mazzulli T, Peret TC, McGeer A, et al J Infect Dis 1999;180:1686-9), several strains of hMPV circulated during a very short period of

time (one month) in our study area. The hMPV strains segregated into two F subgroups in agreement with previous studies although one strain clearly predominated accounting for 58.3% of all isolates (van den Hoogen BG, de Jong JC, Groen J, et al. Nat Med 2001;7:719-24; Peret TC, Boivin G, Li Y, et al. J Infect Dis 5 2002;185:1660-3; Boivin G, Abed Y, Pelletier G, et al. J Infect Dis 2002;186:1330-4).

Due to the small number of hMPV strains belonging to one of the F subgroup, we did not attempt to correlate hMPV genotype with clinical outcome. Such viral heterogeneity may allow multiple reinfections throughout life especially in elderly 10 subjects and immunocompromised patients (Pelletier G, Dery P, Abed Y, Boivin G Emerg Infect Dis 2002;8:976-8).

In conclusion, our study supports the concept of the epidemic nature of hMPV infection and establishes its role as a significant pathogen in severe ARTI of children.

15

Example 5: Determination of the hMPV Load in Clinical Samples Using Quantitative Real-time PCR

20 An evaluation of a quantitative real-time PCR hMPV assay using a TaqMan probe was performed using naso-pharyngeal aspirate specimens collected from children and adults in Québec City during the winter seasons of 2001-02 and 2002-03. Viral RNA was first extracted from 200 µl of naso-pharyngeal aspirates using the QIAamp Viral RNA Mini Kit (QIAGEN) and eluted in a volume of 40 µl. Serial concentrations 25 (5X10² to 1X10⁹ copies) of a transcribed hMPV plasmid were used to construct the standard curve of the assay. Complementary DNA was synthesized using 10 µl of eluted RNA from clinical samples or hMPV plasmid dilutions, 50 ng (0.75 µM) of random hexamer primers (Amersham Pharmacia Biotech), 0.5 mM of dNTPs, 1X Omniscript buffer and 4 units of the Omniscript reverse transcriptase (QIAGEN) in a 30 final volume of 20 µl. The reverse transcription (RT) mixture was incubated during 1.5 hr. at 37°C and inactivated at 70°C during 10 min. Complementary DNA was amplified by real-time PCR with a TaqMan probe for the hMPV L gene and the LC

44

Master Hybridization Probe Mix (Roche Diagnostics) in a LightCycler instrument (Roche Diagnostics). Each reaction had a volume of 20 µl including 2 µl of RT mixture, 4 mM MgCl₂, 0.3 µM of specific hMPV L primers (sequence ID Nos. 112 and 114), 0.6 µM of a specific hMPV L probe (SEQ. ID. No. 127), and 2 µl of the LC Master Hybridization Probe Mix (Roche Diagnostics). Cycling conditions included an initial denaturation step of 30 s at 94°C, followed by 60 cycles of 5 s at 94°C, 5 s at 56°C, and 25 s at 72°C. A fluorescence lecture was taken after each elongation step (Fig. 9A). A standard curve for the PCR reaction was constructed by plotting cycle threshold values against log concentrations of each hMPV plasmid dilution. The number of hMPV copies present in clinical samples was calculated by interpolating cycle threshold values into the assay's standard curve (Fig. 9B).

Example 6: Determination of hMPV genotypes based on amplification and sequencing of the viral F and G genes.

Determination of hMPV genotypes was achieved by PCR amplification of the F or G genes followed by DNA sequencing of the viral amplified products. Different sets of primers were used for PCR amplification and DNA sequencing of the F (SEQ. ID. Nos. 97 and 98) and G (SEQ. ID. Nos. 104 and 107 or 108 and 109) genes. Alignment of viral DNA sequences was done with the CLUSTAL W software version 1.7 for Unix and phylogenetic trees were computed by maximum parsimony, distance and maximum likelihood-based criteria analysis using PAUP version 4.0.d8. Phylogenetic trees for both the F (Fig. 10A) and G (Fig. 10B) genes from Canadian and non-Canadian viral isolates revealed the existence of two hMPV groups (genotypes A and B) with two subgroups (subtypes 1 and 2) within each group.

Example 7: Comparison of viral culture and multiplex PCR assay for detection of influenza A virus.

A multiplex PCR assay (including PCR primers defined in SEQ. ID. Nos. 139, 140, 141, 142, 143, and 144) aimed at detecting influenza A and B viruses and the respiratory syncytial virus was used to assess the etiology of severe acute respiratory

tract infections (ARTI) necessitating hospitalization in children aged 0-3 years during the winter season 2001-02 at the CHUL, an university-based hospital, in Québec City (Canada). Nasopharyngeal (NPA) samples from all 208 children with ARTI were tested by multiplex PCR whereas a subset of 100 NPA specimens was inoculated on

5 Madin Darby Canine Kidney (MDCK) cells for influenza culture. Briefly, viral RNA was preably extracted from 200 µl of naso-pharyngeal aspirates using the QIAamp Viral RNA Mini kit (QIAGEN, Mississauga, ON, Canada). cDNA was then synthesized using 10 µl of the RNA preparation, 0.75 µM of random hexamer primers (Amersham Pharmacia Biotech, Baie d'Urfé, QC, Canada), 300 copies of an internal

10 control template, and the Omniscript Reverse Transcriptase Kit (QIAGEN, Mississauga, ON, Canada) following the manufacturer's instructions. The internal control template consisted of a 558-bp transcribed region of an herpes simplex virus type 2 DNA polymerase region flanked by influenza B complementary primer sequences cloned into the pDrive plasmid (QIAGEN). cDNA was amplified using a

15 real-time PCR procedure with the LC Faststart DNA Master SYBR Green 1 Kit (Roche Diagnostics, Laval, QC, Canada) in a LightCycler instrument (Roche Diagnostics). Each reaction had a total volume of 20 µl including 2 µl of complementary DNA and 18 µl of a reaction mixture containing 4 mM MgCl₂, 2 µl of LC Faststart DNA SYBR Green Master 1 Mix, 3% DMSO, 0.3 µM of influenza A

20 and influenza B primers, and 0.8 µM of human respiratory syncytial virus primers (Table 10). Cycling conditions included an initial denaturation step of 10 min. at 94°C, followed by 50 cycles of 15 s at 94°C, 5 s at 58°C, and 25 s at 72°C. Detection of viral amplicons was characterized using the melting curve analysis feature of a real-time PCR instrument. Briefly, following the last amplification cycle on a

25 LightCycler, the reaction temperature was rapidly increased to 94°C and was then decreased to 60°C for 30 seconds followed by a slow increase to 94°C at a rate of 0.1°C per second, with continuous fluorescence lecture. Using this procedure, the amplified products of influenza A, influenza B, hRSV, and the internal control could be differentiated by virtue of their specific T_m values i.e. 85, 83, 79, and 91°C,

30 respectively (Table 13 and Fig. 11).

Influenza A virus was detected by multiplex PCR (specific T_m value of 85°C) in 45/208 (21.6%) of the samples tested whereas the culture was positive for this virus in 14/100 (14%) tested NPA samples from children. For the subset of 100 NPA specimens tested by both methods, 88 (88%) had concordant results (11 positive, 77 negative) whereas discordant results were obtained for 12 (12%) samples (3 culture⁺/PCR⁻, 9 culture⁻/PCR⁺). Assuming that all positive results (by either culture or PCR) were true positive, the sensitivity values of the culture and the multiplex PCR for influenza A virus were 60.9% and 87.0%, respectively. There was no positive result for influenza B virus by any detection method as part of this study.

10

Example 8: Comparison of antigen detection and multiplex PCR assay for detection of the human respiratory syncytial virus.

Same as example 1, except that the multiplex PCR assay was used to detect hRSV from NPA samples obtained from 211 children with ARTI and compared with an antigen detection test for hRSV (TestPack, Abbott Laboratories, Abbott Park, IL).

hRSV was detected by the multiplex PCR (specific T_m value of 79°C) in 109/211 (51.7%) of the tested samples whereas the hRSV antigen detection test was positive in 93/211 (44.1%) tested NPA samples from children. Concordant results were found in 183 (86.7%) samples (87 positive, 96 negative) whereas discordant results were obtained for 28 (13.3%) samples (6 antigen⁺/PCR⁻, 22 antigen⁻/PCR⁺). A second RT-PCR assay for detection of another hRSV gene (glycoprotein G) was used to resolve the discrepancies (Peret et al., 1998, J Gen Virol. 79:2221-2229). After the resolution of the discrepant results, the sensitivity values of the antigen detection test and the multiplex PCR for hRSV were 80.5% and 96.5%, respectively.

Example 9: Evaluation of multiplex real time PCR assay for Influenza and human respiratory syncytial viruses.

We developed a real-time multiplex PCR assay for influenza viruses and HRSV using the melting curve analysis feature of the LightCycler instrument to rapidly distinguish viral products.

Materials and methods

Study population and antigenic test. Nasopharyngeal aspirates (NPA) were prospectively collected from children aged 0-3 years hospitalized for ARTI during the winter season 2001-02 at a single, university-based hospital, in Québec City (Canada) (Boivin, G., et al. 2003. Emerg. Infect. Dis. 9:634-640). The study was approved by the institutional review board of the Centre Hospitalier Universitaire de Québec. All samples were tested by the multiplex real-time PCR assay whereas antigen detection tests were performed on a subset of samples upon request by the treating physician. An aliquot (750 µl) of fresh NPA was first tested for the presence of HRSV and influenza A/B antigens using respectively the RSV TestPack (Abbott Laboratories, Abbott Park, IL) and the Directigen Flu A+B test (Becton Dickinson, Sparks, MD). The rest of the specimen was frozen at -80°C for a maximum of 3 months before nucleic acid extraction and PCR testing.

RNA extraction and cDNA synthesis. Viral RNA was extracted from 200 µl of NPA samples using the QIAamp Viral RNA Mini kit (QIAGEN, Mississauga, ON, Canada). Complementary DNA was then synthesized using 10 µl of the RNA preparation, 0.75 µM of random hexamer primers (Amersham Pharmacia Biotech, Balc d'Urfé, QC, Canada), and the Omniscript Reverse Transcriptase kit (QIAGEN) in presence of 300 copies of an internal control. The internal control template consisted of a 558-bp transcribed region of the herpes simplex virus type 2 DNA polymerase gene flanked by influenza B complementary primer sequences (see below) cloned into the pDrive plasmid (QIAGEN).

Real-time multiplex and conventional PCR assays. The multiplex PCR respiratory assay was designed to amplify conserved regions of the influenza A (Fouchier, R. A., et al. . 2000.. J. Clin. Microbiol. 38:4096-4101) and B (Li, J., S. Chen, and D. H. Evans. 2001.. J. Clin. Microbiol. 39:696-704) matrix genes as well as the fusion gene of HRSV (Mazzulli, T., et al, 1999. J. Infect. Dis. 180:1686-1689). Complementary DNA was amplified using a real-time PCR procedure with the LC Faststart DNA Master SYBR Green 1 kit (Roche Diagnostics, Laval, QC, Canada) in a LightCycler instrument (Roche Diagnostics). Each reaction had a total volume of 20 µl including 2 µl of cDNA and 18 µl of a reaction mixture containing 2.4 mM MgCl₂, 2 µl of Faststart DNA SYBR Green 1 Master Mix, 3% DMSO, 0.3 mM of influenza A and B primers (SEQ. ID Nos. 139-140 and 141-142), and 0.8 mM of HRSV primers (SEQ. ID Nos. 143-144). Cycling conditions included an initial denaturation step of 10 min at 94°C, followed by 50 cycles of 15 s at 94°C, 5 s at 58°C, and 25 s at 72°C. At the end of each cycle, the fluorescent signal was measured at a wavelength of 530 nm using the LightCycler Fluorimeter. The melting curve analysis program of the LightCycler was used to identify specific PCR products. Briefly, following the last amplification cycle, the reaction temperature was rapidly increased to 94°C and was then decreased to 60°C for 30 s followed by a slow increase to 94°C at a rate of 0.1°C per s, with continuous fluorescence monitoring. For resolution of discrepant results between the multiplex PCR assay and antigenic tests, conventional RT-PCR assays for detection of the glycoprotein G gene of HRSV (Peret, T. C., et al. 1998. J. Gen. Virol. 79:2221-2229) and the M2 gene of influenza A viruses (Klimov, A. I., et al. . 1995. J. Infect. Dis. 172:1352-1355) were performed retrospectively.

25 Results

Assay's characteristics. The sensitivity of each individual real-time PCR assay and of the multiplex PCR respiratory assay was determined for each target by testing serial dilutions of transcribed plasmids containing specific viral sequences. The lower limit of detection for influenza A, influenza B, and HRSV was found to be 10, 50 and 30 50 copies, respectively, in individual PCR assays whereas it was 50 copies for each target (10 out of 10 times for influenza A and 9 out of 10 times for influenza B and

HRSV) in the multiplex PCR test. No amplification signal was detected in the real-time multiplex assay when testing viral DNA or RNA from adenoviruses, rhinoviruses, enteroviruses (echo 11), parainfluenza viruses 1-3, human metapneumovirus, and herpesviruses (herpes simplex virus type 1 and 2, cytomegalovirus, and varicella-zoster virus). Specific melting temperature (T_m) values were first determined by testing a set of clinical isolates and vaccine strains (11 influenza A/H1, 10 influenza A/H3, 10 influenza B, and 17 HRSV) from different years (Table 14). A representative set of results is shown in Fig.11. The mean T_m values were 85.27 ± 0.22 , 83.47 ± 0.46 , and 79.51 ± 0.30 °C for influenza A, influenza B, and HRSV corresponding to fragments of 245, 524, and 380 bp, respectively, after gel electrophoresis (Table 14). We found no influence of influenza A subtype (H1 and H3) and HRSV genotype (A and B) on the T_m values (Table 14). The addition of 300 copies of the internal control in the assay's master mix resulted in a specific amplification peak with a T_m value of 91°C (558 bp) in the absence of amplified viral products.

Evaluation of the multiplex PCR respiratory assay in a pediatric population. The multiplex PCR assay was evaluated using extracted RNA from 208 NPA samples collected during a prospective pediatric study aimed at assessing the role of the human metapneumovirus in hospitalized children (0-3 years) with ARTI (Boivin, G., et al. 2003. *Emerg. Infect. Dis.* 9:634-640). The number of positive samples by multiplex PCR for influenza A, influenza B, and HRSV was found to be 45 (21.6%), 0 (0%), and 106 (51.0%), respectively. Overall, the rate of positivity for any of the 3 viruses was 66.8% (139/208) including 5.8% (12/208) of co-infections with influenza A and HRSV. None of the samples was considered to contain PCR-inhibitory samples as verified by the amplification of the internal control in all PCR-negative samples. The ranges of T_m values for influenza A- and HRSV-positive samples were 85.10 ± 0.39 and 79.60 ± 0.42 °C, respectively. All amplified products with one of the latter T_m values were confirmed as specific viral targets by visualizing the appropriate band on agarose gel. As none of the samples contained influenza B virus sequences (which correlates with the very occasional isolation of influenza B viruses in the Québec City

area during the 2001-2002 season), a clinical evaluation of the multiplex PCR assay was conducted for influenza A and HRSV only.

Evaluation of the multiplex PCR respiratory assay for detecting influenza A virus. From the 208 NPA evaluated by multiplex PCR, a subset of 172 samples were also tested for the presence of influenza antigens using the Directigen Flu A+B test (Table 15). Concordant results were obtained for 145 (84.3%) samples including 17 positive and 128 negative results by the two tests. A total of 27 (15.7%) samples showed discordant results with 25 being positive by the multiplex PCR assay only and two positive by the antigenic test only. In order to resolve the discrepant results, a second RT-PCR test targeting a conserved gene (M2) of influenza A viruses was performed as reported (Klimov, A. L., et al. . 1995. J. Infect. Dis. 172:1352-1355). As shown in Table 15, influenza A virus M2 sequences were detected by the conventional RT-PCR assay in 22 (81.5%) of the 27 discordant samples. Using the M2 RT-PCR assay as the reference test, the sensitivity, specificity, positive predictive and negative predictive values of the multiplex real-time PCR assay and the antigenic test for influenza A virus were 100%, 97.7%, 92.8%, 100% and 43.6%, 98.5%, 89.5%, 85.6%, respectively. The multiplex PCR respiratory assay was positive for HRSV in 12 (44.4%) of the 27 discordant influenza A samples with 10 of these samples showing dual HRSV/influenza A infections by the multiplex PCR assay (Table 16). The mean cycle threshold (C_T) values, which are inversely correlated with the amounts of viral RNA, were 28.1 and 33.0 for PCR-positive/antigen-positive and PCR-positive/antigen-negative samples, respectively (Table 16). These values correspond to approximately 5000 and 50 copies, respectively, when a specific influenza A plasmid was tested in the multiplex PCR assay.

Evaluation of the multiplex PCR respiratory assay for detecting HRSV. A total of 204 NPA samples could be evaluated by both the multiplex real-time PCR assay and the presence of HRSV antigens using the RSV TestPack (Table 17). Concordant results were found in 172 (84.3%) samples including 83 positive and 89 negative results by the two tests. Thirty-two (15.7%) samples had discordant results with 21

being positive by the multiplex PCR assay only and 11 by the antigenic test only. Conventional RT-PCR tests for the HRSV glycoprotein G (A and B subtypes) were performed as previously described (Peret, T. C., et al. 1998. *J. Gen. Virol.* 79:2221-2229) to resolve discrepant results. Specific sequences were detected by RT-PCR assays for the HRSV glycoprotein G gene in 26 (81.2%) of the 32 discordant samples including 11 A and 15 B strains (Table 18). Using the latter test as "gold standard", the sensitivity, specificity, positive predictive and negative predictive values of the multiplex real-time PCR assay and the antigenic test for HRSV were 94.5%, 98.9%, 99.0%, 94.0% and 81.6%, 94.7%, 94.7%, 81.8%, respectively. Four (66.7%) of the six false-negative HRSV results by multiplex PCR were positive for influenza A (Table 18). The mean C_T values of the PCR-positive/antigen-positive and PCR-positive/antigen-negative samples were 28.6 and 32.7, respectively (Table 18). These values correspond to approximately 1000 and 100 copies, respectively, when a specific HRSV plasmid was tested in the multiplex PCR assay.

15

Discussion

In this report, we describe a new multiplex real-time PCR assay for the most commonly detected respiratory viral pathogens, namely influenza viruses (A and B) and HRSV. The key features of this novel assay include its rapidity (turnaround time of approximately 2h30 including 30 min for sample preparation, 60 min for RT and 60 min for real-time PCR including viral identification), and its sensitivity (50 copies per assay for each viral target). Our results clearly demonstrate that our multiplex real-time PCR assay is more sensitive than commercially-available antigenic detection tests for both influenza A and HRSV with the greatest difference in sensitivity noted for influenza viruses.

25

Multiplex RT-PCR assays targeting up to nine different respiratory pathogens have been previously reported (Fan, J., K. J. Henrickson, and L. L. Savatski. 1998. *Clin. Infect. Dis.* 26:1397-1402, Grondahl, B., et al. 1999. *J. Clin. Microbiol.* 37:1-7, Kehl, S. C., et al. 2001. 39:1696-1701, Osiowy, C. 1998. *J. Clin. Microbiol.* 36:3149-3154, Stockton, J., et al. . 1998. *M. J. Clin. Microbiol.* 36:2990-2995, Valassina, M., A. M.

30

Cippone, M. G. Cusi, and P. E. Valensin. 1996. *Clin. Diag. Virol.* 8:227-232). More recently, a TaqMan-based real-time PCR assay for simultaneous detection of influenza A and B viruses has also been described (van Elden, L. J., et al. 2001. *J. Clin. Microbiol.* 39:196-200). Compared to conventional multiplex PCR assays, our
5 real-time PCR test is much more rapid because it avoids additional nested amplification and/or hybridization steps required for identification of viral products. Furthermore, the real-time amplification procedure minimizes the chances of contamination because there is no post-PCR processing of the samples. Although our multiplex real-time PCR assay based on melting curve analysis of amplicons does not
10 permit absolute quantification of the viral targets as in the TaqMan PCR procedure (van Elden, L. J., et al. 2001. *J. Clin. Microbiol.* 39:196-200), it allows for a larger number of viral targets to be simultaneously detected since there is no limitation related to the capability of the system to detect multiple dyes linked to detection probes with distinct emission wavelengths. Eventually, more than three viral targets
15 could be simultaneously detected in the LightCycler assay assuming the absence of interaction between PCR primers and a reproducible and discriminative T_m value for each viral amplicon.

Evaluation of our multiplex real-time PCR assay in a pediatric population with severe
20 ARTI clearly illustrated its clinical potential. Indeed, our assay identified a viral pathogen (influenza or HRSV) in two-thirds of the hospitalized children. These numbers are similar to the ones previously reported by our group in the same population using individual real-time assays for these viral pathogens (Boivin, G., et al. 2003. *Emerg. Infect. Dis.* 9:634-640). Notably, we were able to show that 6% of
25 hospitalized children had dual viral infections, a finding that has been reported albeit at a lower frequency by other investigators (Grondahl, B., et al. 1999. *J. Clin. Microbiol.* 37:1-7, Kehl, S. C., et al. 2001. 39:1696-1701, Osowy, C. 1998. *J. Clin. Microbiol.* 36:3149-3154). Rapid identification of these viral pathogens (influenza and HRSV) is of paramount importance for isolation purposes in the hospital setting
30 and early institution of specific antiviral therapy (Whitley, R. J., et al. 2001. *Pediatr. Infect. Dis. J.* 20:127-133). We are currently designing additional multiplex assays

based on the same technology to cover the whole spectrum of respiratory pathogens including parainfluenza viruses (PIV), adenoviruses, enteroviruses, and coronaviruses (Treanor, J. 2002. Respiratory Infections, p. 7-26, In Richard J. Whitley, Douglas D. Richman and Frederick G. Hayden (eds.), Clinical Virology, 2nd ed. ASM Press, Washington, D.C.). To this list, we should also add the recently described human metapneumovirus which has been reported by our group and others in up to 10% of ARTI in hospitalized children (Boivin, G., et al. 2003. Emerg. Infect. Dis. 9:634-640, Jartti, T., B. van den Hoogen, R. P. Garofalo, A. D. Osterhaus, and O. Ruuskanen. 2002.. Lancet. 360:1393-1394, Nissen, M. D., et al 2002. Med. J. Aust. 176:188, van den Hoogen, B. G., J. C. de Jong, J. Groen, T. Kuiken, R. de Groot, R. A. Fouchier, and A. D. Osterhaus. 2001.. Nat. Med. 7:719-724):

A specific evaluation of the real-time multiplex PCR assay for influenza A revealed that it was twice as sensitive as the rapid antigen detection test in use at our institution during the winter season of 2001-2002. None of the two PCR-negative, antigen-positive samples were confirmed by a second PCR targeting another conserved gene (M2) of influenza A whereas most (22/25, 88.0%) multiplex PCR-positive, antigen-negative were. In general, the false-negative antigenic test results could be explained by low amounts of viral RNA in those NPA samples as demonstrated by high C_T values in the corresponding real-time PCR assays. Three multiplex PCR test results were not confirmed by the influenza A M2 PCR test and were thus considered false-positive results. However, as we have not formally evaluated the sensitivity of the M2 PCR test and since viral cultures were not routinely done, we cannot rule out the possibility that such discrepant results were indeed true positive. We and other investigators have shown the superiority of RT-PCR tests over conventional methods (antigenic tests and viral culture) for detecting influenza viruses in clinical samples including those sent to the laboratory by mail (Boivin, G., I. Hardy, and A. Kress. 2001. J. Clin. Microbiol. 39:730-732, Ellis, J. S., D. M. Fleming, and M. C. Zambon. 1997. J. Clin. Microbiol. 35:2076-2082, Kehl, S. C., et al. 2001. 39:1696-1701, Steininger, C., et al. 2002. J. Clin. Microbiol. 40:2051-2056, van Elden, L. J., et al. 2001. J. Clin. Microbiol. 39:196-200). It is now common procedure at our institution

to confirm any negative antigenic tests results for influenza A and B by either viral culture or RT-PCR considering the high specificity but relatively poor sensitivity of the former assays.

- 5 For HRSV, the multiplex real-time PCR assay was also found to be more sensitive (by approximately 15%) than the rapid antigenic test when a second RT-PCR assay for the HRSV gG gene was used to resolve the discrepancies. Most of the false-negative antigenic test results could be explained by low amounts of viral RNA in the NPA samples as shown by mean C_T values of 32.7 for discordant results compared to
10 28.6 for concordant positive results. The specificity of the multiplex assay was excellent with only one positive result not confirmed by the second PCR test. In contrast, the multiplex PCR assay missed six cases of HRSV which tested positive in both the antigenic test and the second PCR assay for HRSV gG. Notably, four of those six samples tested positive for influenza A in the multiplex PCR assay. Thus,
15 although our multiplex PCR test has the capability of detecting dual infections (as seen in 6% of our cases), it is possible that high amounts of one virus could inhibit amplification of other pathogens. It is noteworthy that the multiplex PCR assay had the ability to amplify both HRSV genotypes as shown by the detection of 15 B and 11 A genotypes using specific PCR assays aimed at detecting variable regions of the
20 HRSV gG gene (Mazzulli, T., et al, 1999. *J. Infect. Dis.* 180:1686-1689). Although the rapid antigenic test had a relatively good performance (sensitivity: 82%, specificity 95%) for detection of HRSV in young children, the real advantage of PCR probably resides in the adult population where viral titers are lower and where antigenic tests are not recommended with upper respiratory tract samples (Englund, J. A., et al 1996.. 34:1649-1653).
25

- There are some limitations to our study which are worth mentioning. First, the absence of circulation of influenza B viruses during the study period prevents adequate validation of our multiplex assay for this pathogen and in particular for its
30 ability to detect mixed (influenza B and HRSV) infections. Second, the performance of our multiplex PCR assay cannot be generalized to other types of samples (throat

swabs, sputum) and to other populations (outpatients, adults) at the present time. Also, the absence of serological testing in our study may have underestimated the rate of viral infections especially for those children who presented late after the onset of symptoms. Finally, although it was relatively easy to discriminate between the
5 different viral pathogens based on their specific T_m values only (Table 14), we sometimes had to confirm viral amplicons by gel electrophoresis, which increases the assay's turnaround time.

In conclusion, we described a rapid and sensitive multiplex real-time PCR assay for
10 detecting influenza viruses and HRSV in children's NPA samples. This new assay is as specific and much more sensitive than currently-available antigen detection tests and could complement the latter in the hospital setting when there is high clinical suspicion despite negative results. Future work is needed to expand the panel of viral pathogens detected by such rapid molecular methods in order to eventually
15 circumvent the need for viral cultures. Also, future evaluation of the multiplex PCR assay is warranted in the adult population with severe ARTI.

Example 10 : Description and Evaluation of the Multiplex Respiratory-1 PCR.

- 5 Same as example 9, except that hMPV primers (as defined in SEQ. ID. Nos. 95 and 96 or 112 and 114) were added to the multiplex PCR assay (already containing primers of SEQ. ID. Nos. 139+140 for influenza A, 141+142 for influenza B, and 143+144 for hRSV) at a concentration of 0.3 μ M. The cycling conditions included an amplification step of 50 cycles of 15 s at 94°C, 5 s at 58°C, and 28 s at 72°C. The T_m values determined with the LightCycler instrument were 91°C, 85°C, 83°C, 79°C, and 82°C/77°C for the internal control, influenza A, influenza B, hRSV, and hMPV N/L genes, respectively (Fig. 7A for hMPV N primers of SEQ. ID. Nos. 95 and 96 and Fig. 7B for hMPV L primers of SEQ. ID. Nos. 112 and 114). The lower limit of detection of the multiplex respiratory-1 PCR was 10, 50, 100, and 100 copies for influenza A, influenza B, hRSV, and hMPV, respectively, in the presence of 500 copies of an internal control (plasmid flanked by influenza B complementary primer sequences). The multiplex respiratory-1 assay was found to be specific with no amplification of adenoviruses, enteroviruses, rhinoviruses, parainfluenza viruses 1-3, coronaviruses, herpes simplex virus type 1 and 2, varicella-zoster virus, cytomegalovirus, HHV-6, HHV-8, and Epstein-Barr virus.
- 10
- 15
- 20
- The multiplex respiratory-1 PCR was evaluated with NPA samples from 248 children ≤ 3 years with acute respiratory tract infections and 108 adults ≥ 50 years with exacerbation of chronic obstructive pulmonary disease or pneumonia requiring hospitalization during the 2002-03 winter/spring period in Québec City, Canada. In children, the detection rate for influenza A, influenza B, hRSV, and hMPV was 9/248 (3.6%), 4/248 (1.6%), 123/248 (49.6%), and 13/248 (5.2%), respectively. Five (2.0%) of the 248 children had a mixed viral infection. None of the NPA samples was found to contain PCR-inhibitory substances as shown by successful amplification of the internal control in all negative samples. In adults, the detection rate for influenza A, influenza B, hRSV, and hMPV was 3/108 (2.8%), 0/108 (0%), 13/108 (12.0%), and
- 25
- 30

2/108 (1.8%), respectively. One patient had a mixed viral infection (influenza A and hMPV). No samples contained PCR-inhibitory samples.

Example 11: Description and evaluation of the multiplex respiratory-2 PCR.

5

Same as example 9, except that the multiplex PCR assay is composed of the following PCR primers for detection of parainfluenza-1 (SEQ. ID. Nos. 147 and 148), parainfluenza-3 (SEQ. ID. Nos. 151 and 152), human adenoviruses (SEQ. ID. Nos. 153 and 155), and rhinoviruses/enteroviruses (SEQ. ID. Nos. 160 and 161). Primers are added in the PCR reaction mixture at a concentration of 0.2 μ M (SEQ. ID. Nos. 147 and 151), 0.3 μ M (SEQ. ID. Nos. 153 and 155), 0.4 μ M (SEQ. ID. Nos. 148 and 152), and 0.5 μ M (SEQ. ID. Nos. 160 and 161). The cycling conditions included an amplification step of 50 cycles of 15 s at 94°C, 5 s at 58°C, and 26 s at 72°C. The T_m values determined with the LightCycler instrument were 91.0°C, 80.0°C, 77.5°C, 89.0°C, 82.0°C, and 85.0°C for the internal control, parainfluenza-1, parainfluenza-3, adenoviruses, rhinoviruses, and enteroviruses, respectively (Fig. 8). The lower limit of detection of the multiplex respiratory-2 PCR was 50, 100, 10, 50, and 200 copies for parainfluenza-1, parainfluenza-3, adenoviruses, rhinoviruses, and enteroviruses, respectively, in the presence of 500 copies of an internal control (plasmid flanked by rhinovirus/enterovirus complementary primer sequences). The multiplex respiratory-2 assay was found to be specific with no amplification of influenza A, influenza B, hRSV, hMPV, coronavirus, herpes simplex virus type 1 and 2, varicella-zoster virus, cytomegalovirus, HHV-6, HHV-8, and Epstein-Barr virus.

25

Example 12 : Description of real-time PCR assays for the SARS-coronavirus.

30 Same as example 9, except that the PCR assay is composed uniquely of primers (two different possible sets) for the SARS-coronavirus as defined in SEQ. ID. Nos. 156-157 and 158-159. The concentration of the primers in the PCR reaction mixture is 0.5 μ M. The amplification step included 50 cycles of 15 s at 94°C, 5 s at 58°C, and 20 s at 72°C. The T_m for both PCR products is 83°C (Fig. 12).

Table 1: List of hMPV sequences revealed in the present invention

SEQ ID NO.	Isolate No.	Target gene
2	hMPV-118	Nucleoprotein
3	hMPV-174	Nucleoprotein
4	hMPV-176	Nucleoprotein
5	hMPV-193	Nucleoprotein
6	C-77807	Nucleoprotein
7	C-77748	Nucleoprotein
8	hMPV-217	Nucleoprotein
10	hMPV-163	Nucleoprotein
11	hMPV-174	Nucleoprotein
12	hMPV-184	Nucleoprotein
13	hMPV-185	Nucleoprotein
14	C-76837	Nucleoprotein
15	C-77807	Nucleoprotein
16	C-78123	Nucleoprotein
17	C-78132	Nucleoprotein
18	C-78163	Nucleoprotein
19	C-91205	Nucleoprotein
20	C-91364	Nucleoprotein
21	C-91830	Nucleoprotein
22	hMPV-Q86	Nucleoprotein
23	hMPV-Q97	Nucleoprotein
24	Can 97-83	Nucleoprotein
25	Can 98-75	Nucleoprotein
26	Can 98-75	Fusion type II
27	C-67345	Fusion type I
28	Can 97-83	Fusion type I
29	C-68661	Fusion type I
30	C-73750	Fusion type I
31	C-74549	Fusion type I
32	C-74783	Fusion type I
33	C-76837	Fusion type I
34	C-77748	Fusion type I
35	C-77807	Fusion type I
36	C-78132	Fusion type I
37	C-78163	Fusion type I
38	C-78406	Fusion type I
39	hMPV-174	Fusion type II
40	hMPV-193	Fusion type I
41	hMPV-194	Fusion type I
42	hMPV-198	Fusion type I
43	hMPV-217	Fusion type II
44	hMPV-208	Fusion type I
45	hMPV-223	Fusion type I
46	hMPV-228	Fusion type I

47	C-85885	Fusion type I
48	C-60298	Fusion type I
49	C-63778	Fusion type II
50	C-64199	Fusion type II
51	C-64200	Fusion type II
52	C-64536	Fusion type II
53	C-73769	Fusion type II
54	C-74501	Fusion type II
55	C-78429	Fusion type II
56	C-78562	Fusion type II
57	hMPV-118	Fusion type II
58	hMPV-200	Fusion type I
59	hMPV-176	Fusion type I
60	hMPV-193	Glycoprotein
61	hMPV-228	Glycoprotein
62	C-77748	Glycoprotein
63	C-78132	Glycoprotein
64	C-78406	Glycoprotein
65	C-73750	Glycoprotein
66	C-60298	Glycoprotein
67	C-74783	Glycoprotein
68	C-74549	Glycoprotein
69	C-77807	Glycoprotein
70	C-78163	Glycoprotein
71	C-68661	Glycoprotein
72	C-88448	Glycoprotein
73	C-88470	Glycoprotein
74	C-86316	Glycoprotein
76	hMPV-118	Polymerase
77	hMPV-163	Polymerase
78	hMPV-174	Polymerase
79	hMPV-184	Polymerase
80	hMPV-196	Polymerase
81	C-76837	Polymerase
82	C-77807	Polymerase
83	hMPV-185	Polymerase
84	C-78123	Polymerase
85	C-91205	Polymerase
86	C-91364	Polymerase
87	C-91749	Polymerase
88	hMPV-Q86	Polymerase
89	hMPV-Q97	Polymerase
90	Can 97-83	Polymerase

Table 2: List of hMPV PCR primers developed in the present invention

SEQ ID NO.	Target gene	Originating DNA	
		Position ^a	SEQ ID NO.
91	Nucleoprotein (N)	25	1
92	Nucleoprotein (N)	97	1
93	Nucleoprotein (N)	146 ^b	1
94	Nucleoprotein (N)	1025 ^b	1
95	Nucleoprotein (N)	58	9
96	Nucleoprotein (N)	986 ^b	9
97	Fusion (F)	3052	1
98	Fusion (F)	3810 ^b	1
99	Fusion (F)	4671 ^b	1
100	Matrix (M)	2165	1
101	Matrix (M)	2943 ^b	1
102	Phosphoprotein (P)	1248	1
103	Phosphoprotein (P)	2124 ^b	1
104	Glycoprotein (G)	6231	1
105	Glycoprotein (G)	6247	1
106	Glycoprotein (G)	6957 ^b	1
107	Glycoprotein (G)	6981 ^b	1
108	Glycoprotein (G)	6235	1
109	Glycoprotein (G)	7227 ^b	1
110	RNA polymerase (L)	142	75
111	RNA polymerase (L)	691 ^b	75
112	RNA polymerase (L)	7308	1
113	RNA polymerase (L)	7857 ^b	1
114	RNA polymerase (L)	7616	1

^a Position refers to nucleotide position of 5' end of primer. Oligonucleotides are 17-20 bp in length.

^b Primer is reverse-complement of target sequence.

Table 3: List of hMPV probes developed in the present invention

SEQ ID NO.	Target gene	Originating DNA	
		Position ^a	SEQ ID NO.
115	Nucleoprotein (N)	786	1
116	Nucleoprotein (N)	808	1
117	Fusion (F)	3163	1
118	Fusion (F)	3187	1
119	Matrix (M)	2636	1
120	Matrix (M)	2659	1
121	Phosphoprotein (P)	1492	1
122	Phosphoprotein (P)	1518	1
123	Glycoprotein (G)	6509	1
124	Glycoprotein (G)	6530	1
125	RNA polymerase (L)	7600	1
126	RNA polymerase (L)	7627	1
127	RNA polymerase (L)	7331	1
128	Nucleoprotein (N)	824	9
129	RNA polymerase (L)	434	75

^a Position refers to nucleotide position of 5' end of probe. Oligonucleotides are 21-25 bp in length.

Table 4: Length of amplicons obtained with different hMPV primer pairs which are objects of the present invention

SEQ ID NO.	Target Gene	Length of amplicon
91 + 94	Nucleoprotein (N)	1001 bp
92 + 94	Nucleoprotein (N)	929 bp
91 + 93	Nucleoprotein (N)	122 bp
97 + 99	Fusion (F)	1620 bp
97 + 98	Fusion (F)	759 bp
100 + 101	Matrix (M)	779 bp
102 + 103	Phosphoprotein (P)	877 bp
104 + 107	Glycoprotein (G)	751 bp
105 + 106	Glycoprotein (G)	711 bp
108 + 109	Glycoprotein (G)	993 bp
112 + 113	RNA polymerase (L)	550 bp
112 + 114	RNA polymerase (L)	308 bp

Table 5: Comparative evaluation of different hMPV primers for detection of hMPV strains

Strains No.	Type of sample	Year	N Primers SEQ. ID. Nos. 92+94		M primers SEQ. ID. Nos. 100+101		F primers SEQ. ID. Nos. 97+98		P primers SEQ. ID. Nos. 102+103		L primers SEQ. ID. Nos. 112+113	
			+/-	Tm (°C)	+/-	Tm (°C)	+/-	Tm (°C)	+/-	Tm (°C)	+/-	Tm (°C)
C-74501	I	2000	+	82.92	-		+	83.65	-		+	78.14
C-74783	I	2000	+	82.76	+	83.49	-		+	81.51	+	78.20
C-76837	I	2000	+	82.94	+	83.43	+	83.17	+	80.86	-	
C-77719	I	2001	+	83.26	+	84.64	-		-		+	77.87
C-77738	I	2001	+	82.99	-		-		-		+	77.51
C-77748	I	2001	+	83.06	+	82.96	+	83.10	+	81.27	+	77.38
C-77807	I	2001	+	82.99	+	83.26	+	83.25	+	80.65	+	78.34
C-78024	I	2001	+	82.86	+	83.95	-		-		-	
C-78028	I	2001	+	82.98	+	83.13	-		-		+	78.32
C-78123	I	2001	+	82.83	-		-		-		+	78.48
C-85984	I	2002	+	82.59	+	83.21	+	83.51	+	79.99	+	77.97
C-85448	I	2002	+	83.66	+	82.72	-		+	80.31	+	78.88
C-85983	I	2002	+	82.59	+	83.03	+	83.88	+	80.46	+	78.01
C-85988	I	2002	+	82.17	-		+	84.39	-		+	78.06
C-86316	I	2002	+	82.64	+	82.75	+	83.18	+	80.18	+	78.02
C-85126	I	2002	+	81.89	+	82.05	+	82.39	+	80.00	+	78.04
C-85473	I	2002	+	82.39	+	82.57	+	82.91	+	80.74	+	77.82
C-85332	I	2002	+	81.98	-		-		-		+	77.72
C-85596	I	2002	+	82.24	+	82.73	+	83.17	+	80.35	+	77.58
C-85542	I	2002	+	82.22	+	82.72	-		+	80.08	+	77.67
HMPV-118	S	2002	+	82.47	-		-		-		ND	ND
HMPV-174	S	2002	+	82.18	+	82.06	+	83.16	+	80.17	+	78.20
HMPV-176	S	2002	+	81.41	-		+	83.97	-		+	77.81
HMPV-193	S	2002	+	82.22	+	81.89	+	83.00	+	80.58	+	77.94
HMPV-194	S	2002	+	82.24	+	81.32	+	82.98	-		+	77.84
HMPV-198	S	2002	+	82.66	-		-		-		-	
HMPV-200	S	2002	+	82.53	-		+	83.46	-		+	77.62
HMPV-208	S	2002	+	82.11	+	81.81	+	83.33	+	80.06	+	78.40
HMPV-217	S	2002	+	82.13	+	81.58	+	83.01	-		+	77.76

64

HMPV- S	2002	+	81.89	+	82.36	+	83.04	-	+	78.10
223										
Mean			82.63 ± 0.87		82.75 ± 0.80		83.29 ± 0.45		80.48 ± 0.46	77.99 ± 0.33

Note: I = viral isolates; S = clinical specimens; Tm = melting temperature as determined by a LightCycler instrument.

Table 6: Size of the viral fusion (F) gene fragments obtained for different hMPV isolates following enzymatic digestion with restriction endonucleases

SEQ ID NO.	Fragments obtained after digestion with the restriction endonuclease		F genotype ^a
	ApaI	Avr II	
35	532 bp + 227 bp	759 bp	I
58	532 bp + 227 bp	759 bp	I
39	759 bp	447 bp + 312 bp	II
40	532 bp + 227 bp	759 bp	I
41	532 bp + 227 bp	759 bp	I
43	759 bp	447 bp + 312 bp	II
44	532 bp + 227 bp	759 bp	I
59	532 bp + 227 bp	759 bp	I
45	532 bp + 227 bp	759 bp	I

^aAs determined by DNA sequencing and phylogenetic studies.

Table 7. Type of laboratory confirmation by type of infection

Laboratory test	hMPV			hRSV			Influenza A			Adenovirus			PIV 2		
	Tests	Positive	done	Tests	Done	Positive	Tests	done	Positive	Tests	done	Positive	Tests	done	Positive
PCR	208	12		208	208	106	208		45	NA	NA	NA	NA	NA	NA
		5.8%				51.0%			21.6%						
Culture	100	2		100	100	31	100		14	100		6	100		1
		2%				31%			14%			6%			1%
Antigen detection (Immunofluorescence)	NA	NA		NA	NA	NA	NA		NA	81		1	76		1
												1%			1.3%
Antigen detection (ELISA)	NA	NA		204	204	94	100		14	NA	NA	NA	NA	NA	NA
						46.1%			14%						
Total (+/-) in at least one test	12 (5.8%)			118 (56.7%)			49 (23.6%)			6 (6%)			2 (2%)		

Delay between onset of symptoms and NPA, days	6.3/5.0	5.2/4.0	8.7/5.0	6.0/6.5	3.0/3.0
Mean/median					
Note: hMPV, human metapneumovirus; hRSV, human respiratory syncytial virus; PIV, parainfluenza virus; NPA, nasopharyngeal aspirate; NA, not available.					

Table 8. Signs and symptoms by type of viral infection

Signs and symptoms	HMPV	hRSV	Influenza A	Single virus	Multiple viruses	No virus detected	Total
	N=12	N=118	N=49	N=141	N=23	N=44	N=208
Fever	8 67%	67 57%	38 78%	85 60%	17 74%	25 57%	127 61%
Cough	12 100%	117 99%	47 96%	138 98%	23 100%	40 90%	201 97%
Rhinorrhea	11 92%	107 91%	41 84%	123 87%	22 96%	42 96%	187 90%
Retractions	11 92%	112 95%	40 82%	125 89%	22 96%	39 89%	186 89%
Wheezing	10 83%	77 65%	28 57%	83 59%	19 83%	31 71%	133 64%
Laerymation	3 25%	37 31%	15 31%	46 33%	6 26%	11 25%	63 30%
Diarrhea	1	19	13	24	5	10	39

	8%	17%	27%	17%	22%	23%	19%
Vomiting	3	9	5	10	4	1	15
	25%	8%	10%	7%	17%	2%	7%
Other	0	31	9	33	4	9	46
		26%	18%	23%	17%	21%	22%

Note: hMPV, human metapneumovirus; hRSV, human respiratory syncytial virus.

Table 9. Final diagnoses (complications) by type of viral infection

Complications	hMPV N=12	HRSV N=118	Influenza A N=49	Single virus N=141	Multiple viruses N=23	No virus detected N=44	Total N=208
Bronchiolitis	8 66.7%	99 83.9%	25 51.0%	98 69.5%	19 82.6%	25 56.8%	142 68.3%
Otitis	6 50.0%	70 59.3%	27 55.1%	78 55.3%	15 65.2%	24 54.6%	117 56.3%
Pneumonia	2 16.7%	30 25.4%	18 36.7%	39 27.7%	7 30.4%	12 27.3%	58 27.9%
Laryngotracheobronchitis	0	10 8.5%	6 12.2%	9 6.4%	4 17.4%	1 2.3%	14 6.7%
Sinusitis	0	3 2.5%	3 6.1%	2 1.4%	2 8.7%	1 2.3%	5 2.4%
Pharyngitis	0	1 0.9%	0	2 1.4%	0	2 4.6%	4 1.9%
Flu syndrome	0	2 1.7%	0	2 1.4%	0	4 9.1%	6 2.9%

Group	0	2	0	2	0	1	3
		1.7%		1.4%		2.3%	1.4%
Other	1	4	3	9	0	5	14
	8.3%	3.4%	6.1%	6.4%		11.4%	6.7%

Note: hMPV, human metapneumovirus; hRSV, human respiratory syncytial virus.

Table 10: List of PCR primers used for amplification of respiratory viruses

Primer SEQ ID NO.	Virus	Target Gene	Originating DNA Position ^a	SEQ ID NO.
139	Influenza A	Matrix	32 / 7	130-131
140	Influenza A	Matrix	276 ^b / 251 ^b	130-131
141	Influenza B	Matrix	82	132
142	Influenza B	Matrix	605 ^b	132
143	RSV	Fusion	1111	133
144	RSV	Fusion	1490 ^b	133
95	hMPV	Nucleocapsid	58	9
96	hMPV	Nucleocapsid	986 ^b	9
110	hMPV	Polymerase	142	75
111	hMPV	Polymerase	691 ^b	75
145	PIV-1	Nucleocapsid	64	134
146	PIV-1	Nucleocapsid	147 ^b	134
149	PIV-3	Nucleocapsid	416	135
150	PIV-3	Nucleocapsid	649 ^b	135
154	Adenovirus	Hexon	154	136
155	Adenovirus	Hexon	368 ^b	136
160	Enterovirus	5'-non coding region	94 / 71	137/138
161	Enterovirus	5'-non coding region / VP2	746 ^b / 603 ^b	137/138
141	IC ^c multiplex 1	DNA polymerase HSV-2	1	170
142	IC ^c multiplex 1	DNA polymerase HSV-2	558 ^b	170
160	IC ^c multiplex 2	DNA polymerase HSV-2	1	171
161	IC ^c multiplex 2	DNA polymerase HSV-2	554 ^b	171
112	hMPV	Polymerase	7308	1
114	hMPV	Polymerase	7616	1
147	PIV-1	Nucleocapsid	822	134
148	PIV-1	Nucleocapsid	1029	134
151	PIV-3	Nucleocapsid	792	135
152	PIV-3	Nucleocapsid	902	135
153	Adenovirus	Hexon	7	136
156	Sars-CoV	nsp9		
157	Sars-CoV	nsp9		
158	Sars-CoV	nsp11		
159	Sars-CoV	nsp11		

^aPosition refers to nucleotide position of 5'-end of primer. Oligonucleotides are 16-26 bp in length.

^bPrimer is reverse-complement of target sequence.

^cIC refers to the internal control used to monitor the efficiency of the PCR amplification

Table 11: List of viral probes developed in the present invention

Probe SEQ ID NO.	Target	Position ^a	Originating DNA SEQ ID NO.
162	Influenza A	114 ^b / 89 ^b	130/131
163	Influenza B	383	132
164	hRSV	1391 ^b	133
128	hMPV	824	9
129	hMPV	434	75
165	PIV-1	96	134
166	PIV-3	616 ^b	135
167	Adenovirus	228	136
168	Rhinovirus	467 ^b	138
169	Enterovirus	644 ^b	137
172	Internal Control ^c	321 ^b / 325 ^b	170/171

^aPosition refers to nucleotide position of 5'-end of primer. Oligonucleotides are 20-23 bp in length.

^bProbe is reverse-complement of target sequence.

^cA modified DNA polymerase gene sequence of herpes simplex virus (HSV) type 2 serves as the internal control template.

Table 12: Length of amplicons obtained with the different viral primer pairs

Primer SEQ ID NO.	Virus	Target gene	Length of amplicons
139 + 140	Influenza A	Matrix	245 bp
141 + 142	Influenza B	Matrix	524 bp
143 + 144	hRSV	Fusion	380 bp
95 + 96	hMPV	Nucleocapsid	929 bp
110 + 111	hMPV	Polymerase	550 bp
145 + 146	PIV-1	Nucleocapsid	84 bp
149 + 150	PIV-3	Nucleocapsid	234 bp
154 + 155	Adenoviruses	Hexon	215 bp
160 + 161	Rhinoviruses	5'-non coding region - VP2	533 bp
160 + 161	Enteroviruses	5'-non coding region	653 bp
141 + 142	IC multiplex 1	DNA polymerase HSV-2	558 bp
160 + 161	IC multiplex 2	DNA polymerase HSV-2	554 bp
112 + 114	hMPV	Polymerase	308 bp
147 + 148	PIV-1	Nucleocapsid	208 bp
151 + 152	PIV-3	Nucleocapsid	111 bp
153 + 155	Adenovirus	Hexon	362 bp
156 + 157	Sars-CoV	nsp9	198 bp
158 + 159	Sars-CoV	nsp11	368 bp

- 75 -

Table 13: Selection of primer sets with resulting melting temperature (T_m) values of the corresponding amplicons for each multiplex assay

Primer SEQ ID NO.	Target	Multiplex NO.	T _m (°C)
139 + 140	Influenza A	1	85
141 + 142	Influenza B	1	83
143 + 144	hRSV	1	79
95 + 96 or 112 + 114	hMPV	1	82 or 77
141 + 142	Internal Control	1	91
147 + 148	Parainfluenza 1	2	80.0
151 + 152	Parainfluenza 3	2	77.5
153 + 155	Adenovirus	2	89.0
160 + 161	Rhinovirus	2	82.0
160 + 161	Enterovirus	2	85.0
160 + 161	Internal Control	2	91

- 76 -

Table 14. Evaluation of the multiplex PCR respiratory assay with temporally-different strains of influenza (Flu) and human respiratory syncytial (HRSV) viruses.

Virus	Subtype / Genotype ^a	Year	Type	Strain	T _m (°C)
Flu A	H1	99-00	clinical	-	85.10
Flu A	H1	99-00	clinical	-	85.39
Flu A	H1	00-01	clinical	-	85.12
Flu A	H1	00-01	clinical	-	85.20
Flu A	H1	00-01	clinical	-	85.04
Flu A	H1	01-02	clinical	-	84.62
Flu A	H1	01-02	clinical	-	85.07
Flu A	H1	02-03	clinical	-	85.50
Flu A	H1	02-03	clinical	-	85.29
Flu A	H1	95	vaccine	A/Beijing/262/95	85.06
Flu A	H1	33	vaccine	A/WSN/33	85.25
Mean AH1					85.15±0.22
Flu A	H3	98-99	clinical	-	85.58
Flu A	H3	98-99	clinical	-	85.18
Flu A	H3	98-99	clinical	-	85.35
Flu A	H3	98-99	clinical	-	85.57
Flu A	H3	98-99	clinical	-	85.37
Flu A	H3	01-02	clinical	-	85.07
Flu A	H3	01-02	clinical	-	85.65
Flu A	H3	02-03	clinical	-	84.92
Flu A	H3	99	vaccine	A/Panama/2007/99	85.18
Flu A	H3	97	vaccine	A/Sydney/5/97	85.06
Mean AH3					85.31±0.25
Mean FluA					85.27±0.22
Flu B	-	98-99	clinical	-	83.87
Flu B	-	98-99	clinical	-	83.12
Flu B	-	00-01	clinical	-	83.01
Flu B	-	00-01	clinical	-	83.03
Flu B	-	00-01	clinical	-	82.88
Flu B	-	00-01	clinical	-	83.75
Flu B	-	00-01	clinical	-	83.17
Flu B	-	02-03	clinical	-	83.89
Flu B	-	02-03	clinical	-	83.84
Flu B	-	97	vaccine	B/Harbin/7/97	84.12
Mean FluB					83.47±0.46

- 77 -

Table 14. Evaluation of the multiplex PCR respiratory assay with temporally-different strains of influenza (Flu) and human respiratory syncytial (HRSV) viruses. (Continued).

HRSV	B	99-00	clinical	-	79.32
HRSV	A	99-00	clinical	-	79.83
HRSV	A	00-01	clinical	-	79.06
HRSV	A	00-01	clinical	-	78.85
HRSV	B	01-02	clinical	-	79.34
HRSV	B	01-02	clinical	-	79.52
HRSV	B	01-02	clinical	-	79.98
HRSV	B	01-02	clinical	-	79.52
HRSV	A	01-02	clinical	-	79.84
HRSV	B	01-02	clinical	-	79.42
HRSV	B	01-02	clinical	-	79.49
HRSV	B	01-02	clinical	-	79.48
HRSV	B	01-02	clinical	-	79.55
HRSV	B	01-02	clinical	-	79.32
HRSV	A	01-02	clinical	-	79.77
HRSV	A	01-02	clinical	-	79.92
HRSV	A	01-02	clinical	-	79.40
Mean HRSV-A					79.52±0.43
Mean HRSV-B					79.49±0.19
Mean HRSV					79.51±0.30

5

* Subtyping was performed by multiplex PCR for the influenza hemagglutinin gene (2) whereas genotyping was done by conventional PCR for the HRSV G gene (22)

- 78 -

Table 15. Evaluation of the multiplex PCR respiratory assay for detection of influenza A (Flu A) virus.

		Antigenic test (Directigen Flu A+B) (n = 172)		Flu A M2 PCR assay ^a (n = 27)	
		+	-	+	-
Multiplex PCR	+	17	25	22	3
	-	2	128	0	2

- 5 Sensitivity: $39/39 = 100\%$
 Specificity: $130/133 = 97.7\%$
 Positive predictive value: $39/42 = 92.8\%$
 Negative predictive value: $130/130 = 100\%$

- 10 ^a Only performed for samples showing discordant results with the
 multiplex PCR assay and the antigenic test.

Table 16: Evaluation of discordant test results for influenza A (Flu A) virus.

Sample No.	Multiplex PCR Flu A results (C _T) ^a	Antigenic test Flu A results	M2 PCR Flu A results	Other PCR- amplified viruses ^b
16	-	+	-	HRSV
23	+(27)	-	+	
24	-	+	-	HRSV
27	+(28)	-	+	HRSV
35	+(28)	-	+	HRSV
36	+(32)	-	+	
40	+(34)	-	+	
43	+(36)	-	+	HRSV
44	+(35)	-	+	
50	+(34)	-	+	
51	+(33)	-	+	HRSV
103	+(35)	-	+	
107	+(35)	-	+	
114	+(37)	-	-	
118	+(38)	-	+	HRSV
120	+(36)	-	+	HRSV
125	+(38)	-	+	
130	+(35)	-	+	HRSV
131	+(36)	-	-	
132	+(24)	-	+	HRSV
134	+(36)	-	-	
141	+(28)	-	+	
152	+(34)	-	+	
158	+(32)	-	+	
160	+(33)	-	+	HRSV
161	+(28)	-	+	HRSV
163	+(32)	-	+	

^a Cycle threshold value as determined by LightCycler.^b Viruses amplified by the multiplex PCR assay i. e. influenza (Flu) A and B as well as the human respiratory syncytial virus (HRSV).

- 80 -

Table 17: Evaluation of the multiplex PCR respiratory assay for detection of human respiratory syncytial virus (HRSV).

		Antigenic test (RSV TestPack) (n = 204)		HRSV gG PCR assays ^a (n = 32)	
		+	-	+	-
Multiplex PCR	+	83	21	20	1
	-	11	89	6	5

S

Sensitivity: 103/109 = 94.5%

Specificity: 94/95 = 98.9%

Positive predictive value: 103/104 = 99.0%

Negative predictive value: 94/100 = 94.0%

- 10 ^a Only performed for samples showing discordant results with the multiplex PCR assay and the antigenic test.

- 81 -

Table 18: Evaluation of discordant test results for human respiratory syncytial virus (HRSV).

Sample No.	Multiplex PCR HRSV results (C _T) ^a	Antigenic test HRSV results	Glycoprotein G PCR HRSV results (genotype)	Other PCR- amplified viruses ^b
16	+(32)	-	+(B)	
17	+(27)	-	+(B)	
26	+(36)	-	+(B)	
34	-	+	-	
35	+(33)	-	+(B)	Flu A
43	+(36)	-	+(A)	Flu A
46	+(34)	-	+(B)	
55	+(34)	-	+(B)	
58	+(33)	-	+(B)	
62	-	+	+(A)	Flu A
70	+(37)	-	+(B)	
71	+(35)	-	+(B)	
72	+(34)	-	+(B)	
73	+(26)	-	+(A)	
79	+(33)	-	+(B)	
80	-	+	-	
91	+(35)	-	+(A)	
95	+(29)	-	+(B)	
100	+(30)	-	+(A)	
112	-	+	-	
129	+(34)	-	+(B)	
130	+(35)	-	+(B)	Flu A
134	-	+	+(A)	Flu A
139	-	+	+(A)	
142	-	+	+(A)	Flu A
144	-	+	+(B)	Flu A
150	+(34)	-	-	
187	+(31)	-	+(A)	
191	+(29)	-	+(A)	

- 82 -

Table 18: Evaluation of discordant test results for human respiratory syncytial virus (HRSV). (Continued).

193	-	+	+(A)
211	-	+	-
212	-	+	-

5

^a Cycle threshold value as determined by Lightcycler.

^b Viruses amplified by the multiplex PCR assay i. e. influenza (Flu) A and B as well as the human respiratory syncytial virus (HRSV).

- 83 -

CLAIMS:

- 1- A method for detecting and/or quantifying human metapneumovirus (hMPV) in a sample said method comprising:
- 5 -providing at least one probe or primer specific for a nucleic acid sequence of said hMPV;
- contacting said at least one probe and/or primer with said sample to allow annealing of said probe and/or primer with said nucleic acid sequence; and
- detecting and/or quantifying said nucleic acid sequence using said annealed probe and/or primer.
- 10 2- The method as claimed in claim 1 wherein said nucleic acid sequence is comprised in a hMPV gene selected from a group consisting of nucleocapsid (N), phosphoprotein (P), matrix (M), fusion (F), membrane (M2), polymerase (L), glycoprotein (G) and small hydrophobic (SH) genes.
- 15 3- The method as claimed in claim 2 wherein said nucleic acid sequence is selected from any one of SEQ ID NO 2-8, 10-74, 76-90 or fragments thereof.
- 4- The method as claimed in claim 3 wherein said probe or primer is at least 10
- 20 nucleotides in length.
- 5- A method for detecting and/or quantifying human metapneumovirus (hMPV) in a sample said method comprising:
- 25 -performing reverse transcription using a template RNA derived from said sample to obtain a nucleic acid product;
- amplifying said nucleic acid product using a pair of oligonucleotide primers wherein said primers are specific for a sequence comprised in a hMPV gene selected from a group consisting of nucleocapsid (N), phosphoprotein (P), matrix (M), fusion

- 84 -

(F), membrane (M2), polymerase (L), glycoprotein (G) and small hydrophobic (SH) genes to produce an amplification product; and
-detecting and/or quantifying said amplification product.

5 6- The method as claimed in claim 5 wherein said oligonucleotide primers are at least 10 nucleotides in length and capable of hybridizing with any one of SEQ ID 2-8, 10-74, 76-90.

7- The method as claimed in claim 5 or 6 wherein said step of performing reverse
10 transcription comprises contacting said template RNA with a primer selected from SEQ ID Nos 91-114 and wherein said oligonucleotide primers for PCR are selected from SEQ ID Nos 91-114.

8- The method as claimed in any one of claim 5-7 wherein said oligonucleotide
15 primers are pairs of primers selected from SEQ ID Nos 91 and 94, 92 and 94, 91 and 93, 97 and 99, 97 and 98, 100 and 101, 102 and 103, 104 and 107, 105 and 106, 108 and 109, 112 and 113 and 112 and 114.

9- The method as claimed in claim 5 wherein said template RNA is contacted
20 with primer selected from SEQ ID NO 91-94 and 112-114 and wherein said oligonucleotide primers are selected from SEQ ID NO 91-94 and 112-114.

10- The method as claimed in any one of claim 5-9 wherein said step of
25 amplifying is performed using PCR.

11- The method as claimed in claim 10 wherein said PCR is RT-PCR.

12- The method as claimed in claim 11 wherein said RT-PCR is real time RT-PCR.

30

- 85 -

- 13- The method as claimed in claim 11 or 12 wherein said RT-PCR comprises detecting said amplification products by characteristic melting temperature of said amplification products.
- 5 14- The method as claimed in claim 11 or 12 wherein said RT-PCR comprises detecting and/or quantifying said amplification products with probes specific for said products.
- 15- The method as claimed in claim 14 wherein said probes are selected from SEQ
10 ID no 115-129.
- 16- A composition for detecting a hMPV said composition comprising a combination of oligonucleotides for performing reverse transcription of a template RNA from said hMPV virus and for an amplification reaction of a product obtained
15 from said reverse transcription wherein said oligonucleotides for said amplification reaction are specific for a sequence comprised in a hMPV gene selected from a group consisting of nucleocapsid (N), phosphoprotein (P), matrix (M), fusion (F), membrane (M2), polymerase (L), glycoprotein (G) and small hydrophobic (SH) genes.
- 20 17- The composition of claim 16 wherein said oligonucleotides for said amplification reaction are primers selected from the group comprising SEQ ID Nos 91-114 and combination thereof.
- 18- A kit comprising a combination of oligonucleotides wherein the oligonucleotides
25 are selected from:
SEQ ID NOs: 91-114 and the pairs of SEQ ID Nos 91 and 94, 92 and 94, 91 and 93, 97 and 99, 97 and 98, 100 and 101, 102 and 103, 104 and 107, 105 and 106, 108 and 109, 112 and 113 and 112 and 114.
- 30 19- A method for typing an hMPV strain which comprises the steps of:

- 86 -

-amplifying an hMPV nucleic acid fragment and detecting specific viral sequences;

-digesting said fragment with one or more restriction endonuclease; and

-analysing the restriction fragment length polymorphisms.

5

20- The method as claimed in claim 19 wherein said nucleotide sequences are found in fusion(F) or glycoprotein (G) gene.

21- The method as claimed in claim 20 wherein said sequences are selected from
10 SEQ ID NO 26-74.

22- A method for detecting and/or quantifying two or more respiratory viruses in a sample said method comprising:

15 -providing at least two probes and/or primers specific for a nucleic acid sequences in said two or more respiratory viruses;

-contacting said at least two probes and/or primers with said sample to allow annealing of said probes and/or primers with said nucleic acid sequences; and

-detecting and/or quantifying said nucleic acid sequences using said annealed probes and/or primers.

20

23- The method as claimed in claim 22 wherein at least one of said nucleic acid sequences is comprised in a hMPV gene selected from a group consisting of nucleocapsid (N), phosphoprotein (P), matrix (M), fusion (F), membrane (M2), polymerase (L), glycoprotein (G) and small hydrophobic (SH) genes.

25

24- The method as claimed in claim 23 wherein said nucleic acid sequence is selected from any one of SEQ ID NO 2-8, 10-74, 76-90 or fragments thereof.

25- The method as claimed in claim 24 wherein said probes and/or primers are at
30 least 10 nucleotides in length.

- 87 -

26- A method for detecting and/or quantifying two or more respiratory viruses in a sample said method comprising:

- 5 -performing reverse transcription using a template RNA derived from said sample to obtain nucleic acid products for each of said viruses;
 -amplifying said nucleic acid products using two or more pairs of oligonucleotide primers to produce an amplification product; and
 -detecting and/or quantifying said amplification products.

10

27- The method as claimed in claim 26 wherein at least one of said primers is specific for a sequence comprised in a hMPV gene selected from a group consisting of nucleocapsid (N), phosphoprotein (P), matrix (M), fusion (F), membrane (M2), polymerase (L), glycoprotein (G) and small hydrophobic (SH) genes.

15

28- The method as claimed in claim 27 wherein said oligonucleotide primers specific for a sequence comprised in a hMPV gene are capable of hybridizing with any one of SEQ ID 2-8, 10-74, 76-90.

20

29- The method as claimed in claim 28 wherein said oligonucleotide primers are at least 10 nucleotides in length.

30- The method as claimed in claim 29 wherein said oligonucleotide primer
25 specific for a sequence comprised in a hMPV gene are selected from SEQ ID Nos 91-114.

31- The method as claimed in any one of claim 26-30 wherein said at least two respiratory viruses comprise influenza A and B, human respiratory syncytial viruses A
30 and B, parainfluenza viruses (PIV) types 1-4, adenoviruses, rhinoviruses,

- 88 -

enteroviruses and coronaviruses or any combination thereof and wherein said primers and said two or more viruses are selected such that the detection and or quantification is specific for each of said viruses.

5 32- The method as claimed in claim 31 wherein said oligonucleotide primers are at least 10 nucleotides in length and capable of hybridizing with any one of SEQ ID NO 130-138 and 170-171.

10 33- The method as claimed in claim 31 or 32 wherein said at least two viruses are influenza A, human Respiratory syncytial virus, human metapneumovirus and influenza B.

15 34- The method as claimed in claim 31 or 32 wherein said at least two viruses are Parainfluenza 1, Rhinovirus, Adenovirus, Parainfluenza 3, Enterovirus, and coronavirus.

20 35- The method as claimed in claim 31 wherein said pairs of oligonucleotide primers are selected from SEQ ID Nos 139 and 140, 141 and 142, 143 and 144, 92 and 94, 112 and 113, 112 and 114, 145 and 146, 147 and 148, 149 and 150, 151 and 152, 154 and 155, 153 and 155, 156 and 157, 158 and 159, 160 and 161.

25 36- The method as claimed in any one of claim 26-35 wherein said step of amplifying is performed using PCR and said primer are selected such that the products of the amplifications can be distinguished.

37- The method as claimed in claim 36 wherein said PCR is RT-PCR.

38- The method as claimed in claim 37 wherein said RT-PCR is real time RT-PCR.

30

- 89 -

39- The method as claimed in claim 37 or 38 wherein said RT-PCR comprises detecting said amplification products by characteristic melting temperature of said amplification products.

5 40- The method as claimed in claim 37 or 38 wherein said RT-PCR comprises detecting and/or quantifying said amplification products with probes specific for said products.

41- The method as claimed in claim 40 wherein said probes are selected from SEQ
10 ID no 162-169 and 172.

42- A composition for detecting at least two respiratory viruses said viruses comprising at least hMPV said composition comprising a combination of oligonucleotides for performing reverse transcription of a template RNA from said at
15 least two respiratory viruses and for an amplification reaction of a product obtained from said reverse transcription wherein said oligonucleotides for said amplification reaction comprise oligonucleotides specific for a sequence comprised in a hMPV gene selected from a group consisting of nucleocapsid (N), phosphoprotein (P), matrix (M), fusion (F), membrane (M2), polymerase (L), glycoprotein (G) and small hydrophobic
20 (SH) genes.

43- The composition of claim 42 wherein said oligonucleotides for said amplification reaction are primers selected from the group comprising SEQ ID Nos 91-114 and combination thereof.

25

44- A method for simultaneously detecting two or more respiratory tract viruses in a sample, said method comprising:

- 90 -

Selecting primers specific for a nucleic acid sequence in each of said two or more viruses, said primers to be used in an amplification reaction for detection of said nucleic acid sequences;

5 distributing said primers in one or more assay mixture comprising said sample and at least two pairs of primers; and

wherein said selecting and said distributing is based on said detection of said nucleic acid sequence of said two or more viruses being substantially free of overlap and
10 enabling to distinguish between said two or more viruses.

45- The method as claimed in claim 44 wherein said amplification reaction is PCR.

15

46- The method as claimed in claim 45 wherein said PCR is RT-PCR.

47- The method as claimed in claim 45 wherein said RT-PCR is real time RT-PCR.

20

48- The method as claimed in claim 24 wherein said detection is melting temperature.

49- An isolated nucleic acid comprising a sequence that hybridizes under stringent
25 conditions to a hybridization probe the nucleotide sequence of which consists of any one of SEQ ID NO 2-8, 10-74, 76-90 or complements thereof.

50- A kit comprising a nucleic acid probe that is at least 10 nucleotides in length and is selected from a group consisting sequences hybridizable to SEQ ID Nos 2-8, 10-74,
30 76-90 or fragments thereof.

- 91 -

51- A human metapneumovirus (hMPV) deposited with the International Depository Authority of Canada under Accession number IDAC 111203-01 (CAN97-83).

- 5 52- A human metapneumovirus (hMPV) deposited with the International Depository Authority of Canada under Accession number IDAC 111203-02 (CAN98-75).

1/13




FIGURE 1




FIGURE 2

3/13




FIGURE 3

4/13




FIGURE 4

5/13




FIGURE 5

6/13




FIGURE 6

7/13




FIGURE 7

8/13




FIGURE 8

9/13




FIGURE 9




FIGURE 10 A

11/13




FIGURE 10 B

12/13




FIGURE 11

13/13




FIGURE 12

1/52

SEQUENCE LISTING

<110> Boivin, Guy

<120> METHOD FOR DETECTING RESPIRATORY VIRUSES

<130> 6013-148PCT

<150> CA 2,411,264

<151> 2002-12-19

<150> CA 2,418,004

<151> 2003-01-24

<160> 172

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 13350

<212> DNA

<213> human metapneumovirus

<220>

<221> gene

<222> (1)...(13450)

<223> human metapneumovirus, strain 00-1
Accession number af371337

<400> 1

gtataaatta	gattccaaaa	aaatatggga	caagtgaaaa	tgtctcttca	agggattcac	60
ctgagtgatt	tatcatataca	gcatgctata	ttaaaagagt	ctcagtacac	aataaaaaaga	120
gatgtgggta	caacaactgc	agtgacaccc	tcatcattgc	aacaagaaat	aacactgttg	180
tgtggagaaa	ttctgtatgc	taaacaatgc	gactacaaat	atgctgcaga	aataggaata	240
caatatatta	gcacagcttt	aggatcagag	agagtgcagc	agattctgag	gaactcaggg	300
agtgaagtc	aagcggctct	aaccagaacg	tactctctgg	ggaaaattaa	aaacaataaa	360
ggagaagatt	tacagatggt	agacatacac	ggggtagaga	agagctgggt	agaagagata	420
gacaaaaga	caaggaaaac	aatggcaacc	ttgcttaagg	aatcatcagg	taatatccca	480
caaaatcaga	ggccctcagc	accagacaca	cccataatct	tattatgtgt	aggtgcctta	540
atattcacta	aactagcatc	aaccatagaa	gtgggactag	agaccacagt	cagaagggtc	600
aaccgtgtac	taagtgargc	actcaagaga	taccctagaa	tggacatacc	aaagattgcc	660
agatccttct	atgacttatt	tgaacaaaaa	gtgtatcaca	gaagtttgtt	cattgagtat	720
ggcaaaagcat	taggctcatc	atctacaggg	agcaaaagcag	aaagtctatt	tgtaaatata	780
ttcatgcaag	cttatggggc	cgggtcaaca	atgctaagggt	gggggggtcat	tgccaggtca	840
tccaacaata	taatgttagg	acatgtatcc	gtccaagctg	agttaaaaa	ggtcacagaa	900
gtctatgact	tgggtgcgaga	aatggggcct	gaatctggac	ttctacattt	aaggcaaaagc	960
ccaaaagctg	gactgtttatc	actagccaac	tgtcccaact	ttgcaagtgt	tggtctcgga	1020
aatgcctcag	gcttaggcat	aatcgggtatg	tatcgaggga	gagtaccaa	cacagaatta	1080
ttttcagcag	ctgaaagtta	tgccaaaaagt	ttgaaagaaa	gcaataaaat	aaatttctct	1140
tcattaggac	ttacagatga	agagaaagag	gctgcagaa	atctcttaaa	tgtgagtgac	1200
gacagtcaaa	atgattatga	gtaattaaaa	aagtgggaca	agtcaaaatg	tcattccctg	1260
aaggaaaaaga	tattcttttc	atgggtaatg	aagcagcaaa	attagcagaa	gctttccaga	1320
aatcattaag	aaaaccaggt	cataaaaagt	ctcaatctat	tataggagaa	aaagtgaata	1380
ctgtatcaga	aacattggaa	ttacctacta	tcagttagacc	tgcaaaacca	accataccgt	1440
cagaaccaaa	gttagcatgg	acagataaag	gtggggcaac	caaaactgaa	ataaagcaag	1500
caatcaaaagt	catggatccc	attgaagaag	aagagtctac	cgagaagaag	gtgctaccct	1560
ccagtgatgg	gaaaaccctt	gcagaaaaga	aactgaaacc	atcaactaac	accaaaaaga	1620
aggttttcatt	tacaccaaat	gaaccaggga	aatatacaaa	gttggaaaaa	gatgctctag	1680
atttgctctc	agataatgaa	gaagaagatg	cagaatcttc	aatcttaacc	tttgaagaaa	1740
gagatacttc	atcattaagc	attgaggcca	gattggaatc	aatagaggag	aaattaagca	1800
tgtatttagg	gctatttaaga	acactcaaca	ttgctacagc	aggaccaca	gcagcaagag	1860
atgggatcag	agatgcaatg	attggcgtaa	gagagggaat	aatagcagac	ataataaagg	1920
aagctaaagg	gaaagcagca	gaaatgatgg	aagaggaaat	gaagtcaacga	tcaaaaatag	1980
gaaatggtag	tgtaaaaata	acagaaaaag	caaaaagagct	caacaaaatt	gttgaagatg	2040
aaagcacaag	tggagaatcc	gaagaagaag	aagaaccaaa	agacacacaa	gacaatagtc	2100
aagaagatga	catttiaccag	ttaattatgt	agtttaataa	aaataaacaa	tgggacaagt	2160
aaaaatggag	tcctacctag	tagacaccta	tcaaggcatt	ccttacacag	cagctgttca	2220
agttgatcta	atagaaaagg	acctgttacc	tgcaagccta	acaatatgggt	tccctttgtt	2280
tcaggccaac	acaccaccag	cagtgctgct	cgatcagcta	aaaaccctga	caataaccac	2340
tctgtatgct	gcatcacaaa	atggtccaat	actcaaagtg	aatgcatcag	cccaagggtgc	2400

2/52

agcaatgtct	gtacttccca	aaaaatttga	agtcaatgcy	actgtagcac	tcgatgaata	2460
tagcaaatg	gaatttgaca	aaactcacagt	ctgtgaagta	aaaacagttt	acttaaccaac	2520
catgaaacca	tacgggatgg	tatcaaaaatt	tgtgagctca	gccaaatcag	ttggcaaaaa	2580
aacacatgat	ctaactgcac	tatgtgattt	tatggatcta	gaaaagaaca	cacctgttac	2640
aataaccagca	ttcatcaaat	cagtttcaat	caaagagagt	gagtcagcta	ctgttgaagc	2700
tgttataagc	agtgaagcag	accaagctct	aaacacaggcc	aaaattgcac	cttatgctgg	2760
attaattatg	atcatgacta	tgaacaatcc	caaaggcata	ttcaaaaagc	ttggagctgg	2820
gactcaagtc	atagtagaac	taggagccta	tgtccaggct	gaaagcataa	gcaaaaatag	2880
caagagctgg	agcctacaa	ggacaagata	tgtcttgaag	tccagataac	aaccaagcac	2940
cttggccaa	agctactaac	cctatctcat	agatcataaa	gtcaccattc	tagttatata	3000
aaaatcaagt	tagaacaga	attaaatcaa	tcaagaacgg	gacaaataaa	aatgtcttgg	3060
aaagtgggtg	tcattttttc	attgttaata	acacctcaac	acggtcttaa	agagagctac	3120
ttagaagaat	catgtagcac	tataactgaa	ggatattctca	gtgttctgag	gacaggttgg	3180
tacaccaatg	tttttacact	ggaggtaggc	gatgtagaga	accttacatg	tgccgatgga	3240
cccagcttaa	taaaaaacaga	attagacctg	acaaaaagtg	cactaagaga	gtcagagaac	3300
gtttctaggag	atcaactggc	aagagaggag	caaattgaaa	atcccagaca	atctagattc	3360
attgccaaaa	caatagcact	cggtgttgca	actgcagctg	cagttacagc	aggtgttgca	3420
accatgaaag	ccatccggct	tgaagtgaa	gtaacagcaa	ttaagaatgc	cctcaaaaag	3480
gagctgaaag	artttgtgag	caagaatcta	ggagttcgtg	tgttggcaac	tgcatgtaga	3540
attgctgacc	tgaaaaatggc	cggttagcttc	acacgtgcaa	tcaacaaaaa	caagtgcgac	3600
gtgcgggaat	tttcagacaa	cgctggaata	agtcatttca	acagaagggt	cctaaatgtt	3660
gatgctgaac	tagccagagc	tggttccaac	atgccaaat	ctgcaggaca	aataaaaactg	3720
atgtttggaga	accgtgcaat	ggtaagaga	aaaggggtcg	gattcctgat	aggagtttac	3780
ggaaagctccg	taattttacat	gggtcaactg	ccaattcttg	gggttataga	cagcctttgc	3840
tggaatagtaa	aagcagcccc	ttcttgttca	ggaaaaaagg	gaaactatgc	ttgctcttta	3900
agagaagacc	aaggatgtgt	ttgtcaaaaat	gcagggtcaa	ctgtttacta	cccaaatgaa	3960
aaagactgtg	agaccatgtc	ttttgcgaca	ttttgcgaca	cagcagcagg	aatcaatgtt	4020
gctgagcagt	caaaggagtg	caacataaac	atatctacta	ctaattacc	atgcaaagtt	4080
agcacaggaa	gacatccctat	cagtatgggt	gcactatctc	ctcttggggc	tttgggttgc	4140
tgctacaagg	gagtgagctg	ttccattggc	agcaacagag	tagggatcat	caagcaactg	4200
aacaaaggct	gctcttatat	aaccaacca	gacgcagaca	cagtgaacat	agacaacact	4260
gtataccagc	taagcaaatg	tgaaggcgaa	cagcatgtta	taaaaggaa	gccagtgtca	4320
agcagcttgg	acccagtcac	gtttcctgaa	gatcaattca	atgttgcact	tgaccaagrt	4380
ttcgagagca	ttgagaacag	tcaggccctg	gtggatcaat	caaacagaat	cctaagcagt	4440
gcagagaaag	gaaacactgg	cttcatcatt	gtaataatc	taattgctgt	ccttggctct	4500
accatgatcc	tagtgagtg	ttttatcata	ataaagaaaa	caaagaaacc	cacaggagca	4560
cctccagagc	tgagtgggtg	cacaaacaat	ggcttcatac	cacataatta	gttaattaaa	4620
aataaagttaa	attaaaataa	attaaaatta	aaaataaaaa	tttgggacaa	atcataatgt	4680
ctcgcaaggc	tcctgtgcaaa	tatgaagtgc	ggggcaaatg	caatagagga	agtgagtgtca	4740
agtttaacca	caattactgg	agttggccag	atagataact	arttaataaga	tcaaatatt	4800
tattaaatca	acttttaagg	aacactgata	gagctgatgg	cttatcaata	atatcaggag	4860
caggcagaga	agataggaca	caagattttg	tcctaggttc	caccaatgtg	gttcaaggtt	4920
ataattgatga	taaccaaaagc	ataacaaaag	ctgcagcctg	ttacagtcta	cataatataa	4980
tcaaaacaact	acaagaagtt	gaagtttaggc	aggctagaga	taacaaacta	ctgcacagca	5040
aacatgtagc	acttcacaac	ttagtcttat	cttatatgga	gatgagcaaa	actcctgcat	5100
ctttaatcaa	caatctcaag	agactgcca	gagagaaaact	gaaaaaatta	gcaaggtcca	5160
taattgactt	atcagcaggt	gctgaaaaatg	actcttcata	tgccctgcaa	gacagtga	5220
gcactaatca	agtgcagtga	gcatgggtcca	gttttcatta	ctatagaggt	tgatgacatg	5280
atatggactc	acaaggactt	aaaagaagct	ttatctgatg	ggatagtga	gtctcatact	5340
aacatttaca	attgttatrt	agaaaacata	gaaattatat	atgtcaaggc	ttacttaagt	5400
tagtaaaaaa	atcatcagagt	gggataaatg	acaatgataa	cattagatgt	cattaaaagt	5460
gatgggtctt	caaaaaacatg	tactcacctc	aaaaaaataa	ttaaagacca	ctctggtaaa	5520
gtgcttattg	tacttaagtt	aatattagct	ttactaacat	ttctcacagt	aacaatcacc	5580
atcaattata	taaaagtggg	aaacaatctg	caaatatgcc	agtcaaaaac	tgaatcagac	5640
aaaaaggact	catcatcaaa	taccacatca	gtcacacaac	agactactct	aaatcatgat	5700
atcacacagt	atttttaaaag	tttgattcaa	aggatatcaa	actctgcaat	aaacagtgac	5760
acatgctgga	aaataaacag	aaatcaatgc	acaaatataa	caacatacaa	atttttatgt	5820
tttaaatctg	aagacacaaa	aaccaacaat	tgtgataaac	tgacagattt	atgcagaaac	5880
aaacaaaaac	cagcaggttg	agtgtatcac	atagtagaat	gccattgtat	atacacagtt	5940
aaatgggaag	gctatcatta	cccaaccgat	gaaacccaat	cctaaatgrr	aacaccagat	6000
taggaatccat	ccaagtctgt	tagttcaaca	atrtagttat	rtaaaaatat	rttgaaaaa	6060
agtaagtctt	tatgataact	cataataata	agtaataatt	aattgcttaa	tcatacatcac	6120
aacattatcc	gaaaccataa	ctattcaatt	taaaaagtta	aaaacaataa	catgggacaa	6180
gtagtatttg	aggtgaaagt	ggagaacatt	cgaaacaatag	atatgtctca	agcaagagta	6240
aaaaatctgt	tgccacgcag	caaatgcttt	aaaaatgcct	cttgggtcct	catagggata	6300
actacattga	gtattgccct	caatatctat	ctgatcataa	actataaaat	gcaaaaaaac	6360
acatctgaat	cagaaacatca	caccagctca	tcacccatgg	aatccagcag	agaaactcca	6420
acgggtcccca	cagacaactc	agacaccaac	tcaagccac	agcatccaac	tcaacagctcc	6480
acagaaggct	ccacactcta	ctttgcagcc	tcagcaagct	caccagagac	agaaccaaac	6540
tcaacaccag	atacaacaaa	ccgcccgcgc	ttcgtcgaca	cacacacaa	accaccaagc	6600
gcaagcagaa	caaagacaag	tccggcagtc	cacacaaaaa	acaacccaag	gacaagctct	6660

3/52

agaacacatt	ctccaccacg	ggcaacgaca	aggacggcac	gcagaaccac	cactctccgc	6780
acaagcagca	caagaaagag	accgctccaca	gcacacagtc	aacctgacat	cagcgcaaca	6840
accacaaaaa	acgaagaagc	aagtcacagcg	agcccacaaa	catctgcaag	cacacaagca	6900
atacaaaagga	aaagcgtgga	ggccaacaca	tcaacaacat	acaaccaaac	tagttaacaa	6960
aaaatacaaa	ataactctaa	gataaaacat	gcagacacca	acaatggaga	agccaaaaga	7020
caattcacaa	tctccccaaa	aaggcaacaa	caccatatta	gctctgcca	aatctccctg	7080
gaaaaaacac	tcgcccata	accaaaaata	ccacaaccac	ccaagaaaa	aaactgggca	7140
aaacaacacc	caagagacaa	ataacaatgg	atcctctcaa	tgaatccagt	gttaatgtct	7200
atcttcttga	ctcatatctt	aaaggagtga	ttcctttag	tgagactaat	gcaattgggt	7260
catgtctctt	aaaaagacct	tacctaataa	atgacaacac	tgcaaaagrt	gccatagaga	7320
atcctgttat	cgagcatgrr	agactcaaaa	atgcagtcaa	ttctaagatg	aaaatatcag	7380
attacaagat	agtagagcca	gtaaacatgc	aacatgaaat	tatgaagaat	gtacacagtt	7440
gtgagctcac	attattaaaa	cagtttttaa	caaggagtaa	aaatattagc	actctcaaat	7500
taaatatgat	atgtgattgg	ctgcagttaa	agtctacatc	agatgatacc	tcaatcctaa	7560
gttttataga	tgtagaatttt	atacctagct	gggttaagca	ttggtttagt	aattgggtaca	7620
atctcaacaa	gttgattctg	gaattcagga	aagaagaagt	aataagaact	ggttcaatct	7680
tgtgtagggtc	attgggttaaa	ttagtttttg	ttgtatcatc	atatggatgt	atagtcaaga	7740
gcaacaaaaa	caaaagagtg	agcttcttca	catacaatca	actgttaaca	tggaaagatg	7800
tgatgttaag	tagattcaat	gcaaatrrrr	gtatatgggt	aagcaacagt	ctgaatgaaa	7860
atcaagaagag	gctaggggtg	agaagtaatc	tgcaaggcat	atttaactaat	aagctatatg	7920
aaaactgtaga	ttatatgctt	agttttatgt	gcaatgaagg	tttctcaett	gtgaaagatg	7980
tcgaaggctt	tattatgagt	gaaattctta	ggattactga	acatgctcaa	ttcagtaacta	8040
gatttagaaaa	tactttatta	aatggattaa	ctgatcaatt	aacaaaatta	aaaaataaaa	8100
acagactcag	agttcatggt	accgtgttag	aaaaaataga	ttatccaatg	tacgaagttg	8160
tacttaagtt	attaggagat	actttgagat	gtattaaatt	attaatcaat	aaaaacttag	8220
agaatgctgc	tgaattatac	tatatattta	gaattattcg	tcaccaatg	gtagatgaaa	8280
gagatgcaat	ggatgctgct	aaattaaaca	atgaaatcac	aaaaatcctt	aggtgggaga	8340
gcttgacaga	actaagaggg	gcattcatat	taaggattat	caaaggattt	gtagacacaa	8400
acaaaagatg	gccccaaatt	aaaaacttaa	aagtgttag	taagagatgg	actatgtact	8460
tcaaagcaaa	aagttacccc	agtcaacttg	aattaaagca	acaagatttt	rtagagcttg	8520
ctgcaatata	gtttgaacaa	gagttttctg	tccctgaaaa	aaccaacctt	gagatggtat	8580
taaatgatata	agctatatca	cctcctaaaa	gattaatatg	gtctgtgrrt	ccaaaaaatt	8640
acttacttga	gaaaaataaa	aatcgatatc	tagaagagac	tttcaatgca	agtgatgttc	8700
tcaaaaacag	aagagtacta	gagtactatt	tgaagataaa	taaarttcgac	caaaaagaac	8760
rtaaaagttta	tggtgttaaa	caagaatatt	taaatgataa	ggatcaattt	gtctcgctaa	8820
ctggaataaga	aagagaattta	agtgtaggta	gaattgtttg	targcaacca	ggaaaacagc	8880
gacaaatata	aatattggct	gaaaaattgt	tagctgataa	tattgtacct	tttttccag	8940
aaaccttaac	aaagtatggt	gatctagatc	ttcagagaat	aatggaaatc	aaatcggaac	9000
ttcttcttat	taaaactaga	agaaatgata	gttataataa	ttacattgca	agagactcca	9060
tagtaacaga	tttaagtaag	ttcaaccaag	cctttaggta	tgaaaactaca	gcgatctgtg	9120
cggatgtagc	agatgaacta	catggaaacac	aaagcctatt	ctgttggtta	catcttatcg	9180
ttccttatgac	aacaatgata	tggtcctata	gacatgcacc	accagaaaca	aaagggtgaat	9240
atgatataga	taagatagaa	gagcaaaagt	gtttatatag	atatcatatg	gggtggtattg	9300
aaggatgggt	tcaaaaaactc	tggaacaatg	aagctatatc	tctattagat	gttgatctcg	9360
taaaaaacag	atgtcnaatg	acatctttat	taaacgggtga	caaccaatca	atagatgtaa	9420
gtaaaaccagt	taagtattct	gagggttttag	atgaagtga	agcagattat	agcttggtcg	9480
taaaaaatgrr	aaaagaaata	agagatgcac	acagaaatat	agggcataaa	citaaagaag	9540
gggaacacata	tatatcaaga	gatcttcagt	ttataagtaa	gggtgattcaa	tctgaaggag	9600
taatgcatcc	tacccctata	aaaaagatct	taagagtggg	accatggata	aacacaatat	9660
tagatgacat	taaaaaccagt	gcagagtcaa	tagggagtct	atgtcaggaa	ttagaattta	9720
ggggggaaag	cataatagtt	agtctgatatt	taaggaaatt	ttggctgtat	aatttatata	9780
tgcatgaatc	aaagcaacac	cccctagcag	ggaagcagtt	attcaaacaa	ctaaataaaa	9840
cattaacatc	agtgcagaga	ttttttgaaa	taaaaaagga	aatgaagta	gtagatctat	9900
ggatgaaacat	accaatgcag	tttgaggag	gagatccagt	agctctctat	agatctttct	9960
atagaaggac	ccctgatttt	tttaactgaag	caatcagta	tgtggatatt	ctgttaagaa	10020
tatcagccaa	cataagaat	gaagcgaaaa	taagtttctt	caaagcctta	ctgtcaatag	10080
aaaaaaatga	acgtgctaca	ctgacaacac	taattgagaga	tcctcaagct	gttggctcag	10140
agcgacaagc	aaaagttaaca	agtgatatca	atagaacagc	agttaccagc	atcttaagtc	10200
tttctccaaa	tcaacttttc	agcgatagtg	ctatacacta	cagtagaaat	gaagaagagg	10260
tcggaaatcat	ttgtgacaa	ataaaccttg	tttatcctca	tggactgaga	gttttgtatg	10320
aatcattacc	ttttcataaa	gctgaaaaag	ttgtgaatat	gatatacagga	acgaaatcca	10380
taaccaactt	attacagaga	acatctgcta	ttaatgttga	agatattgac	agagctgtat	10440
ccatgatgct	ggagaaccra	ggattattat	ctagaatatt	gtcagtagtt	gttgatagta	10500
tagaaattcc	aaccaaatct	aatggtaggc	tgatatgttg	tcagatatct	agaaccttaa	10560
gggagacatc	atggaataat	atggaaatag	ttggagtaac	atcccctagc	atcactacat	10620
cctatgtatg	catatatgca	actagctctc	atttgaaagg	gataatcatt	gaaaagtcca	10680
gtactgacag	aactacaaga	ggctcaagag	gtccaaagag	cccttgggta	gggtcgagca	10740
ctcaagagaa	aaaattagtt	cctgtttata	acagacaaat	tctttcaaaa	caacaaagag	10800
acagctaga	agcaattgga	aaaatgagat	gggtatataa	agggacacca	ggtttaagac	10860
gattactcaa	taagatttgt	cttggaagtt	taggcattag	ttacaatgtt	gtaaaaactt	10920
tattacctag	gtttatgagt	gtaaaatttc	tacacaggtt	atctgtcagt	agtagacctta	10980
tggaattccc	agctacagtt	ccagcttata	gaacaacaaa	ttaccatttt	gacactagtc	11040

4/52

ctattaatca	agcactaagt	gagagatttg	ggaatgaaga	tattaatttg	gtcttccaaa	11100
atgcaatcag	ctgtgggaatt	agcataatga	gtgtagtaga	acaattaact	ggtaggagtc	11160
caaaacagtt	agttttaata	cctcaattag	aagaaataga	cattatgcca	ccaccagtgt	11220
ttcaaggga	attcaattat	aagctagtag	ataagataac	ttctgatcaa	catatcttca	11280
gtccagacaa	aatagatarg	ttaacactgg	ggaaaatgct	catgcccact	ataaaaaggtc	11340
agaaaaacaga	tcagttcctg	aacaagagag	agaattattt	ccatgggaat	aatcttattg	11400
agtctttgtc	agcagcgtra	gcattgtcatt	gggtggggat	attaacagag	caatgtatag	11460
aaaataaata	tttcaagaaa	gactgggggtg	acgggttcatt	atcgatcat	gcttttatgg	11520
acttcaaaat	attcctatgt	gtctttaaata	ctaaactttt	atgtagttgg	gggtcccaag	11580
ggaaaaacat	taaagatgaa	gatatagtag	atgaatcaat	agataaactg	ttaaggattg	11640
ataatacttt	ttggagaatg	ttcagcaagg	ttatgtttga	atcaaagggt	aagaaaagga	11700
taatgtttat	tgatgtaaaa	tttcratcat	tagtaggta	tatagggttt	aagaattggt	11760
ttatagaaca	gttgagatca	gctgagttgc	atgaggtacc	ttggattgtc	aatgccgaag	11820
gtgatctggg	tgagatcaag	tcaattaaaa	tcattttgca	actgatagag	caaagtttat	11880
ttttaagaa	aactgttttg	aactatacag	ataatggcaca	tgctctcaca	agattaatca	11940
gaaagaagtt	gatgtgtgat	aatgcactat	taactccgat	tccatcccca	atggttaatt	12000
taactcaagt	tattgatcct	acagaacaat	tagcttattt	ccctaagata	acatttgaaa	12060
ggctaaaaaa	ttatgaact	agttcaaat	atgttaaagg	aaagctaaca	aggaattaca	12120
tgatactgtt	gccatggcaa	catgttaata	gatataactt	tgtctttagt	tctactggat	12180
gtaaagrtag	tctaaaaaca	tgcartggaa	aacttatgaa	agatctaaac	cctaaagtcc	12240
gttactttat	ttgagaagg	gcaggaaatt	ggatggccag	aacagcatgt	gaatatcctg	12300
acatcaaatt	tgtatacaga	agtttaaaag	atgaccttga	tcattcattat	cctttggaat	12360
accagagagt	tataggagaa	taaagcagga	taatagatag	cggtgaaggg	ctttcaatgg	12420
aaacaacaga	tgcaactcaa	ggatattgat	acacagagta	agcaaagatg	agcaaagatg	12480
ctttattaat	aactttatgt	gatgcagaat	tttaaggacag	agatgatttt	tttaagatgg	12540
taattctatg	gaggaaacat	gtattatcat	gcagaatttg	cactacttat	gggacagacc	12600
tcattttatt	cgcaaatgat	catgtcaaat	actgcaaatg	aaaattacct	ttttttgtga	12660
gatcagtagc	cacttttatt	atgcaaggta	gtaaacigt	aggctcagaa	tgctacatac	12720
ctttaacact	agggcaccac	aacaattttac	cctgccatgg	agaaatacaa	aattctaaga	12780
tgaaaatagc	agtggttaat	gattttttatg	ctgcaaaaaa	acttgacaat	aaatctattg	12840
aagccaactg	taaatcactt	ttatcagggc	taagaataacc	gataaataag	aaagaattaa	12900
atagacagag	aaggttatta	acactacaaa	gcaaccattc	ttctgtagca	acagttggag	12960
gtagcaagg	catagagtc	aaatggtraa	caaacaaggc	aaacacataa	attgattggt	13020
tagaacatat	tttaaatctt	ccaaaagggtg	aattaaatta	tgattttttt	gaagcattag	13080
aaaatactta	ccctaataatg	attaaactaa	tagataatct	agggaaatga	gagataaaaa	13140
aactgatcaa	agtaactgga	tatatgcttg	taagtataaa	atgaaaaatg	ataaaaaatga	13200
taaaataggt	gacaacttca	tactattcca	aagtaaatcat	ttgattatgc	aattatgtaa	13260
tagttaatta	aaaactaaaa	atcaaaagrt	agaaactaac	aactgtcatt	aagtttatta	13320
aaaataagaa	attataattg	gatgtatcag				13350<210> 2
<211>	782					
<212>	DNA					
<213>	human metapneumovirus					
<220>						
<221>	gene					
<222>	(1)...(782)					
<223>	human metapneumovirus, strain Quebec Isolate No hmpv-118					
<400>	2					
atgtaggcac	cacaactgca	gtgacacctt	catcattgca	gcaagagata	acacttttgt	60
gtggagagat	tcttttactt	aaacatactg	attacaaaata	tgctgcagag	atagggatcac	120
aatatattttg	cacggctcta	ggatcagaaa	gagtacaaca	gatttttaaga	aattcaggca	180
gtgaagttica	gggtggttcta	accaagacat	actcrttagg	gaaaggtaaa	aatagtaaaag	240
gggaagaggtt	gcaaatgtta	gatatacatg	gagtggaataa	gagttgggta	gaagaaatag	300
acaaaagaggc	aagaaaaaca	atggtgactt	tgctaaagga	atcatcaggt	aacatccac	360
aaaaccagag	gccttcagca	ccagacacac	caataatttt	attatgtgta	ggtgctttaa	420
tatttactaa	actagcatca	acaatagaag	ttggactaga	gactacagtt	agaagagcta	480
acagagtgct	aagtgatgag	actcaaagat	accctagggt	agatatacca	aagattgcta	540
gatctttcta	tgaactattt	gagcagaaag	tgtattacag	gagtcatttc	attgagtagt	600
ggaaaagcttt	aggctcatct	tgcacagga	gcaaaagcaga	aagttttgttt	gtaaaatatat	660
ttatggcagc	ttatggagcc	ggtaaaacaa	tgctaaaggtg	gggtgtcatt	gccagatcat	720
ctaataatfac	attgctaggg	catgtatctg	tgcaacctgg	atttgaagca	agtcacagag	780
gt						782
<210>	3					
<211>	878					
<212>	DNA					
<213>	human metapneumovirus					
<220>						
<221>	gene					

5/52

<222> (1)...(878)

<223> human metapneumovirus, strain Quebec isolate No
hMPV-174

<400> 3

atgtgggtac	caccactgca	gtgacaccct	catcattgca	acaagaaata	acgctgttgt	60
gtggagaaat	tctgtatgct	aaacatgctg	attacaaata	tgctgcagaa	ataggaatac	120
aatatattag	cacagcttta	ggatcagaga	gagtgacaga	gattctgagg	aactcaggca	180
gtgaagtcca	agtggcttta	accagaacgt	actctctggg	gaaagttaaa	aacaataaag	240
gagaagattt	acagatgtta	gacatacacg	gggtagagaa	gagctgggta	gaagagatag	300
acaaagaagc	aaggaaaaa	atggcaacct	tgcttaagga	atcatcaggt	aatatcccg	360
aaaatcagag	gccctcagca	ccagacacac	ccataatctt	attatgtgta	ggtgccttaa	420
tatttactaa	actagcatca	acatagaaag	tggtactaga	gaccacagtc	agaagggcta	480
accgtgtact	aagtgtatga	ctcaagagat	accctagaat	ggacatccca	aaaattgcca	540
gatccttcta	tgacttattt	gaacaaaaag	tgtatcacag	aagtttgttc	attgagtatg	600
gcaaagcatt	aggctcatca	tctacaggca	gcaaagcaga	aagtctatit	gttaatatat	660
tcatgcaagc	ttatggagcc	gggtcaaacaa	tgctaagggt	gggggtcatt	gccagggtcat	720
ccaacaatat	aatgttatga	catgtatccg	tccaagctga	gttaaacaca	ggtcacagaa	780
gtctatgact	tggtgcgaga	aatgggtcct	gaatctggac	ttctacattt	aaggcaaagc	840
ccaaaagctg	gactgtatca	ctagccaact	tcccaact			878

<210> 4

<211> 876

<212> DNA

<213> human metapneumovirus

<220>

<221> gene

<222> (1)...(876)

<223> human metapneumovirus, strain Quebec isolate No
hMPV-176

<400> 4

atgtaggccc	cacaactgca	gtgacacctt	catcattgca	gcaagagata	acacttttgt	60
gtggagagat	tctttacact	aaacatactg	attacaaata	tgctgcagag	atagggatag	120
aatatatttg	cacggctcta	ggatcagaaa	gagtacaaca	gattttaaga	aattcaggta	180
gtgaagtcca	ggtgggtcta	accaagacat	actctttagg	gaaaggtaaa	aatagttaaag	240
gggaagagtt	gcaaatgtta	gatatacatg	gagtggaana	gagttgggta	gaagaaatag	300
acaaagaggg	aagaaaagaca	atgggtgactt	tgctaaagga	atcatcaggt	aacatccac	360
aaaaccagag	gccttcagca	ccagacacac	caataatctt	attatgtgta	ggtgcgttaa	420
tatttactaa	actagcatca	acaatagaag	ttggactaga	gactacagtt	agaagagcta	480
acagagtgct	aagtgtatgc	ctcaaaagat	accctagggt	agatatacca	aagattgcta	540
gatcttttta	tgaactatit	gagcagaaag	tgtattacag	gagctctatc	attgagtatg	600
ggaaagcttt	aggctcatct	tcaacaggaa	gcaaagcaga	aagtttgttt	gtaaatatat	660
ttatgcaagc	ttatggagcc	gggtcaaacaa	tgctaagggt	gggtgtcatt	gccagatcat	720
ctaacaacat	aatgtctagg	catgtatctg	tgcaagctga	attgaaacaa	gttacagagg	780
ttatgatatt	ggtgagagaa	atgggtcctg	aatctgggct	tttactctta	agaacaagtc	840
caaaagccag	actgttatcg	tggctaattg	cccaat			876

<210> 5

<211> 869

<212> DNA

<213> human metapneumovirus

<220>

<221> gene

<222> (1)...(869)

<223> human metapneumovirus, strain Quebec isolate No
hMPV-193

<400> 5

atgtgggtac	aacaactgca	gtgacaccc	catcattgca	acaagaaata	acgctgttgt	60
gtggagaaat	tctgtatgct	aaacatgctg	attacaaata	tgctgcagaa	ataggaatac	120
aatatattag	cacagcttta	ggatcagaga	gagtgacaga	gattctgagg	aactcaggta	180
gtgaagtcca	agtggcttta	accagaacgt	actctctggg	gaaagttaaa	aacaataaag	240
gagaagattt	acagatgtta	gacatacacg	gggtagagaa	gagctgggta	gaagagatag	300
acaaagaagc	aaggaaaaa	atggcaacct	tgcttaagga	atcatcaggt	aatatcccg	360
aaaatcagag	gccctcagca	ccagacacac	ccataatctt	attatgtgta	ggtgccttaa	420
tatttactaa	actagcatca	acatagaaag	tggtactaga	gaccacagtc	agaagggcta	480
accgtgtact	aagtgtatga	ctcaagagat	accctagaat	ggacatccca	aaaattgcca	540
gatccttcta	tgacttattt	gaacaaaaag	tgtatcacag	aagtttgttc	attgagtatg	600
gtaaagcatt	aggatcatca	tctacaggca	gcaaagcaga	aagtctatit	gttaatatat	660

6/52

tcattgcaagc	ttatggagcc	ggtcaaacaa	tgctaagggtg	gggggtcatt	gccaggtcat	720
ccaacaatat	aatgttttaga	catgtatccg	tccaaggctg	agttaaaaca	ggtcacagaa	780
gtctatgact	tggtgcgaga	aatgggtcct	gaatctggac	ttctacattt	aaggcaaacg	840
ccaaaagctg	gactgtttac	actagccaa				869

<210> 6
 <211> 848
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(848)
 <223> human metapneumovirus, strain Quebec Isolate No C-77807

<400> 6						
atgtaggcac	cacaactgca	gtgacacctt	catcattgca	acaagaaata	acacttttgt	60
gtggggaaat	tctttacact	aaacacactg	attacaata	tgctgctgag	ataggaatgc	120
aatatatttg	cacagctcta	ggatcagaaa	gagtacaaca	gattttgaga	aactcaggta	180
gtgaagtcca	ggtggtttta	accaaaccat	actccttagg	gaaaggcaa	aacagtaaag	240
gggaagagct	gcagatgtta	gatatacatg	gagrggaaaa	gagttgggta	gaagaaatag	300
acaaagaggc	aagaaagaca	atggtaactt	tgcttaagga	gtcatcaggt	aacatcccac	360
aaaaccagag	accttcagca	ccagacacac	caataatttt	attatgtgta	ggtgccctaa	420
tattacttaa	actagcatca	acaatagaag	ttggattaga	gactacagtt	agaagagcta	480
atagagtgtc	aagtgatgca	ctcaaaagat	accctagggt	agatatacca	aagattgcta	540
gactttttta	tgaactattt	gaacaaaag	tgtactacag	aagttttattc	attgagtacg	600
gaaaagcttt	aggctcatct	tcaacaggaa	gcaaagcaga	aagttttgtt	gtaaatatat	660
ttatgcaagc	ttatggagct	ggtcaaacac	tgctaagggtg	gggtgtcatt	gccagatcat	720
ccaacaacat	aatgctaggg	catgtatctg	tgcaatctga	attgaagcaa	ttcacagagg	780
tttatgactt	ggtgagagaa	atgggtcctg	aatctgggct	tttatcatcta	agacaaagtc	840
caaaggca						848

<210> 7
 <211> 894
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(894)
 <223> human metapneumovirus, strain Quebec Isolate No C-77748

<400> 7						
atgtgggtac	aacaactgca	gtgacaccc	catcattgca	acaagaaata	acgctgttgt	60
gtgggaaat	tctgtatgct	aaacatgctg	attacaata	tgctgcagaa	ataggaatgc	120
aatatatttg	cacagcttta	ggatcagaga	gagtgcagca	gattctgagg	aactcaggca	180
gtgaagtcca	agtggcttta	accagaacgt	actctctggg	gaaagttaa	aacaataaag	240
gagaagattt	acagatgtta	gacatacacg	gggtagagaa	gagctgggta	gaagagatag	300
acaaagaagc	aaggaaaaca	atggcaacct	tgcttaagga	atcatcaggt	aatatcccac	360
aaaatcagag	gccctcagca	ccagacacac	ccataatctt	attatgtgta	ggtgccctaa	420
tatttactaa	gctagcatca	accatagaag	tgggactaga	gaccacagtc	agaagggcta	480
accgtgtact	aagtgatgca	ectaagagat	accctagaat	ggacatccca	aaaattggca	540
gatccttcta	tgactttatt	gaacaaaaag	tgtatcacag	aagttttgtc	attgagtatg	600
gcaaagcatt	aggctcatca	tctacaggca	gcaaacgaga	aagtctattt	gcttatatat	660
tcatgctgagc	ttatggagcc	ggtccaacaa	tgctaagggtg	gggggtcatt	gccagatcat	720
ccaacaatat	aatgttttaga	catgtatctg	tccaagctga	atttaaaca	gacacagaag	780
tccatgactt	ggtggtagaa	atggggcctg	aatctggact	ttctacttta	aggcaaacgc	840
caaaggctgg	actgtttaca	ctagccaact	gtcccaactt	tgcaagtgtt	gttc	894

<210> 8
 <211> 850
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(850)
 <223> human metapneumovirus, strain Quebec Isolate No hMPV-217

7/52.

```

<400> 8
atgtgggtac aacaactgca gtgacacccat catcattgca acaagaaata acgctgttgt 60
gtggagaaat tctgtatgct aaacatgctg attacaataa tgctgcagaa atagggaatac 120
aatatattag cacagcttta ggatcagaga gagtgcagca gattctgagg aactcaggta 180
gtgaagtcca agtgggtctta accagaacgt actctctggg gaaagttaaa aacaataaag 240
gagaagattt acagatgtta gacatacacg gggtagagaa gagctgggta gaagagatag 300
acaaagaagc aaggaaaaca atggcaacct tgcctaaagg atcatcagg atatatccac 360
aaaatcagag gccctcagca ccagacacac ccataatctt attatgtgta ggtgccttaa 420
tatttactaa gctagcatca accatagaag tgggactaga gaccacagtc agaagggtta 480
accgtgtrac aagtgtatgca ctcaagagat accctagaat ggacatccca aaaattgcca 540
gatccttcta tgacttattt gaacaaaaag tgtatcacag aagtttggtc attgagtatg 600
gtaaagcatt aggatcatca tctacaggca gcaagcaga aagtcctatt gttaatatat 660
tcatgcaagc ttatggagcc ggtcaaaaca tgctaagggt ggggggtcatt gccaaagtc 720
ccaacaatat aatgttagga catgtatccg tccaagctga attaaaaaca ggtcccagaa 780
gtccatgact tgggtgcgaga aatggggcct gaatctggac ttctactttt aaggcaagc 840
ccaaaagctg

```

```

<210> 9
<211> 1185
<212> DNA
<213> human metapneumovirus

```

```

<220>
<221> gene
<222> (1)...(1185)
<223> human metapneumovirus, strain 00-1, accession
number af371337

```

```

<400> 9
argtctcttc aagggattca cctgagtgat ttatcataga agcatgctat attaaaagag 60
tctcagtaca caataaaaag agatgtgggt acaacaactg cagtgacacc ctcatcattg 120
caacaagaaa taacactgtt gtgtggagaa attctgtatg ctaaacatgc tgactacaaa 180
tatgtcgcag aaatagggaat acaatatatt agcacagctt taggatcaga gagagtgcag 240
cagattctga ggaactcagg cagtgaagtc caagtgggtc taaccagaac gtactctctg 300
gggaaaatta aaaacaataa aggagaagat ttacagatgt tagacataca cggggtagag 360
aagagctggg tagaagagat agacaaagaa gcaaggaaaa caatggcaac ctgtcttaag 420
gaatcatcag gtaatatccc acaaaatcag aggcctcag caccagacac acccataatc 480
ttattatgtg taggtgcctt aatattcact aaactagcat caaccataga agtgggacta 540
gagaccacag tcagaagggc taaccgtgta ctaagtgatg cactcaagag ataccctaga 600
atggacatac caaagattgc cagatccttc tatgacttat ttgaacaaaa agtgtatcac 660
agaagtttgt tcattgagta tggcaaaagca ttaggctcat catctacagg cagcaaaagca 720
gaaagttctat ttgttaatat attcatgcaa gcttatgggg ccggtcaaac aatgctaagg 780
tgggggggtca ttgccaggtc atccaacaat ataattgttag gacatgtrac cgtccaagct 840
gagttaaaac aggtcacaga agtctatgac ttggtgcgag aaatgggccc tgaatctgga 900
cttctacatt taaggcaaaag cccaaaagct ggactgttat cactagccaa ctgtcccaac 960
tttgcaagtg ttgttctcgg aaatgcctca ggtctaggca taatcgggtat gtatcgaggg 1020
agatgaccaa acacagaatt atttcagca gctgaaagtt atgccaaaag ttgaaaagaa 1080
agcaataaaa taaatttctc ttcattagga cttacagatg aagagaaaga ggctgcagaa 1140
catttcttaa atgtgagtga cgacagtcaa aatgattatg agtaa 1185

```

```

<210> 10
<211> 700
<212> DNA
<213> human metapneumovirus

```

```

<220>
<221> gene
<222> (1)...(700)
<223> human metapneumovirus, strain Quebec Isolate No
HMPV-163

```

```

<400> 10
tgcagaaata ggaatacaat atattagcac agcttttaga tcagagagag tgcagcagat 60
tctgaggaaac tcaggcagtg aagtcgaagt ggtcttaacc agaactgact ctctggggaa 120
agttaaaaac aataaaggag aagatttaca gatgtrtagc atacacgggg tagagaagag 180
ctgggtagaa gagatagaca agagaagcaag gaaaacaatg gcaaccttgc ttaagggaatc 240
atcaggtaat atccacaaaa atcagaggcc ctacgacca gacacacca taatcttatt 300
atgtgttagt gccttaatat ttactaaact agcatcaacc atagaagtgg gactagagac 360
cacagtcaga agggccaacc gtgtactaag tgatgcactc aagagatacc ctagaatgga 420
catcccaaaa attgccagat cttctatga cttatttgaa caaaaagtgt atcacagaag 480
tttgttcatt gagtatggca aagcattagg ctcatcatct acaggcagca aagcagaag 540
tctatttgtt aatatattca tgcaagctta tggagccggt caaacaatgc taaggttggg 600

```

8/52

ggtaattgcc aggtcatcca acaatataat gttaggacat	660
aaaacaggtr acagaagtct atgacttggg gcgagaaatg	700

<210> 11
 <211> 705
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(705)
 <223> human metapneumovirus, strain Quebec isolate No
 hmpv-174

<400> 11 tattagcaca gctttaggat cagagagagt gcagcagatt ctgaggaact caggcagtga	60
agtccaagt gtcttaacca gaacgtactc ictggggaaa gttaaaaaca ataaaggaga	120
agattttacag atgttagaca tacacggggg agagaagagc tgggtagaag agatagacaa	180
agaagcaagg aaaacaatgg caaccttgct taagggaatca tcaggtaata tcccgcaaaa	240
tcagaggccc tcagcaccag acacacccat aatcttatta tgtgtagggt ccttaatat	300
tactaaacta gcatcaacca tagaagtggg actagagacc acagtcagaa gggctaaccg	360
tgtactaagt gatgactca agagataccc tagaatggac atcccaaaaa ttgccagatc	420
cttctatgac ttatttgaac aaaaagtgt tccagagaat ttgttcattg agtatggcaa	480
agcattaggc tcatcatcta caggcagcaa agcagaaagt gtatttgtaa atatattcat	540
gcaagcttat ggagccggtc aaacaatgct gagggtgggg gtcattgcca ggtcatccaa	600
caatataatg ttaggacatg tatccgtcct agctgagcta aaaccgggtc cagaagtcta	660
tgacttgggt cgagaaatga ttcttgaatc tngacttcta catta	705

<210> 12
 <211> 700
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(700)
 <223> human metapneumovirus, strain Quebec isolate No
 hmpv-184

<400> 12 aaatatgctg cagaatagg aatacaatat attagcacag ctttaggac agagagagt	60
cagcagattc tgaggaaactc aggcagtga gtcacagttg tcttaaccag aacgtactct	120
ctggggaaag ttaaaaacaa taaaggagaa gattttacaga tgttagacat acacggggta	180
gagaagagct gggtagaaga gatagacaaa gaagcaagga aaacaatggc aaccttgctt	240
aaggaatcat caggtaatat cccacaaaat cagaggccct cagcaccaga cacaccata	300
atcttattat gtgtagggtc ctttaatttt actaaactag catcaaccat agaagtggga	360
ctagagacca cagtcagaag ggccaaccgt gtactaagt atgcactcaa gagataccct	420
agaattggaca tcccaaaaat tgcagatcc ttctatgact tatttgaaca aaaagtgtat	480
cacagaagtt gtattcattga gtatggcaca gcattaggct catcatctac aggcagcaaa	540
gcagaaagtc tatttgtaa tatattcatg caagcttatg gagccgggtc aacaatgcta	600
aggtgggggg taattgccag atcatccaac aatataatgt taggacatgt atccgtccaa	660
gctgagttaa aacaggctac agaagtctat gacttagtgc	700

<210> 13
 <211> 650
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(650)
 <223> human metapneumovirus, strain Quebec isolate No
 hmpv-185

<400> 13 taggaatca atatattagc acagctctag gatcagagag agtacagcag attctaagaa	60
artcaggcag tgaagtccaa gtgggtttta ccagaacgta ctccrtgggg aaagttaaaa	120
acaacaaagg agaagattta cagatgttag acatacacgg agtagagaaa agctgggtgg	180
aagagataga caaagaagca agaaaaacaa tggcaacttt gcttaaaaga tcatcaggca	240
atattccaca aaatcagagg ccttcagcac cagacacacc tataatctta ttatgtgtag	300
gtgccttaat atttaccaaa ctatagatcaa ctatagaagt gggattagag accacagtca	360
gaagagctaa ccgtgtacta agtgatgcac tcaaaagata ccctaggatg gacataccaa	420

9/52

aaatcgctag atctttctat gacttatttg aacaaaaagt gtattacaga agcttggtca 480
 ttgagtatgg caaagcatta ggctcatcct ctacaggcag caaagcagaa agttttattcg 540
 ttaatatatt catgcaagct tacgggtgctg gtcaaaacaat gctgagggtgg ggagtcattg 600
 ccagggtcatc taacaatata atgttaggac atgtaictgt ccaagctgag 650

<210> 14

<211> 650

<212> DNA

<213> human metapneumovirus

<220>

<221> gene

<222> (1)...(650)

<223> human metapneumovirus, strain Quebec Isolate No C-76837

<400> 14

acaaatattgc atcataaata ggaatacaat atattagcac agcttttagag atcagagaga 60
 gtgcagcaga ttctaaggaa ctacaggcagt gaagtcacag tgggtctaac cagaacgtac 120
 tctctgggga aggttaaaaa caataaaggga gaagatttac agatgttaga catcacggg 180
 gtagagaaga gctgggtaga agagatagac aaagaagcaa ggaaaaacaat ggcaaccttg 240
 cttaaggaat catcaggtaa tatccacaa aatcagaggg cctcagcacc agacacacc 300
 ataactttat tatgtgtagg tgcttaata ttactaaac tagcatcaac catagaagt 360
 ggactagaga ccacagtcag aagggtctaac cgtgtactaa gtgatgcact caagagatac 420
 cctagaatgg acatcccaaa aattgccaga tccttctatg acttatttga acaaaaagt 480
 tatcacagaa gtttgttcat tgagtatggc aaagcattag gctcatcatc tacaggcagc 540
 aaagcagaaa gtctatttgt taatatattc atgcaagctt atggagccgg tcaacaagt 600
 ctaagggtggg ggggtcattgc caggtcatcc aacaatataa tgtaggata 650

<210> 15

<211> 650

<212> DNA

<213> human metapneumovirus

<220>

<221> gene

<222> (1)...(650)

<223> human metapneumovirus, strain Quebec Isolate No C-77807

<400> 15

gtatgctaaa catgctgatt acaaatatgc tgcagaaata ggaatacaat atattagcac 60
 agcttttagga tcagagagag tgcagcagat tctgaggaa tcaggcagtg aagtcacagt 120
 ggtctttaacc agaaccgtact cctctgggga agttaaaaa aataaaaggga gaagatttac 180
 agatgttaga catcacggg gtagagaaga gctgggtaga agagatagac aaagaagcaa 240
 ggaaaaacaat ggcaaccttg cttaaggaat catcaggtaa tatccacaa aatcagaggg 300
 cctcagcacc agacacacc ataactttat tatgtgtagg tgcttaata ttactaagc 360
 tagcatcaac catagaagt ggactagaga ccacagtcag aagggtctaac cgtgtactaa 420
 gtgatgcact caagagatac cctagaatgg acatcccaaa aattgccaga tccttctatg 480
 acttatttga acaaaaagt tatcacagaa gtttgttcat tgagtatggc aaagcattag 540
 gctcatcatc tacaggcagc aaagcagaaa gtctatttgt taatatattc atgcaagctt 600
 atggagccgg tcaacaagt ctaagggtggg ggggtcattgc caggtcatcc 650

<210> 16

<211> 650

<212> DNA

<213> human metapneumovirus

<220>

<221> gene

<222> (1)...(650)

<223> human metapneumovirus, strain Quebec Isolate No hmpv-C-78123

<400> 16

rtaggatcag agagagtgc gcagattcta aggaactcag gcagtgaagt ccaagtggtc 60
 ttaaccagaa cgtactctct ggggaaagt aaaaacaata aaggagaaga ttacagatg 120
 ttagacatac acggggtaga gaagagctgg gtagaagaga tagacaaaga agcaaggaaa 180
 acaatggcaa ccttgcttaa ggaatcatca ggtaatatcc cacaaaatca gaggccctca 240
 gcaccagaca caccataat ctattatgt gtaggtgcct taatatttac taaactagca 300
 tcaaccatag aagtgggact agagaccaca gtcagaagg ctaactgtgt actaagtgat 360
 gcactcaaga gataccctag aatggacatc ccaaaaattg ccagatcctt ctatgactta 420

10/52

tttgaacaaa	aagtgtatca	cagaagtttg	ttcattgagt	atggcaaaagc	attaggctca	480
tcatctacag	gcagcaaagc	agaaagtcta	tttgtaata	tattcatgca	agcttatgga	540
gccgggtcaa	caatgctaag	gtgggggggc	attgccagg	catccaacaa	tataatgtta	600
ggacatgtat	ccgtccaagc	tgagttaaaa	caggtcacag	aagtctatga		650

<210> 17
 <211> 650
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(700)
 <223> human metapneumovirus, strain Quebec Isolate No C-78132

<400> 17						60
cagaagagat	acaacagatt	ttgagaaact	caggtagtga	agttcagggt	gttttaacca	120
aaacatactc	cttagggaaa	ggcaaaaaa	gtaaagggga	agagctgcag	atgttagata	180
tacatggagt	ggaaaagagt	tggttagaag	aaatagacaa	agaggcaaga	aagacaatgg	240
taactttgct	taaggagtc	tcaggtaca	ccccacaaa	ccagagacct	tcagcaccag	300
acacaccaat	aattttatta	tggttaggtg	ccctaataat	cactaaacta	gcactacaa	360
tagaagttgg	attagagact	acagtttagaa	gagctaata	agtgttaagt	gatgcactca	420
aaagataccc	tagggtagat	ataccaaaga	ttgttagatc	tttttatgaa	ctatttgaac	480
aaaaagtgta	ctacagaagt	ttatttcattg	agtacggaaa	agcttttaggc	tcattctcaa	540
caggaagcaa	agcagaaagt	ttgtttgtaa	atatatttat	gcaagcttat	ggagctgggtc	600
aaacactgct	aaggtggggt	gtcattgcca	gatcatccaa	caacataatg	ctagggcatg	650
tatctgtgca	atctgaattg	aagcaagtta	cagaggttta	tgacttgggtg		

<210> 18
 <211> 650
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(650)
 <223> human metapneumovirus, strain Quebec Isolate No C-78163

<400> 18						60
ggatcagaaa	gagtacaaca	gattttgaga	aactcaggta	gtgaagttca	ggtgggttta	120
accaaaacat	actccttagg	gaaaggcaaa	aacagtaaa	gggaagagct	gcagatgtta	180
gatatacatg	gagtggaaaa	gagttgggtg	gaagaataag	acaaagaggc	aagaaagaca	240
atgggttaact	tgcttaagga	gtcatcagg	aacatccac	aaaaccagag	accttcagca	300
ccagacacac	caataatttt	attatgtgta	ggtgccctaa	tattcactaa	actagcatca	360
acaatagaag	ttggattaga	gactacagtt	agaagagcta	atagagtgtc	aagtgtatga	420
ctcaaaaagat	accctagggg	agatatacca	aagattgcta	gatcttttta	tgaactattt	480
gaacaaaag	tgtactacag	aagttttattc	attgagtagc	gaaaagcttt	aggctcatct	540
tcaacaggaa	gcaaaagcaga	aagttttgttt	gtaaatatat	ttatgcaagc	ttatggagct	600
ggtcaaacac	tgctaagggtg	gggtgtcatt	gccagatcat	ccaacacat	aatgctaggg	650
catgtatctg	tgcaatctga	attgaagcaa	gttacagagg	tttatgactt		

<210> 19
 <211> 600
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(600)
 <223> human metapneumovirus, strain Quebec Isolate No C-91205

<400> 19						60
atattagcac	agctctaggg	atcagagaga	gtacagcaga	ttctaagaaa	ctcaggcagt	120
gaagtccaag	tggttttaac	cagaacgtac	tccttgggga	aagttaaaaa	caacaaagga	180
gaagatttac	agatgttaga	catcacgga	gtagagaaaa	gctgggtgga	agagatagac	240
aaagaagcaa	gaaaaacaat	ggcaactttg	cttaaaagaat	catcaggcaa	tattccacaa	300
aatcagaggc	cttcagcacc	agacacacct	ataatcttat	tatgtgtagg	tgcccttaata	360
tttaccaaa	tagcatcaac	tatagaagtg	ggattagaga	ccacagtcag	aagagctaac	420
cgtgtactaa	gtgatgcact	caaaagatac	cctaggatgg	acataccaaa	aatcgctaga	

11/52

tctttctatg	acctatttga	acaaaaagtg	tattacagaa	gcttgttcat	tgagtatggc	480
aaagcattag	gctcatcctc	tacaggcagc	aaagcagaaa	gtttattcgt	taatatattc	540
atgcaagctt	acgggtgctg	tcaaacaatg	ctgaggtggg	gagtcattgc	caggtcattc	600

<210> 20
 <211> 700
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(700)
 <223> human metapneumovirus, strain Quebec Isolate No C-91364

<400> 20						
gattacaaat	atgctgctga	gataggaatg	caatatattt	gcacagctct	aggatcagaa	60
agagtacaac	agattttgag	aaactcaggt	agtgaagttc	aggtggttct	aaccaaaaca	120
tactccttag	ggaaaaggcaa	aaacagtaaa	ggggaagagc	tgcatagtgt	agatatcat	180
ggagtggaaa	agagttgggt	agaagaaaata	gacaaagagg	caagaaaagac	aatggtaact	240
tigcttaagg	agtcattcagg	taacatccca	caaaaccaga	gaccttcagc	accagacaca	300
ccaataat	tattatgtgt	agggtgccct	atattcacta	aactagcatc	aacaatagaa	360
gttgatttag	agactacagt	tagaagagct	aatagagtgc	taagtgatgc	actcaaaaga	420
taccttaggg	tagatatacc	aaagattgct	agatcttttt	atgaactatt	tgaacaaaaa	480
gtgtactaca	gaagtttatt	cattgagtac	ggaaaagctt	taggctcatt	ttcaacagga	540
agcaaaagcag	aaagtttgtt	tgtaaatata	tttatgcaag	cttatggagc	tggtcaaaaca	600
ctgctaaggt	ggggtgtcat	tgccagatca	tccaacaaca	taatgctagg	gcatgtatct	660
gtgcaatctg	aattgaagca	agttacagag	gtttatgact			700

<210> 21
 <211> 600
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(600)
 <223> human metapneumovirus, strain quebec isolate No C-91830

<400> 21						
cagctctagg	atcagaaaga	gtacaacaga	ttttgagaaa	ctcaggtagt	gaagttcagg	60
tggttctaac	caaaacatac	tccttaggga	aaggcaaaaa	cagtaaaggg	gaagagctgc	120
agatgttaga	tatacatgga	gtggaaaaga	gttggttaga	agaaatagac	aaagaggcaa	180
gaaagacaat	ggttaactttg	cttaaggagt	catcaggtaa	catccacaa	aaccagagac	240
cttcagcacc	agacacacca	ataattttat	tatgtgtagg	tgcccttaata	ttcactaaac	300
tagcatcaac	aatagaagtt	ggattagaga	ctacagttag	aagagcta	agagtgtctaa	360
gtgatgcact	caaaagatac	cctagggttag	atataccaaa	gattgctaga	tctttttatg	420
aactatttga	acaaaaagtg	tactacagaa	gcttattcat	tgagtacgga	aaagctttag	480
gctcatcttc	aacaggaagc	aaagcagaga	gtttgtttgt	aaatatattt	atgcaagctt	540
atggagctgg	tcaaacactg	ctaagggtgg	gtgtcattgc	cagatcatcc	aacaacataa	600

<210> 22
 <211> 650
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(650)
 <223> human metapneumovirus, strain Quebec isolate No hMPV-Q86

<400> 22						
atgctgatta	caaataatgct	gcagaaatag	gaatacaata	tattagcaca	gctctagggt	60
cagagaaagt	acagcagatt	ctaagaaact	caaggcagtga	agtccaagcg	gttttaacca	120
gaacgtactc	tttggggaaa	gttaaaaaa	ataaaggaga	agattacag	atgttagaca	180
tacacggagt	agaaaaaagc	tgggtggaag	agatagacaa	agaagcaaga	aaaaaatgg	240
caactttact	taaaagaatca	tcaggtaata	ttccacaaaa	tcagaggcct	tcagcaccag	300
acacacctat	aattttatta	tggttaggtg	ccttaatat	caccaaaacta	gcatcaacta	360
tagaagtggg	attagagacc	acagtcagaa	gagctaaccg	tgtactaagt	gatgcactca	420
aaagataccc	tagaatggac	ataccaaaaa	ttgctagatc	cittctatgac	ttatttgaac	480

12/52

aaaaagtgtgta tcacagaagt ttgttcattg agtatgggca aagcattagg ctcacccctc 540
 acagggcagca aagcagaaag tttattcgtc aatatattca tgcaagctta tgggtgctgg 600
 caaacaatgc tgaggtg999 ggggttatcgc cagggtcarct aacaatataa 650

<210> 23
 <211> 650
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(650)
 <223> human metapneumovirus, strain Quebec isolate No
 hMPV-Q97

<400> 23
 tagcacagct ctaggatcag agagagtaca gcagattcta agaaactcag gcagtgaagt 60
 ccaagtgggtt ttaaccagaa cgtactcctt ggggaaagt aaaaacaaca aaggagaaga 120
 tttacagatg ttagacatac acggagtaga gaaaagctgg gtggaagaga tagacaaaga 180
 agcaagaaaa acaatggcaa ctttgcttaa agaactatca ggcaatattc cacaaaatca 240
 gaggccttca gcaccagaca cacctataat cttattatgt gtagggtgct taatatttac 300
 caaactagca tcaactatag aagtgggatt agagaccaca gtcagaagag ctaaccgtgt 360
 actaagtgtg gcactcaaaa gataccctag gatggacata ccaaaaatcg ctatgctttt 420
 ctatgactta ttgaaacaaa aagtgtatta cagaagcttg ttcattgagt atggcaaaagc 480
 attaggctca tcctctacag gcagcaaaagc agaaagtttt attcgttaat atattcatgc 540
 aagcttacgg tgctgggtcaa acaatgctga ggtggggagt cattgcccagg tcatctaaca 600
 atataatggt aggacaigta tctgtccaag ctgaggttaa acaagtcacg 650

<210> 24
 <211> 741
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(741)
 <223> human metapneumovirus, strain Can 97-83

<400> 24
 agtgacaccc tcatcattgc aacaagaaat aacactattg tgtggagaaa ttctatatgc 60
 taagcatgct gattacaaat atgctgcaga aataggaata caatatatta gcacagctct 120
 aggatcagag agagtacagc agattctaag aaactcaggc agtgaagtcc aagtgggttt 180
 aaccagaacg tactccttgg ggaaggttaa aaacaacaaa ggagaagatt tacagatggt 240
 agacatacac ggagttagaga aaagctgggt ggaagagata gacaaagaag caagaaaaac 300
 aatggcaact ttgcttaaaag aatcatcagg caatatcca caaatcaga ggccttcagc 360
 accagacaca cctataatct tattatgtgt aggtgcctta atattacca aactagcatc 420
 aactatagaa gtgggattag agaccacagt cagaagagct aaccgtgtac taagtgtatg 480
 actcaaaaga taccctagga tggacatacc aaaaatcgt agatctttct atgatttatt 540
 tgaacaaaaa gtgtattaca gaagrttgtt cattgagtat ggcaagcat taggctcatc 600
 ctctacaggc agcaagcag aaagtttatt cgttaataata tcatgcaag cttacgggtg 660
 tgggtcaaaac atgctgaggt ggggagtcac tgccagggtca tctaacaata taatgttagg 720
 acatgtatct gtccaagctg a 741

<210> 25
 <211> 650
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(650)
 <223> human metapneumovirus, strain Can 98-75

<400> 25
 caatatattt gcacggctct aggatcagaa agagtacaac agattttaag aaattcaggc 60
 agtgaagtgc aggtggttct aaaccaagaca tacicttttag ggaaaggtta aaatagtaaa 120
 ggggaagagt tgcaaatggt agatatacat ggagtggaaa agagtgggt agaagaaata 180
 gacaaagagg caagaaaaac aatgggtgact ttgctaaagg aatcatcagg taacatccca 240
 caaaaccaga ggccttcagc accagacaca ccaataattt tattatgtgt aggtgcttta 300
 atattcacta aactagcatc aacaatagaa gttggactag agactacagt tagaagagct 360
 aacagagtgc taagtgtatg gctcaaaaga taccctaggg tagatatacc gaagattgct 420
 agatctttct atgaactatt tgagcagaaa gtgtattaca ggagtcattt cattgagtat 480

13/52

gggaaagcct taggctcatc ttcaacagga agcaaagcag aaagtgtgtt tgtaaatata 540
 ttatgcaag cttatggagc cgggtcaaca atgctaaggt ggggtgtcat tgccagatca 600
 tctaacaaca taatgctagg acatgtgtct gtgcaagctg aattgaagca 650

<210> 26
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Can 98-75

<400> 26
 atgtcttgga aagtgggtgat tatcatttcg ttactcataa cacctcagca cggactaaaa 60
 gaaagtatt tagaagaatc atgtagtact ataactgaag gataatctcag tgttttaaga 120
 acaggttggt acaccaatgt ctttacatta gaagttgggtg atgttgaaaa tcttacatgt 180
 actgatggac ctatcttaat caaaacagaa cttgacctaa ccaaaagtgc tctaagagaa 240
 ctcaaaaacag tttctgctga tcagtttagcg agagaagaac aaattgaaa tcccagacaa 300
 tcaaggtttg tcctaggtgc aatagctctt ggtgttgcca cagcagcagc agtcacagca 360
 ggcattgcga tagccaaaac cataaggctt gagagtgaag tgaatgcaat caaaggtgct 420
 ctcaaaaacaa ccaacgagggc agtatccaca ctaggaaatg gagtgcgagt cctagccacc 480
 gcagtaagag agctgaaaaga atttgtgagc aaaaacctga ctagtgcatt taacaagaac 540
 aaatgtgaca ttgctgacct gaagatggct gtcagcttca gtcaattcaa cagaagattc 600
 ctcaatgttg tgcggcagtt ttcagacaaat gcagggataa caccagcaat atcattggag 660
 ctaatgactg atgctgagct ggccagagct gtatcataca tgccaacatc tgcaggacag 720
 ataaaactaa tgttagagaa ccgtgcaatg gtaagaaga 759

<210> 27
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No C-67345

<400> 27
 atgtcttgga aagtgggtgat ctttttttca ttgctaataa cacctcaaca cggctctaaa 60
 gagagctacc tagaagaatc atgtatcact ataactgagg gataatcttag tgttctgaga 120
 acaggttggt ataccaacgt ttttacatta gaggtgggtg atgtagaana ccttacatgt 180
 tctgatggac ctatcttaat aaaaacagaa tttagatctga ccaaaagtgc actaagagag 240
 ctcaaaaacag tctctgctga ccaatrggca agagaggaac aaattgagaa tcccagacaa 300
 tctaggtttg ttctgggagc aatagcactc ggtgttgcaa cagcagctgc agtcacagca 360
 ggtgttgcaa ttgccaanaac catccggctt gagagtgaag tcacagcaat taagaatgcc 420
 ctcaaaaacga ccaatgaagc agtatctaca ttggggaaatg gagtctcagat gttggcaact 480
 gcagtgagag agctgaaaaga ctttgtgagc aagaatttaa ctctgtgcaat caacaaaaac 540
 aagtgcgaca ttgatgacct aaaaatggcc gtttagcttca gtcaattcaa cagaaggttt 600
 ctcaatgttg tgcggcaatt ttcagacaaat gctggaataa caccagcaat atctttggac 660
 ttaatgacag atgctgaact agccagggcc gtttccaaca tgccgacatc tgcaggacaa 720
 ataaaattga tgttggagaa ccgtgcatg gtaagaaga 759

<210> 28
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Can 97-83

<400> 28
 ataaaaatgt cttggaaagt ggtgatcatt ttttcattgc taataacacc tcaaacgggt 60
 cttaaagaga gctacctaga agaatcatgt agcactataa ctgagggata tcttagtgtt 120
 ctgaggacag gttggtatc caacgttttt acatttagagg tgggtgatgt agaaaaactt 180
 acatgttctg atggacctag cctaataaaa acagaattag atctgaccaa aagtgacta 240
 agagagctca aaacagctctc tgcagaccaa ttggcaagag aggaacaaat tgagaatccc 300
 agacaatcta ggtttgttctc agtagcaata gcactcgggt ttgcaacagc agctgcagtc 360

14/52

acagcaggtg	ttgcaattgc	caaaaccatc	cggcttgaga	gtgaagtcac	agcaattaag	420
aatgccctca	aaacgaccaa	tgaagcagta	tctacattgg	ggaatggagt	tcgagtgttg	480
gcaactgcag	tgagagagct	gaaagacttt	gtgagcaaga	atttaactcg	tgcaatcaac	540
aaaaacaagt	gcgacattga	tgacctaaaa	atggccggtta	gcttcagtc	attcaacaga	600
aggtttctaa	atgttgtgcg	gcaattttca	gacaatgctg	gaataacacc	agcaatatct	660
ttggacttaa	tgacagatgc	tgaactagcc	agggccggtt	ctaacatgcc	gacatctgca	720
ggacaataaa	aattgatgtt	ggagaaccgt	gcgatgggtg			759

<210> 29
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain quebec Isolate No
 C-68661

<400> 29						
atgtcttgga	aagtgggtgat	cattttttca	ttgttaataa	cacctcaaca	cggctcttaaa	60
gagagctatt	tagaagagtc	atgtagcact	ataactgaag	gatatactcag	tggtctgagg	120
acaggttggg	ataccaacgt	ttttacactg	gaggtaggtg	atgtagagaa	ccttacatgt	180
gctgatggac	ctagcttaat	aaaaacagaa	ttagacctga	ccaaaagtgc	actaagagag	240
ctcagaacag	ttttctgctga	tcaactggca	agagaggagc	aaattgagaa	tcccagacaa	300
tctagatttg	ttctaggagc	aatagcactc	ggtgttgcaa	cagcagctgc	agttacagca	360
ggtgttgcaa	ttgccaaaac	catccggctt	gaaagtgaag	taacagcaat	taagaatgcc	420
ctcaaaaaga	ccaatgaagc	agtatctaca	ttggggaaatg	gagttcgagt	gttggcaact	480
gcagtggagg	agctgaaaga	ttttgtgagc	aagaatctaa	cacgtgcaat	caacaaaaac	540
aaagtgcgaca	ttgctgacct	gaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
ctaaatgttg	tgccggcaatt	ttcagacaat	gctggaataa	caccagcaat	atccttggac	660
ttaatgacag	atgctgaact	agccagagct	gtttccaaca	tgccaacatc	tcaggacaa	720
ataaaaactga	tgttggagaa	ccgtgcaatg	gtaagaaga			759

<210> 30
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No
 C-73750

<400> 30						
atgtcttgga	aagtgggtgat	cattttttca	ttgttaataa	cacctcaaca	cggctcttaaa	60
gagagctatt	tagaagagtc	atgtagcact	ataactgaag	gatatactcag	tggtctgagg	120
acaggttggg	ataccaacgt	ttttacactg	gaggtaggtg	atgtagagaa	ccttacatgt	180
gctgatggac	ctagcttaat	aaaaacagaa	ttagacctga	ccaaaagtgc	actaagagag	240
ctcagaacag	ttttctgctga	tcaactggca	agagaggagc	aaattgagaa	tcccagacaa	300
tctagatttg	ttctaggagc	aatagcactc	ggtgttgcaa	cagcagctgc	agttacagca	360
ggtgttgcaa	ttgccaaaac	catccggctt	gaaagtgaag	taacagcaat	taagaatgcc	420
ctcaaaaaga	ccaatgaagc	agtatctaca	ttggggaaatg	gagttcgagt	gttggcaact	480
gcagtggagg	aactgaaaga	ttttgtgagc	aagaatctaa	cacgtgcaat	caacaaaaac	540
aaagtgcgaca	ttgctgacct	gaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
ctaaatgttg	tgccggcaatt	ttcagacaat	gctggaataa	caccagcaat	atccttggac	660
ttaatgacag	atgctgaact	agccagagct	gtttccaaca	tgccaacatc	tcaggacaa	720
ataaaaactga	tgttggagaa	ccgtgcaatg	gtaagaaga			759

<210> 31
 <211> 759
 <212> pNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No
 C-74549

<400> 31

15/52

atgtcttggg	aagtgggtgat	catttttttca	ttgctaataa	cacctcaaca	cggtctttaa	60
gagagctacc	tagaagaatc	atgtagcact	ataactgagg	gatatcttag	tggtctgagg	120
acagggttgg	ataccaacgt	ttttacatta	gaggtgggtg	atgtagaaaa	ccttacatgt	180
tctgatggac	ctagcctaata	aaaaacagaa	ttagatctga	ccaaaagtgc	actaagagag	240
ctcaaaacag	tctctgctga	ccaattggca	agagaggaa	aaattgagaa	tcccagacaa	300
tctagggttg	ttctaggagc	aatagcactc	ggtgttgcaa	cagcagctgc	agtcacagca	360
ggtgttgcaa	ttgccaaaac	catccggctt	gagagtgaag	tcacagcaat	taagaatgcc	420
ctcaaaacga	ccaatgaagc	agtatctaca	ttggggaatg	gagttcgagt	gttggcaact	480
gcagtggagag	agctgaaaga	ctttgtgagc	aagaatttaa	ctcgtgcaat	caacaaaaac	540
aagtgcgaca	ttgatgacct	aaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
ctaaatgrrg	tgccggcaatt	ttcagacaat	gctggaataa	caccagcaat	atccttggac	660
ttaatgacag	atgctgaact	agccagggcc	gtttctaaca	tgccgacatc	tgaggacaa	720
ataaaattga	tggtggagaa	ccgcgcgatg	gtaagaaga			759

<210> 32
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No C-74783

<400> 32						
atgtcttggg	aagtgggtgat	catttttttca	ttgctaataa	cacctcaaca	cggtctttaa	60
gagagctacc	tagaagaatc	atgtagcact	ataactgagg	gatatcttag	tggtctgagg	120
acagggttgg	ataccaacgt	ttttacatta	gaggtgggtg	atgtagaaaa	ccttacatgt	180
tctgatggac	ctagcctaata	aaaaacagaa	ttagatctga	ccaaaagtgc	actaagagag	240
ctcaaaacag	tctctgctga	ccaattggca	agagaggaa	aaattgagaa	tcccagacaa	300
tctagggttg	ttctaggagc	aatagcactc	ggtgttgcaa	cagcagctgc	agtcacagca	360
ggtgttgcaa	ttgccaaaac	catccggctt	gagagtgaag	tcacagcaat	taagaatgcc	420
ctcaaaacga	ccaatgaagc	agtatctaca	ttggggaatg	gagttcgagt	gttggcaact	480
gcagtggagag	agctgaaaga	ctttgtgagc	aagaatttaa	ctcgtgcaat	caacaaaaac	540
aagtgcgaca	ttgatgacct	aaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
ctaaatgrrg	tgccggcaatt	ttcagacaat	gctggaataa	caccagcaat	atccttggac	660
ttaatgacag	atgctgaact	agccagggcc	gtttctaaca	tgccgacatc	tgaggacaa	720
ataaaattga	tggtggagaa	ccgcgcgatg	gtaagaaga			759

<210> 33
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No C-76837

<400> 33						
atgtcttggg	aagtgggtgat	catttttttca	ttgttaataa	cacctcaaca	cggtctttaa	60
gagagctatt	tagaagagtc	atgtagcact	ataactgaag	gatatctcag	tggtctgagg	120
acagggttgg	ataccaacgt	ttttacactg	gaggttaggtg	atgtagagaa	ccttacatgt	180
gctgatggac	ctagcctaata	aaaaacagaa	ttagatctga	ccaaaagtgc	actaagagag	240
ctcagaacag	tttctgctga	tcaactggca	agagaggagc	aaattgagaa	tcccagacaa	300
tctagatttg	ttctaggagc	aatagcactc	ggtgttgcaa	cagcagctgc	agttacagca	360
ggtgttgcaa	ttgccaaaac	catccggctt	gaaagtgaag	taacagcaat	taagaatgcc	420
ctcaaaaaga	ccaatgaagc	agtatctaca	ttggggaatg	gagttcgagt	gttggcaact	480
gcagtggagg	aactgaaaga	ttttgtgagc	aagaatctaa	cacgtgcaat	caacaaaaac	540
aagtgcgaca	ttgctgacct	gaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
ctaaatgrrg	tgccggcaatt	ttcagacaat	gctggaataa	caccacaaat	atccttggac	660
ttaatgacag	atgctgaact	agccagagct	gtttccaaca	tgccaacatc	tgaggacaa	720
ataaaaactga	tggtggagaa	ccgtgcaatg	gtaagaaga			759

<210> 34
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>

16/52

<221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No
 C-77748

```

<400> 34
atgtcttggga aagtgggtgat catTTTTTtca ttgttaataa cacttcaaca cggctcttaaa 60
gagagctatt tagaagagtc atgttagcact ataactgaag gatattctcag tgttctgagg 120
acagggttggg ataccaacgt ttttacctg aaggtaggtg atgtagagaa ccttacatgt 180
gctgatggac ctactttaat aaaaacagaa ttagacctga ccaaaagtgc actaagagaa 240
ctcagaacag tttctgctga tcaactggca agagaggagc aaattgagaa tcccagacaa 300
tctagatttg ttctaggagc aatagcactc ggtgttgcaa cagcagctgc agttacagca 360
ggtgttgcaa ttgccaaaac catccggctt gaaagtgaag taacagcaat taagaatgcc 420
ctcaaaaaga ccaatgaagc agtatctaca ttggggaaatg gagttcgagt gttggcaact 480
gcagtgaggg agctgaaaga ttttgtgagc aagaatctaa cacgtgcaat caacaaaaac 540
aagtgcgaca ttgctgacct gaaaatggcc gttagcttca gtcaattcaa cagaagggtt 600
ctaaatgttg tgcggcaatt ttcagacaat gctggaataa caccagcaat atccttggac 660
ttaatgacag atgctgaact agccagagct gtttccaaca tgccaacatc tgcaggacaa 720
ataaaactga tgttggagaa ccgtgcaatg gtaagaaga 759
  
```

<210> 35
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No
 C-77807

```

<400> 35
atgtcttggga aagtgggtgat catTTTTTtca ttgttaataa cacttcaaca cggctcttaaa 60
gagagctatt tagaagagtc atgttagcact ataactgaag gatattctcag tgttctgagg 120
acagggttggg ataccaacgt ttttacctg aaggtaggtg atgtagagaa ccttacatgt 180
gctgatggac ctactttaat aaaaacagaa ttagacctga ccaaaagtgc actaagagaa 240
ctcagaccag tttctgctga tcaactggca agagaggagc aaattgagaa tcccagacaa 300
tctagatttg ttctaggagc aatagcactc ggtgttgcaa cagcagctgc agttacagca 360
ggtgttgcaa ttgccaaaac catccggctt gaaagtgaag taacagcaat taagaatgcc 420
ctcaaaaaga ccaatgaagc agtatctaca ttggggaaatg gagttcgagt gttggcaact 480
gcagtgaggg aactgaaaga ttttgtgagc aagaatctaa cacgtgcaat caacaaaaac 540
aagtgcgaca ttgctgacct gaaaatggcc gttagcttca gtcaattcaa cagaagggtt 600
ctaaatgttg tgcggcaatt ttcagacaat gctggaataa caccagcaat atccttggac 660
ttaatgacag atgctgaact agccagagct gtttccaaca tgccaacatc tgcaggacaa 720
ataaaactga tgttggagaa ccgtgcaatg gtaagaaga 759
  
```

<210> 36
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No
 C-78132

```

<400> 36
atgtcttggga aagtgggtgat catTTTTTtca ttgttaataa cacttcaaca cggctcttaaa 60
gagagctatt tagaagagtc atgttagcact ataactgaag gatattctcag tgttctgagg 120
acagggttggg ataccaacgt ttttacctg aaggtaggtg atgtagagaa ccttacatgt 180
gctgatggac ctactttaat aaaaacagaa ttagacctga ccaaaagtgc actaagagaa 240
ctcagaacag tttctgctga tcaactggca agagaggagc aaattgagaa tcccagacaa 300
tctagatttg ttctaggagc aatagcactc ggtgttgcaa cagcagctgc agttacagca 360
ggtgttgcaa ttgccaaaac catccggctt gaaagtgaag taacagcaat taagaatgcc 420
ctcaaaaaga ccaatgaagc agtatctaca ttggggaaatg gagttcgagt gttggcaact 480
gcagtgaggg aactgaaaga ttttgtgagc aagaatctaa cacgtgcaat caacaaaaac 540
aagtgcgaca ttgctgacct gaaaatggcc gttagcttca gtcaattcaa cagaagggtt 600
ctaaatgttg tgcggcaatt ttcagacaat gctggaataa caccagcaat atccttggac 660
ttaatgacag atgctgaact agccagagct gtttccaaca tgccaacatc tgcaggacaa 720
ataaaactga tgttggagaa ccgtgcaatg gtaagaaga 759
  
```

17/52

<210> 37
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No
 C-78163

<400> 37	atgtcttgga aagtggtgat cattttttca ttgttaataa cacctcaaca cggctctaaa	60
	gagagctatt tagaagagtc atgtagcact ataactgaag gatattctcag tgttctgagg	120
	acaggttggg ataccaacgt ttttacctg gaggtagggt atgtagagaa cttacatgt	180
	gctgatggac ctactttaat aaaaacagaa ttagatctga ccaaaagtgc actaagagag	240
	ctcagaacag tttctgctga tcaactggca agagaggagc aaattgagaa tcccagacaa	300
	tctagatttg ttctaggagc aatagcactc ggtgttgcaa cagcagctgc agttacagca	360
	ggtgttgcaa ttgccaaaac catccggctt gaaagtgaag taacagcaat taagaatgcc	420
	ctcaaaaaga ccaatgaagc agtatctaca ttgggggaatg gagttcgagt gttggcaact	480
	gcagtgaagg aactgaaaaga ttttgtgagc aagaatctaa cacgtgcaat caacaaaaac	540
	aagtgcgaca ttgctgacct gaaaatggcc gttagcttca gtcaattcaa cagaagggtt	600
	ctaaatgttg tgcggcaatt ttcagacaat gctggaataa caccagcaat atccttggac	660
	ttaatgacag atgctgaact agccagagct gtttccaaca tgccaacatc tgcaggacaa	720
	ataaaaactga tgttggagaa ccgtgcaatg gtaagaaga	759

<210> 38
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No
 C-78406

<400> 38	atgtcttgga aagtggtgat cattttttca ttgttaataa cacctcaaca cggctctaaa	60
	gagagctatt tagaagagtc atgtagcact ataactgaag gatattctcag tgttctgagg	120
	acaggttggg ataccaacgt ttttacctg gaggtagggt atgtagagaa cttacatgt	180
	gctgatggac ctactttaat aaaaacagaa ttagatctga ccaaaagtgc actaagagag	240
	ctcagaacag tttctgctga tcaactggca agagaggagc aaattgagaa tcccagacaa	300
	tctagatttg ttctaggagc aatagcactc ggtgttgcaa cagcagctgc agttacagca	360
	ggtgttgcaa ttgccaaaac catccggctt gaaagtgaag taacagcaat taagaatgcc	420
	ctcaaaaaga ccaatgaagc agtatctaca ttgggggaatg gagttcgagt gttggcaact	480
	gcagtgaagg aactgaaaaga ttttgtgagc aagaatctaa cacgtgcaat caacaaaaac	540
	aagtgcgaca ttgctgacct gaaaatggcc gttagcttca gtcaattcaa cagaagggtt	600
	ctaaatgttg tgcggcaatt ttcagacaat gctggaataa caccagcaat atccttggac	660
	ttaatgacag atgctgaact agccagagct gtttccaaca tgccaacatc tgcaggacaa	720
	ataaaaactga tgttggagaa ccgtgcaatg gtaagaaga	759

<210> 39
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No
 hmpv-174

<400> 39	atgtcttgga aagtggtgat cattttttca ttgttaataa cacctcaaca cggctctaaa	60
	gagagctatt tagaagagtc atgtagcact ataactgaag gatattctcag tgttctgagg	120
	acaggttggg ataccaacgt ttttacctg gaggtagggt atgtagagaa cttacatgt	180
	gctgatggac ctactttaat aaaaacagaa ttagacttga ccaaaagtgc actaagagag	240
	ctcagaacag tttctgctga tcaactggca agagaggagc aaattgagaa tcccagacaa	300
	tctagatttg ttctaggagc aatagcactc ggtgttgcaa cagcagctgc agttacagca	360
	ggtgttgcaa ttgccaaaac catccggctt gaaagtgaag taacagcaat taagaatgcc	420
	ctcaaaaaga ccaatgaagc agtatctaca ttgggggaatg gagttcgagt gttggcaact	480

18/52

gcagtgagag	agctgaaaga	ttttgtgagc	aagaatctaa	cacgtgcaat	caacaaaaac	540
aagtgcgaca	ttgctgacct	gaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
ctaaatgttg	tgcggaactt	ttcagacaat	gctggataaa	caccagcaat	atccttggac	660
ttaatgacag	atgctgaact	agccagagct	gtttccaaca	tgccaacatc	tgaggagaaa	720
ataaaactga	tggtggagaa	ccgtgcaatg	gtaagaaga			759

<210> 40
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No hmpv-193

<400> 40	atgtcttggg	aagtgggtgat	cattttttca	ttgttaataa	cacctcaaca	cggtctttaa	60
	gagagctatt	tagaagagtc	atgtagcact	ataactgaag	gatattctcag	tggtctgagg	120
	acagggttgg	ataccaacgt	ttttacactg	gaggttaggtg	atgtagagaa	ccttacatgt	180
	gctgatggac	ctagcttaat	aaaaacagaa	ttagacctga	ccaaaagtgc	actaagagag	240
	ctcagaacag	tttctgctga	tcaactggca	agagaggagc	aaattgagaa	tcccagacaa	300
	tctagatttg	ttctaggagc	aatagcactc	ggtgttgcaa	cagcagctgc	agttacagca	360
	ggtgttgcaa	ttgccaaaac	catccggctt	gaaagtgaag	taacagcaat	taagaatgcc	420
	ctcaaaaaga	ccaatgaagc	agtattctaca	ttggggaaatg	gagttcaggt	gttggcaact	480
	gcagtgagag	agctgaaaga	ttttgtgagc	aagaatctaa	cacgtgcaat	caacaaaaac	540
	aagtgcgaca	ttgctgacct	gaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
	ctaaatgttg	tgcggaactt	ttcagacaat	gctggaataa	caccagcaat	atccttggac	660
	ttaatgacag	atgctgaact	agccagagct	gtttccaaca	tgccaacatc	tgaggagaaa	720
	ataaaactga	tggtggagaa	ccgtgcaatg	gtaagaaga			759

<210> 41
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No hmpv-194

<400> 41	atgtcttggg	aagtgggtgat	cattttttca	ttgttaataa	cacctcaaca	cggtttttaa	60
	gagagctatt	tagaagagtc	atgtagcact	ataactgaag	gatattctcag	tggtctgagg	120
	acagggttgg	ataccaacgt	ttttacactg	gaggttaggtg	atgtagagaa	ccttacatgt	180
	gctgatggac	ctagcttaat	aaaaacagaa	ttagacctga	ccaaaagtgc	actaagagag	240
	ctcagaacag	tttctgctga	tcaactggca	agagaggagc	aaattgagaa	tcccagacaa	300
	tctagatttg	ttctaggagc	aatagcactc	ggtgttgcaa	cagcagctgc	agttacagca	360
	ggtgttgcaa	ttgccaaaac	catccggctt	gaaagtgaag	taacagcaat	taagaatgcc	420
	ctcaaaaaga	ccaatgaagc	agtattctaca	ttggggaaatg	gagttcaggt	gttggcaact	480
	gcagtgagag	agctgaaaga	ttttgtgagc	aagaatctaa	cacgtgcaat	caacaaaaac	540
	aagtgcgaca	ttgctgacct	gaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
	ctaaatgttg	tgcggaactt	ttcagacaat	gctggaataa	caccagcaat	atccttggac	660
	ttaatgacag	atgctgaact	agccagagct	gtttccaaca	tgccaacatc	tgaggagaaa	720
	ataaaactga	tggtggagaa	ccgtgcaatg	gtaagaaga			759

<210> 42
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No hmpv-198

<400> 42	atgtcttggg	aagtgggtgat	cattttttca	ttgttaataa	cacctcaaca	cggtctttaa	60
	gagagctatt	tagaagagtc	atgtagcact	ataactgaag	gatattctcag	tggtctgagg	120

19/52

acaggttggt	ataccaacgt	ttttacactg	gaggtaggtg	atgttagagaa	ccttacatgt	180
gctgatggac	ctagcttaat	aaaaacagaa	tttagacctga	ccaaaagtgc	actaagagag	240
ctcagaacag	tttctgctga	tcaactggca	agagaggagc	aaattgagaa	tcccagacaa	300
tctagatttg	ttctaggagc	aatagcactc	ggtgttgcaa	cagcagctgc	agttacagca	360
ggtgttgcaa	ttgccaaaac	catccggctt	gaaagtgaag	taacagcaat	taagaatgcc	420
ctcaaaaaga	ccaatgaagc	agtatctaca	ttggggaaatg	gagttcgagt	gtrggcaact	480
gcagtgagag	agctgaaaga	ttttgtgagc	aagaatctaa	cacgtgcaat	caacaaaac	540
aagtgcgaca	ttgctgacct	gaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
ctaaatgttg	tgctggcaatt	ttcagacaat	gctgggataa	caccagcaat	atccttggac	660
ttaatgacag	atgctgaact	agccagagct	gtttccaaca	tgccaacatc	tgaggacaa	720
ataaaaactga	tgtrggagaa	ccgtgcaatg	gtaagaaga			759

<210> 43
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No
 hmpv-217

<400> 43	atgtcttggg	aagtgggtgat	cattttttca	ttgtaataa	cacctcaaca	cggtcttaaa	60
	gagagctatt	tagaagagtc	atgtagcact	ataactgaag	gatatctcag	tggtctgagg	120
	acaggttggt	ataccaacgt	ttttacactg	gaggtaggtg	atgttagagaa	ccttacatgt	180
	gctgatggac	ctagcttaat	aaaaacagaa	tttagacctga	ccaaaagtgc	actaagagag	240
	ctcagaacag	tttctgctga	tcaactggca	agagaggagc	aaattgagaa	tcccagacaa	300
	tctagatttg	ttctaggagc	aatagcactc	ggtgttgcaa	cagcagctgc	agttacagca	360
	ggtgttgcaa	ttgccaaaac	catccggctt	gaaagtgaag	taacagcaat	taagaatgcc	420
	ctcaaaaaga	ccaatgaagc	agtatctaca	ttggggaaatg	gagttcgagt	gtrggcaact	480
	gcagtgagag	agctgaaaga	ttttgtgagc	aagaatctaa	cacgtgcaat	caacaaaac	540
	aagtgcgaca	ttgctgacct	gaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
	ctaaatgttg	tgctggcaatt	ttcagacaat	gctgggataa	caccagcaat	atccttggac	660
	ttaatgacag	atgctgaact	agccagagct	gtttccaaca	tgccaacatc	tgaggacaa	720
	ataaaaactga	tgtrggagaa	ccgtgcaatg	gtaagaaga			759

<210> 44
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No
 hmpv-208

<400> 44	atgtcttggg	aagtgggtgat	cattttttca	ttgtaataa	cacctcaaca	cggtcttaaa	60
	gagagctatt	tagaagagtc	atgtagcact	ataactgaag	gatatctcag	tggtctgagg	120
	acaggttggt	ataccaacgt	ttttacactg	gaggtaggtg	atgttagagaa	ccttacatgt	180
	gctgatggac	ctagcttaat	aaaaacagaa	tttagacctga	ccaaaagtgc	actaagagag	240
	ctcagaacag	tttctgctga	tcaactggca	agagaggagc	aaattgagaa	tcccagacaa	300
	tctagatttg	ttctaggagc	aatagcactc	ggtgttgcaa	cagcagctgc	agttacagca	360
	ggtgttgcaa	ttgccaaaac	catccggctt	gaaagtgaag	taacagcaat	taagaatgcc	420
	ctcaaaaaga	ccaatgaagc	agtatctaca	ttggggaaatg	gagttcgagt	gtrggcaact	480
	gcagtgagag	agctgaaaga	ttttgtgagc	aagaatctaa	cacgtgcaat	caacaaaac	540
	aagtgcgaca	ttgctgacct	gaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
	ctaaatgttg	tgctggcaatt	ttcagacaat	gctgggataa	caccagcaat	atccttggac	660
	ttaatgacag	atgctgaact	agccagagct	gtttccaaca	tgccaacatc	tgaggacaa	720
	ataaaaactga	tgtrggagaa	ccgtgcaatg	gtaagaaga			759

<210> 45
 <211> 759
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(759)

20/52

<223> human metapneumovirus, strain Quebec Isolate No
hMPV-223

<400> 45

atgtcttggg	aagtgggtgat	catttttttca	ttgttaataa	cacctcaaca	cggtctttaa	60
gagagctatt	tagaagagtc	atgttagcact	ataactgaag	gatatctcag	tggtctgagg	120
acaggttggg	ataccaacgt	ttttacactg	gaggttaggtg	atgtagagaa	ccttacatgt	180
gctgatggac	ctagcttaat	aaaaacagaa	ttagacctga	ccaaaagtgc	actaagagag	240
ctcagaacag	tttctgctga	tcaactggca	agagaggagc	aaattgagaa	tcccagacaa	300
tctagatttg	ttctaggagc	aatagcactc	gggtgttgcaa	cagcagctgc	agttacagca	360
gggtgttgcaa	ttgccaaaac	catccggcct	gaaagtgaag	taacagcaat	taagaatgcc	420
ctcaaaaaga	ccaatgaagc	agtatctaca	ttgggggaatg	gagttcgagt	gttggcaact	480
gcagtgagag	agctgaaaga	ttttgtgagc	aagaatctaa	cacgtgcaat	caacaaaaac	540
aagtgcgaca	ttgctgacct	gaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
ctaaatgttg	tgccgcaatt	ttcagacaat	gctgggaata	caccagcaat	atccttggac	660
ttaatgacag	atgctgaact	agccagagct	gtttccaaca	tgccaacatc	tcaggacaa	720
ataaaaactga	tggtggagaa	ccgtgcaatg	gtaagaaga			759

<210> 46

<211> 759

<212> DNA

<213> human metapneumovirus

<220>

<221> gene

<222> (1)...(759)

<223> human metapneumovirus, strain Quebec Isolate No
hMPV-228

<400> 46

atgtcttggg	aagtgggtgat	catttttttca	ttgttaataa	cacctcaaca	cggtctttaa	60
gagagctatt	tagaagagtc	atgttagcact	ataactgaag	gatatctcag	tggtctgagg	120
acaggttggg	ataccaacgt	ttttacactg	gaggttaggtg	atgtagagaa	ccttacatgt	180
gctgatggac	ctagcttaat	aaaaacagaa	ttagacctga	ccaaaagtgc	actaagagag	240
ctcagaacag	tttctgctga	tcaactggca	agagaggagc	aaattgagaa	tcccagacaa	300
tctagatttg	ttctaggagc	aatagcactc	gggtgttgcaa	cagcagctgc	agttacagca	360
gggtgttgcaa	ttgccaaaac	catccggcct	gaaagtgaag	taacagcaat	taagaatgcc	420
ctcaaaaaga	ccaatgaagc	agtatctaca	ttgggggaatg	gagttcgagt	gttggcaact	480
gcagtgagag	agctgaaaga	ttttgtgagc	aagaatctaa	cacgtgcaat	caacaaaaac	540
aagtgcgaca	ttgctgacct	gaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
ctaaatgttg	tgccgcaatt	ttcagacaat	gctgggaata	caccagcaat	atccttggac	660
ttaatgacag	atgctgaact	agccagagct	gtttccaaca	tgccaacatc	tcaggacaa	720
ataaaaactga	tggtggagaa	ccgtgcaatg	gtaagaaga			759

<210> 47

<211> 759

<212> DNA

<213> human metapneumovirus

<220>

<221> gene

<222> (1)...(759)

<223> human metapneumovirus, strain Quebec Isolate No
C-85885

<400> 47

atgtcttggg	aagtgggtgat	catttttttca	ttgttaataa	cacctcaaca	cggtctttaa	60
gagagctatt	tagaagagtc	atgttagcact	ataactgaag	gatatctcag	tggtctgagg	120
acaggttggg	ataccaacgt	ttttacactg	gaggttaggtg	atgtagagaa	ccttacatgt	180
gctgatggac	ctagcttaat	aaaaacagaa	ttagacctga	ccaaaagtgc	actaagagag	240
ctcagaacag	tttctgctga	tcaactggca	agagaggagc	aaattgagaa	tcccagacaa	300
tctagatttg	ttctaggagc	aatagcactc	gggtgttgcaa	cagcagctgc	agttacagca	360
gggtgttgcaa	ttgccaaaac	catccggcct	gaaagtgaag	taacagcaat	taagaatgcc	420
ctcaaaaaga	ccaatgaagc	agtatctaca	ttgggggaatg	gagttcgagt	gttggcaact	480
gcagtgagag	agctgaaaga	ttttgtgagc	aagaatctaa	cacgtgcaat	caacaaaaac	540
aagtgcgaca	ttgctgacct	gaaaatggcc	gttagcttca	gtcaattcaa	cagaagggtt	600
ctaaatgttg	tgccgcaatt	ttcagacaat	gctgggaata	caccagcaat	atccttggac	660
ttaatgacag	atgctgaact	agccagagct	gtttccaaca	tgccaacatc	tcaggacaa	720
ataaaaactga	tggtggagaa	ccgtgcaatg	gtaagaaga			759

<210> 48

<211> 759

21/52

<212> DNA
<213> human metapneumovirus

<220>
<221> gene
<222> (1)...(759)
<223> human metapneumovirus, strain Quebec Isolate No
C-60298

```

<400> 48
atgtcttggg aagtgggtgat catcttttca ttgttaataa caccitcaaca cggctcttaaa    60
gagagcttatt tagaagagtc atgtagcact ataactgaag gatattctcag tgttctgagg    120
acagggttggg ataccaacgt ttttaccactg gaggttaggtg atgtagagaa ccttacatgt    180
gctgatgggac ctagtcttaat aaaaacagaa ttagatctga ccaaaagtgc actaagagag    240
ctcagaacag ttctgctga tcaactggca agagaggagc aaattgagaa tcccagacaa    300
tctagatttg ttctaggagc aatagcactc ggtgttgcaa cagcagctgc agttacagca    360
ggtgttgcaa ttgccaaaac catccggctt gaaagtgaag taacagcaat taagaatgcc    420
ctcaaaaaga ccaatgaagc agtatctaca ttgggggaatg gagttcgagt gttggcaact    480
gcagtggagg aactgaaaga tttgtgagc aagaatctaa cagtgcaat caacaaaaac    540
aagtgcgaca ttgctgacct gaaaatggcc gttagcttca gtcaattcaa cagaaggttt    600
ctaaatgttg tgcggcaatt ttcagacaat gctgggaata caccagcaat atccttggac    660
ttaatgacag atgctgaact agccagagct gtttccaaca tgccaacatc tgcaggacaa    720
ataaaactga tgttggagaa ccgtgcaatg gtaagaaga    759

```

<210> 49
<211> 759
<212> DNA
<213> human metapneumovirus

<220>
<221> gene
<222> (1)...(759)
<223> human metapneumovirus, strain Quebec Isolate No
C-63778

```

<400> 49
atgtcttggg aagtgggtgat tatcatttcg ttactcataa caccitcagca cggactaaag    60
gaaagtttatt tagaagaatc atgtagtact ataactgaag gatattctcag tgttttaaga    120
acagggttggg acaccaatgt ctttacatta gaagttgggtg atgttgaaaa tcttacatgt    180
actgatgggac ctagtcttaat caaaacagaa cttgacctaa ccaaaagtgc tctaagagaa    240
ctcaaaaacag tttctgctga tcagttagcg agagaagaac aaattgaaaa tcccagacaa    300
tcaaggtttg tcttaggtgc aatagctctt ggtgttgcca cagcagcagc agtcacagca    360
ggcatttgcga tagccaaaac cataaggctt gagagtgaag tgaatgcaat caaagggtgt    420
ctcaaaaacaa ccaacgaggc agtatccaca ctaggaaatg gagtgcgagt cctagccacc    480
gcagtaagag agctgaaaga attgtgagc aaaaacctga ctagtcaat taacaagaac    540
aaaatgtgaca ttgctgatct gaagatggct gtcagcttca gtcaattcaa cagaagattc    600
ctaaatgttg tgcggcagtt ttcagacaat gcagggataa caccagcaat atcattggac    660
ctaattgactg atgctgagct ggccagaact gtatcataca tgccaacatt tgcaggacag    720
ataaaactaa tgttagagaa ccgtgcaatg gtaagaaga    759

```

<210> 50
<211> 759
<212> DNA
<213> human metapneumovirus

<220>
<221> gene
<222> (1)...(759)
<223> human metapneumovirus, strain Quebec Isolate No
C-64199

```

<400> 50
atgtcttggg aagtgggtgat tatcatttcg ttactcataa cacttcagca cggactaaaa    60
gaaagtttatt tagaagaatc atgtagtact ataactgaag gatattctcag tgttttaaga    120
acagggttggg acaccaatgt ctttacatta gaagttgggtg atgttgaaaa tcttacatgt    180
actgatgggac ctagtcttaat caaaacagaa cttgacctaa ccaaaagtgc tctaagagaa    240
ctcaaaaacag tttctgctga tcagttagcg agagaagaac aaattgaaaa tcccagacaa    300
tcaaggtttg tcttaggtgc aatagctctt ggtgttgcca cagcagcagc agtcacagca    360
ggcatttgcga tagccaaaac cataaggctt gagagtgaag tgaatgcaat caaagatgct    420
ctcaaaaacaa ccaacgaggc agtatccaca ctaggaaatg gagtgcgagt cctagccacc    480
gcagtaagag agctgaaaga attgtgagc aaaaacctga ctagtcaat taacaagaac    540
aaaatgtgaca ttgctgatct gaagatggct gtcagcttca gtcaattcaa cagaagattc    600

```

22/52

ctaaatgttg	tgccggcagtt	ttcagacaat	gcaggggataa	caccagcaat	atcattggac	660
ctaattgactg	atgctgagct	ggccagagct	gtatcatata	tgccaacatc	tcaggagacag	720
ataaaaactaa	tgtttagagaa	ccgtgcaatg	gtaagaaga			759

<210> 51
 <211> 759
 <212> DNA
 <213> human metapneumovirus
 <220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No C-64200

<400> 51						
atgtcttgga	aagtgggtgat	tatcatttcg	ttactcataa	cacctcagca	cggactaaaa	60
gaaagtatt	tagaagaatc	atgtagtact	ataactgaag	gatattctcag	tgttttaaga	120
acaggttgg	acaccaatgt	ctttacatta	gaagttgggtg	atgttgaaaa	tcttacatgt	180
actgatggac	ctagcttaat	caaaaacagaa	cttgacctaa	ccaaaagtgc	tctaagagaa	240
ctcaaaacag	tttctgctga	tcagttagcg	agagaagaac	aaattgaaaa	tcccagacaa	300
tcaaggtttg	tcctaggtgc	aatagctctt	gggtgttgcca	cagcagcagc	agtcacagca	360
ggcattgcca	tagccaaaac	cataaggcct	gagagtgaag	tgaatgcaat	caaaggtgct	420
ctcaaaaacaa	ccaacgaggc	agtatccaca	ctaggaaatg	gagtgcgagt	cctagccacc	480
gcagtaagag	agctgaaaga	atttgtgagc	aaaaacctga	ctagtgcatt	taacaagaac	540
aaatgtgaca	ttgctgatct	gaagatggct	gtcagcttca	gtcaattcaa	cagaagattc	600
ctaaatgttg	tgccggcagtt	ttcagacaat	gcaggggataa	caccagcaat	atcattggac	660
ctaattgactg	atgctgagct	ggccagagct	gtatcatata	tgccaacatc	tcaggagacag	720
ataaaaactaa	tgtttagagaa	ccgtgcaatg	gtaagaaga			759

<210> 52
 <211> 759
 <212> DNA
 <213> human metapneumovirus
 <220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No C-64536

<400> 52						
atgtcttgga	aagtgggtgat	tatcatttcg	ttactcataa	cacctcagca	cggactaaaa	60
gaaagtatt	tagaagaatc	atgtagtact	ataactgaag	gatattctcag	tgttttaaga	120
acaggttgg	acaccaatgt	ctttacatta	gaagttgggtg	atgttgaaaa	tcttacatgt	180
actgatggac	ctagcttaat	caaaaacagaa	cttgacctaa	ccaaaagtgc	tctaagagaa	240
ctcaaaacag	tttctgctga	tcagttagcg	agagaagaac	aaattgaaaa	tcccagacaa	300
tcaaggtttg	tcctaggtgc	aatagctctt	gggtgttgcca	cagcagcagc	agtcacagca	360
ggcattgcca	tagccaaaac	cataaggcct	gagagtgaag	tgaatgcaat	caaaggtgct	420
ctcaaaaacaa	ccaacgaggc	agtatccaca	ctaggaaatg	gagtgcgagt	cctagccacc	480
gcagtaagag	agctgaaaga	atttgtgagc	aaaaacctga	ctagtgcatt	taacaagaac	540
aaatgtgaca	ttgctgatct	gaagatggct	gtcagcttca	gtcaattcaa	cagaagattc	600
ctaaatgttg	tgccggcagtt	ttcagacaat	gcaggggataa	caccagcaat	atcattggac	660
ctaattgactg	atgctgagct	ggccagagct	gtatcatata	tgccaacatc	tcaggagacag	720
ataaaaactaa	tgtttagagaa	ccgtgcaatg	gtaagaaga			759

<210> 53
 <211> 759
 <212> DNA
 <213> human metapneumovirus
 <220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No C-73769

<400> 53						
atgtcttgga	aagtgggtgat	tatcatttcg	ttactcatac	cacctcagca	cggactaaaa	60
gaaagtatt	tagaagaatc	atgtagtact	ataactgaag	gatattctcag	tgttttaaga	120
ccaggttgg	acaccaatgt	ctttacatta	gaagttgggtg	atgttgaaaa	tcttacatgt	180
actgatggac	ctagcttaat	caaaaacagaa	cttgacctaa	ccaaaagtgc	tctaagagaa	240

23/52

ctcaaaacag	tttctgctga	tcagtttagcg	agagaagaac	aaattgaaaa	tcccagacaa	300
tcaaggtttg	tccttaggtgc	aatagctctt	gggtgttgcca	cagcagcagc	agtcacagca	360
ggcattgcga	tagccaaaac	cataaggctt	gagagtgaag	tgaatgcaat	caaaggtgct	420
ctcaaaacaa	ccaacgaggc	agtatccaca	ctaggaaatg	gagtgcgagt	cctagccacc	480
gcagtaagag	agttgaaaga	atttgtgagc	aaaaacctga	ctagtgcatt	taacaagaac	540
aaatgtgaca	ttgctgatct	gaagatggct	gtcagcttca	gtcaattcaa	cagaagattc	600
ctaaatgttg	tgcggcagtt	ttcagacaat	gcagggataa	caccagcaat	atcatttagac	660
ctaattgactg	atgctgagct	ggccagagct	gtatcatata	tgccaacatc	tgcaggacag	720
ataaaactaa	tgtttagagaa	ccgtgcaatg	gtaagaaga			759

<210> 54

<211> 759

<212> DNA

<213> human metapneumovirus

<220>

<221> gene

<222> (1)...(759)

<223> human metapneumovirus, strain Quebec Isolate No C-74501

<400> 54

atgtcttgga	aagtgggtgat	tatcatttcg	ttactcataa	cacttcagca	cggactaaaa	60
gaaggttatt	tagaagaatc	atgtagtact	ataactgaag	gatatctcag	tgttttaaga	120
acaggttggt	acaccaatgt	ctttacatta	gaagtgggtg	atgttgaaaa	tcttacatgt	180
actgatggac	ctagcttaat	caaaacagaa	cttgacctaa	ccaaaagtgc	tctaagagaa	240
ctcaaaacag	tttctgctga	tcagtttagcg	agagaagaac	aaattgaaaa	tcccagacaa	300
tcaaggtttg	tccttaggtgc	aatagctctt	gggtgttgcca	cagcagcagc	agtcacagca	360
ggcattgcga	tagccaaaac	cataaggctt	gagagtgaag	tgaatgcaat	caaaggtgct	420
ctcaaaacaa	ccaacgaggc	agtatccaca	ctaggaaatg	gagtgcgagt	cctagccact	480
gcagtaagag	agctgaaaga	atttgtgagc	aaaaacctga	ctagtgcatt	taacaagaac	540
aaatgtgaca	ttgctgatct	gaagatggct	gtcagcttca	gtcaattcaa	cagaagattc	600
ctaaatgttg	tgcggcagtt	ttcagacaat	gcagggataa	caccagcaat	atcatttagac	660
ctaattgactg	atgctgagct	ggccagagct	gtatcatata	tgccaacatc	tgcaggacag	720
ataaaactaa	tgtttagagaa	ccgtgcaatg	gtaagaaga			759

<210> 55

<211> 759

<212> DNA

<213> human metapneumovirus

<220>

<221> gene

<222> (1)...(759)

<223> human metapneumovirus, strain Quebec Isolate No C-78429

<400> 55

atgtcttgga	aagtgggtgtt	catcatttcg	ttactcataa	caccccagca	cgggctaaag	60
gagagtttatt	tggaagaatc	atgcagtact	ataactgagg	gatacctcag	tgttttaaga	120
acaggttggt	acattaatgt	cttcacatta	gaagtgggtg	atgttgaaaa	tcttacatgt	180
actgatggac	ctagcttaat	caaaacagaa	cttgacctaa	ccaaaagtgc	tttaagggaa	240
ctcaaaacag	tttctgctga	tcagtttagcg	agagaggagc	aaattgaaaa	tcccagacaa	300
tcaagatttg	tccttaggtgc	aatagctctc	ggagrtgcta	cagcagcagc	agtcacagca	360
ggcattgcaa	tagccaaaac	cataaggctt	gagagtgaag	tgaatgcaat	taaaggtgct	420
ctcaaaacaa	ctaattgaagc	agtatccaca	ttaggaaatg	gtgtgcgggt	cctagccact	480
gcagtgagag	agctaaaaga	atttgtgagc	aaaaatctga	ctagtgcatt	caacagggaac	540
aaatgtgaca	ttgctgatct	gaagatggct	gtcagcttca	gtcaattcaa	cagaagattt	600
ctcaatgttg	tgcggcagtt	ttcagacaat	gcagggataa	caccagcaat	atcattggac	660
ctaattgactg	atgctgaatt	ggccagagct	gtatcatata	tgccaacatc	tgcaggacag	720
ataaaactga	tgtrtgagaa	ccgtgcaatg	gtaagaaga			759

<210> 56

<211> 759

<212> DNA

<213> human metapneumovirus

<220>

<221> gene

<222> (1)...(759)

<223> human metapneumovirus, strain Quebec Isolate No C-78652

24/52

```

<400> 56
atgtcrttga aagtgggtgat catcatttcg ttactcataa caccctcagca cgggctaaga 60
gagagttatt tggagaagaatc atgcagtagt ataactgagg gatactctcag tgttttaaga 120
acgggctggg acactaatgt cttcacatta gaagttgggt atgttgaaaa tcttacatgt 180
actgatggac ctactttaat caaacacagaa cttgacctaa caaaaagtgc ttaaggga 240
ctcaaaacag tctctgctga tcagttaacg agagaggagc aaattgaaaa tcccagacaa 300
tcaagatttg tcttaggtgc aatagctctc ggagttgcta cagcagcagc agtcacagca 360
ggcattgcaa tagccaaaac cataaggctt gagagtgagg tgaatgcaat taaagggtgct 420
ctcaaaacaa ctaatgaagc agtatccaca ttaggaaatg gtgtgagggt cctagccact 480
gcagtggagc agctaaaaga atttgtgagc aaaaatctga ctagtgcatt caacaggaaac 540
aaatgtgaca ttgctgatct gaagatggct gtcagcttca gtcaattcaa cagaagattt 600
ctaaatgttg tgcggcagtt ttccagacaat gcagggataa caccagcaat atcattggac 660
ctaattgactg atgctgaatt ggccagagct gtatcatata tgccaacatc tgcaggacag 720
ataaaactga tgttgagaa ccgtgcaatg gtaagaaga 759

```

```

<210> 57
<211> 759
<212> DNA
<213> human metapneumovirus

<220>
<221> gene
<222> (1)...(759)
<223> human metapneumovirus, strain Quebec Isolate No
hmpv-118

```

```

<400> 57
atgtccttga aagggggtgat tatcatttcg ttactcataa caccctcagca cggactaaaa 60
gaaagttatt tagaagaatc atgtagtact ataactgaag gatactctcag tgttttaaga 120
acagggttgg acaccaatgt ctttacatta gaagttgggt atgttgaaaa tcttacatgt 180
actgatggac ctactttaat caaacacagaa cttgacctaa caaaaagtgc tctgagagaa 240
ctcaaaacag tttctgctga tcagttaacg agagaagaac aaattgaaaa tcccagacaa 300
tcaagggttg tcttaggtgc aatagctctt ggtgttgcca cagcagcagc agtcacagca 360
ggcattgcga tagccaaaac cataaggctt gagagtgaag tgaatgcaat caaagggtgct 420
ctcaaaacaa ccaacgaggc agtatccaca ctaggaaatg gagtgcgagt cctagccacc 480
gcagtaagag agctgaaaga atttgtgagc aaaaacctga ccagtgcatt taacaagaac 540
aaatgtgaca ttgctgatct gaagatggct gtcagcttca gtcaattcaa cagaagattc 600
ctaaatgttg tgcggcagtt ttccagacaat gcagggataa caccagcaat atcattagac 660
ctaattgactg atgctgagct ggccagagct gtatcatata tgccaacatc tgcagggcag 720
ataaaactaa tgttagagaa ccgtgcaatg gtaagaaga 759

```

```

<210> 58
<211> 759
<212> DNA
<213> human metapneumovirus

<220>
<221> gene
<222> (1)...(759)
<223> human metapneumovirus, strain Quebec Isolate No
hmpv-200

```

```

<400> 58
atgtccttga aagtgggtgat catcttttcg ttactcataa caccctcagca cggactaaaa 60
gaaagttatt tagaagaatc atgtagtact ataactgaag gatactctcag tgttttaaga 120
acagggttgg acaccaatgt ctttacatta gaagttgggt atgttgaaaa tcttacatgt 180
actgatggac ctactttaat caaacacagaa cttgacctaa caaaaagtgc tctaagagaa 240
ctcaaaacag tttctgctga tcagttaacg agagaagaac aaattgaaaa tcccagacaa 300
tcaagggttg tcttaggtgc aatagctctt ggtgttgcca cagcagcagc agtcacagca 360
ggcattgcga tagccaaaac cataaggctt gagagtgaag tgaatgcaat caaagggtgct 420
ctcaaaacaa ccaacgaggc agtatccaca ctaggaaatg gagtgcgagt cctagccacc 480
gcagtaagag agctgaaaga atttgtgagc aaaaacctga ccagtgcatt taacaagaac 540
aaatgtgaca ttgctgatct gaagatggct gtcagcttca gtcaattcaa cagaagattc 600
ctaaatgttg tgcggcagtt ttccagacaat gcagggataa caccagcaat atcattagac 660
ctaattgactg atgctgagct ggccagagct gtatcatata tgccaacatc tgcagggcag 720
ataaaactaa tgttagagaa ccgtgcaatg gtaagaaga 759

```

```

<210> 59
<211> 759
<212> DNA
<213> human metapneumovirus

```

25/52

<220>
 <221> gene
 <222> (1)...(759)
 <223> human metapneumovirus, strain Quebec Isolate No
 hmpv-176

<400> 59
 atgtcttgga aagtgggtgat tatcatttcg ttactcataa caccctcagca cggactaaaa 60
 gaaagttatt tagaagaatc atgtagtact ataactgaag gatattctag ttttctaaga 120
 acagggttgg acaccaatgt ctttacatta gaagttgggt atgttgaaaa tcttacatgt 180
 actgatggac ctatgcttaac caaacacagaa cttgacctaa ccaaaagtgc ttaagagaa 240
 ctcaaaacag tttctgctga tcagttagcg agagaagaac aaattgaaaa tcccagacaa 300
 tcaagggttg tcctagggtgc aatagctctc ggagttgcca cagcagcagc agtcacagca 360
 ggcatgtgcaa tagccaaaac tataaggctt gagagtgaag tgaatgcaat caaagggtgt 420
 ctcaaaacaa ccaatgaggc agtatcaaca ctaggaaatg gagtgcgggt cctagccact 480
 gcagtaagag agctgaaaga atttgtgagc aaaaacctga ctagtgcgat caacaagaac 540
 aaatgtgaca ttgctgattt gaagatggct gtcagcttca gtcagttcaa cagaagattc 600
 cttaaattgtg tgcggcagtt ttcagacaat gcagggataa caccagcaat atcattggac 660
 ctgatgactg atgtctgagc gccacagagc gtatcatata tgccaacatc tgcaggacag 720
 ataaaactaa tggtagagaa ccgtgcaatg gtaagaaga 759

<210> 60
 <211> 711
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(711)
 <223> human metapneumovirus, strain Quebec Isolate No
 hmpv-193

<400> 60
 atggaggtga aagtggagaa cattcgaaca atagatatgc tcaaagcaag agtgaaaaat 60
 cgtgtggcac gcagcaaatg ctttaaaaaat gcctctttga tcttaatagg aataactaca 120
 ttgagtatag ccttcaatat ctatctgac ataaactata caatgcaaga aaacacatcc 180
 gaatcagaac atcacaccag ctcatcacc atggaatcca gcagggaac tccaacggtc 240
 cccatggact acccagacac caatccaggc tcacagtatc caactcaaca gtccacagaa 300
 ggctccacac tctactttgc agcctcagca agctcaccag agacagaacc aacatcaaca 360
 ccaaacacaa caagccggcc gcccttcgtc gacacacaca caacaccacc aagtgaagc 420
 agaacaaaga caagtccggc agtccacaca aaaaacaatc caaggataag cctcagaaca 480
 cattccccac catggggcaat gacaaggacg gtccgaggaa ccaccactct ccgcacaagc 540
 agcacaagaa aaagaccgtc tacagcatca gtccaacctg acagcagcgc aacaaccac 600
 aaacacgaag aagcaagccc agtgagcccg caaacatctg caagcacagc aagaccataa 660
 aggaaggtca tggaggccag cacatcaaca acatacaatc aaactagtta a 711

<210> 61
 <211> 711
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(711)
 <223> human metapneumovirus, strain Quebec Isolate No
 hmpv-228

<400> 61
 atggaggtga aagtggagaa cattcgaaca atagatatgc tcaaagcaag agtgaaaaat 60
 cgtgtggcac gcagcaaatg ctttaaaaaat gcctctttga tcttaatagg aataactaca 120
 ttgagtatag ccttcaatat ctatctgac ataaactata caatgcaaga aaacacatcc 180
 gaatcagaac atcacaccag ctcatcacc atggaatcca gcagggaac tccaacggtc 240
 cccatggact acccagacac caatccaggc tcacagtatc caactcaaca gtccacagaa 300
 ggctccacac tctactttgc agcctcagca agctcaccag agacagaacc aacatcaaca 360
 ccaaacacaa caagccggcc gcccttcgtc gacacacaca caacaccacc aagtgaagc 420
 agaacaaaga caagtccggc agtccacaca aaaaacaatc caaggataag cctcagaaca 480
 cattccccac catggggcaat gacaaggacg gtccgaggaa ccaccactct ccgcacaagc 540
 agcacaagaa aaagaccgtc tacagcatca gtccaacctg acagcagcgc aacaaccac 600
 aaacacgaag aagcaagccc agtgagcccg caaacatctg caagcacagc aagaccataa 660
 aggaaggtca tggaggccag cacatcaaca acatacaatc aaactagtta a 711

26/52

<210> 62
 <211> 711
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(711)
 <223> human metapneumovirus, strain Quebec Isolate No C-77748

<400> 62							
atggagggtga	aagtggagaa	cattcgaaca	atagatatgc	tcaaagcaag	agtgaaaaat		60
cgtgtggcac	gcagcaaatg	ctttaaaaaat	gcctctttga	tcctaataagg	aataactaca		120
ttgagtatatg	ccctcaatat	ctatctaatac	ataaactata	caatgcaaga	aaacacatcc		180
gaatcagaac	atcacaccag	ctcatcaccc	atggaatcca	gcagggaaac	tccaacagtc		240
cctatggaca	actcagacac	caatccaggc	tcacagtatc	caactcaaca	gtccacagaa		300
ggctccacac	tctactttgc	agcctcagca	agctcaccag	agacagaacc	aacatcaaca		360
ccagacacaa	caagccgccc	gcccttcgtc	gacacacaca	caacaccacc	aagtgcgaagc		420
agaacaagga	caagtccggc	agtccacaca	aaaaacaatc	caaggacaag	ccccagaata		480
cattccccac	catgggcaat	gacaaggacg	gtccgtggaa	ccaccactct	ccgcacaagc		540
agcacaagaa	aaagaccgtc	tacagcatca	gtccaacctg	acagcagcgc	aacaaccac		600
aaacacgaag	aagcaagccc	agtgcagccc	caaacatctg	cgagcacagc	aagaccacaa		660
aggaaggcca	tggaggccag	cacatcaaca	acatacaacc	aaactagtta	a		711

<210> 63
 <211> 711
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(711)
 <223> human metapneumovirus, strain Quebec Isolate No C-78132

<400> 63							
atggagggtga	aagtggagaa	cattcgaaca	atagatatgc	tcaaagcaag	agtgaaaaat		60
cgtgtggcac	gcagcaaatg	ctttaaaaaat	gcctctttga	tcctaataagg	aataactaca		120
ttgagtatatg	ccctcaacat	ctatctgatac	ataaactata	caatgcaaga	aaacacatcc		180
gaatcagaac	atcacaccag	ctcatcaccc	atggaatcca	gcagggaaac	tccaacggtc		240
cctatagaca	actcagacac	caatccaggc	tcacagtatc	caactcaaca	gtccacagaa		300
ggctccacac	tccactttgc	agcctcagca	agctcaccag	agacagaacc	aacatcaaca		360
ccagacacaa	caagccgccc	gcccttcgtc	gacacacaca	caacaccacc	aagtgcgaagc		420
agaacaagga	caagtccggc	agtccacaca	aaaaacaatc	taaagataag	ccccagaata		480
cattccccac	catgggcaat	gacaaggacg	gtccgtggaa	ccaccactct	ccgcacaagc		540
agcacaagaa	aaagaccgtc	tacagcatca	gtccaacctg	acagcagcac	aacaactcac		600
aaacacgaag	aagcaagccc	agtgcagccc	caaacatctg	caagcacagc	aagaccacaa		660
aggaaggcca	tggaggccag	cacatcaaca	acatacaacc	aaactagtta	a		711

<210> 64
 <211> 711
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(711)
 <223> human metapneumovirus, strain Quebec Isolate No C-78406

<400> 64							
atggagggtga	aagtggagaa	cattcgaaca	atagatatgc	tcaaagcaag	agtgaaaaat		60
cgtgtggcac	gcagcaaatg	ctttaaaaaat	gcctctttga	tcctaataagg	aataactaca		120
ttgagtatatg	ccctcaacat	ctatctgatac	ataaactata	caatgcaaga	aaacacatcc		180
gaatcagaac	atcacaccag	ctcatcaccc	atggaatcca	gcagggaaac	tccaacggtc		240
cctatagaca	actcagacac	caatccaggc	tcacagtatc	caactcaaca	gtccacagaa		300
ggctccacac	tccactttgc	agcctcagca	agctcaccag	agacagaacc	aacatcaaca		360
ccagacacaa	caagccgccc	gcccttcgtc	gacacacaca	caacaccacc	aagtgcgaagc		420
agaacaagga	caagtccggc	agtccacaca	aaaaacaatc	taaagataag	ccccagaata		480
cattccccac	catgggcaat	gacaaggacg	gtccgtggaa	ccaccactct	ccgcacaagc		540
agcacaagaa	aaagaccgtc	tacagcatca	gtccaacctg	acagcagcac	aacaactcac		600

27/52

aaacacgaag aagcaagccc agtgagccca caaacatctg caagcacagc aagaccacaa 660
 aggaaggcca tggaggccag cacatcaaca acatacaacc aaactagtta a 711

<210> 65
 <211> 711
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(711)
 <223> human metapneumovirus, strain Quebec Isolate No
 C-73750

<400> 65
 atggagggtga aagtggagaa cattcgaaca atagatatgc tcaaagcaag agtgaaaaat 60
 cgtgtggcac gcagcaaatg ctttaaaaaat gcctctttga tcctaataagg aataactaca 120
 rtgagtatag ccctcaacat ctatctgac ataaactata caatgcaaga aaacacatcc 180
 gaatcagaac atcacaccag ctcacacccc atggaatcca gcagggaac tccaacgggtc 240
 cctatagaca actcagacac caatccaggc tcacagtatc caactcaaca gtccacagaa 300
 ggctccacac tccactttgc agcctcagca agctcaccag agacagaacc aacatcaaca 360
 ccagacacaa caagccgccc gcccttcgtc gacacacaca caacaccacc aagtgaagc 420
 agaacaagga caagtccggc agtccacaca aaaaacaatc taaagataag ccccgaaaca 480
 cattccccac catgggcaat gacaaggacg gtccgtggaa ccaccactct ccgcacaagc 540
 agcataagaa aaagaccgtc cacagcatca gtccacctg acagcagcac aacaaccac 600
 aaacacgaag aagcaagccc agtgagccca caaacatctg caagcacagc aagaccacaa 660
 aggaaggcca tggaggccag cacatcaaca acacacaacc aaactagtta a 711

<210> 66
 <211> 660
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(660)
 <223> human metapneumovirus, strain Quebec Isolate No
 C-60298

<400> 66
 atggagggtga aagtggagaa cattcgaaca atagatatgc tcaaagcaag agtgaaaaat 60
 cgtgtggcac gtagcaaatg ctttaaaaaat gcttctttaa tcctcatagg aataactaca 120
 ctgagtatag ctctcaatat ctatctgac ataaactaca caatacaaaa aacctcatct 180
 gaatcagaac accacaccag ctcaccaccc acagaatcca acaaggaagc ttcaacaatc 240
 tccacagaca acccagacat caatccaaac tcacagcatc caactcaaca gtccacagaa 300
 aacccccacac tcaacccccgc agcatcagtg agcccatcag aaacagaacc agcatcaaca 360
 ccagacacaa caaacccgct gtccctcgta gacaggtcca cagcacaacc aagtgaagc 420
 agaacaagaa caaaaaccgac agtccacaca agaaacaacc caagcacagc ttccagtaga 480
 caatccccac caccgggcaac aacgaaggca atccgcagag ccaccacttt ccgcagtagc 540
 agcacaggaa aaagaccaac cacaacatca gtccagtccg acagcagcac cacaacccaa 600
 aatcatgaag aaacagggtc agcgaaccca caggcatctg taagcacat gcaaaactag 660

<210> 67
 <211> 654
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(654)
 <223> human metapneumovirus, strain Quebec Isolate No
 C-74783

<400> 67
 atggagggtga aagtggagaa cattcgaaca atagatatgc tcaaagcaag agtgaaaaat 60
 cgtgtggcat gtagcaaatg ctttaaaaaat gcttctttaa tcctcatagg aataactaca 120
 ctgagtatag ctctcaatat ctatctgac ataaaccaca caatacaaaa aaccacatct 180
 gaatcagaac accacaccag ctcaccaccc acagaatcta acaaggaagc ttcaacaatc 240
 tccacagaca acccagacat caatccaaac tcacagcatc caactcaaca gtctacagaa 300
 aacccccacac tcaacccccgc agtatcaaca agcccatcag aaacagaacc aacatcaaca 360
 ccagacacaa caaacccgct gtccctcgta gacaggtcca cagcacaacc aagtgaagc 420
 agaacaagaa caaaatcgac agtccacaca agaaacaacc caagcacagc ttccagcaca 480

28/52

caatccccac	cacgggcaac	aacgaaggca	atccgcagag	ccaccacttt	ccgcatgagc	540
agcacagaaa	aaagaccaac	cacaacacta	gtccagtacg	acagcagcac	cacaacccaa	600
aatcatgaag	aaacaggttc	agcgaacca	caggcatctg	caagcacaat	gtaa	654

<210> 68
 <211> 660
 <212> pNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(660)
 <223> human metapneumovirus, strain Quebec Isolate No C-74549

<400> 68						
atggagggtga	aagtggagaa	cattcgagca	atagacatgc	tcaaagcaag	agtgaaaaat	60
cggtgtggcac	gtagcaaatg	cttcaaaaat	gttccccaa	tcctcacagg	aataaccaca	120
ctgagcatag	ctctcaatat	ccacctgatc	ataaactaca	caatacaaaa	aaccacatct	180
gaatcagaac	accacaccag	ctcaccaccc	acagaatcta	acaaggaagc	ttcaacaatc	240
tccacagaca	acccagacat	caatccaaac	tcacagcatc	caactcaaca	gtctacagaa	300
aaccccacac	tcaacccgc	agtatcaaca	agcccatcag	aaacagaacc	aacatcaaca	360
ccagacacaa	caaaccgcct	gtcctccgta	gacagggtcca	cagcacaacc	aagtgaagc	420
agaacaaaga	caaaatcgac	agtccacaca	agaaacaacc	caagcacagc	ttccagcaca	480
caatccccac	cacgggcaac	aacgaaggca	atccgcagag	ccaccacttt	ccgcatgagc	540
agcacagaaa	aaagaccaac	cacaacatta	gtccagtacg	acagcagcac	cacaacccaa	600
aatcatgaag	aaacaggttc	agcgaacca	caggcatctg	taagcacaat	gtaaaactag	660

<210> 69
 <211> 711
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(711)
 <223> human metapneumovirus, strain Quebec Isolate No C-77807

<400> 69						
atggagggtga	aagtggagaa	cattcgaaca	atagatatgc	tcaaagcaag	agtgaaaaat	60
cggtgtggcac	gcagcaaatg	ctttaaaaat	gcctctttga	tcctaataagg	aataacgaca	120
ttgagttag	ccctcaacat	ctatctgatc	ataaactata	caatgcaaga	aaacacatcc	180
gaatcagaac	atcacaccag	ctcatcacc	atgggaatcca	gcagggaaac	tccaacggtc	240
ccatagaca	actcagacac	caatccaggc	tcacagtatc	caactcaaca	gtccacagaa	300
ggctccacac	tccactttgc	agcctcagca	agctcaccag	agacagaacc	aacatcaaca	360
ccagacacaa	caagccgccc	gcccttcgtc	gacacacaca	caacaccacc	aagtgcagc	420
agaacaagga	caagtccggc	agtccacaca	aaaaacaatc	taaagataag	ccccagaata	480
cattccccac	catgggcaat	gacaaggacg	gtccgtggaa	ccaccactct	ccgcacaagc	540
agcacaagaa	aaagaccgtc	tacagcatca	gtccaacctg	acagcagcac	aacaactcac	600
aaacacgaag	aagcaagccc	agtgagccca	caaacatctg	caagcacagc	aagaccacaa	660
aggaaggga	tggaggccag	cacatcaaca	acatacaacc	aaactagtta	a	711

<210> 70
 <211> 711
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(711)
 <223> human metapneumovirus, strain Quebec Isolate No C-78163

<400> 70						
atggagggtga	aagtggagaa	cattcgaaca	atagatatgc	tcaaagcaag	agtgaaaaat	60
cggtgtggcac	gcagcaaatg	ctttaaaaat	gcctctttga	tcctaataagg	aataactaca	120
ttgagttag	ccctcaacat	ctatctgatc	ataaactata	caatgcaaga	aaacacatcc	180
gaatcagaac	atcacaccag	ctcatcacc	atgggaatcca	gcagggaaac	tccaacggtc	240
ccatagaca	actcagacac	caatccaggc	tcacagtatc	caactcaaca	gtccacagaa	300
ggctccacac	tccactttgc	agcctcagca	agctcaccag	agacagaacc	aacatcaaca	360
ccagacacaa	caagccgccc	gcccttcgtc	gacacacaca	caacaccacc	aagtgcagc	420

29/52

```

agaacaagga caagtccggc agtccacaca aaaaacaatc taaagataag cccagaata 480
cattccccac catgggcaat gacaaggacg gtccgtggaa ccaccactct ccgcacaagc 540
agcacaagaa aaagaccgtc tacagcatca gtccaacctg acagcagcac aacaactcac 600
aaacacgaag aagcaagccc agtgagccca caaacatctg caagcacagc aagaccacaa 660
aggaagggca tggaggccag cacatcaaca acatacaacc aaactagtta a 711

```

<210> 71
 <211> 711
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(711)
 <223> human metapneumovirus, strain Quebec isolate No C-68661

```

<400> 71
atggagggtga aagtggagaa cattcgaaca atagatagcg tcaaagcaag agtgaaaaat 60
cgtgtggcac gcagcaaatg ctttaaaaat gcctctttga tcctaataagg aataactaca 120
ttgagtatag ccctcaatat ctatctgacg ataaactata caatgcaaga aaacacatcc 180
aaatcagaac atcacaccag ctcatcaccg atggaatcca gcagggaac tccaacgggtc 240
cctatgagac actcagacac caatccaggc tcacagtatc caactcaaca gtccacagaa 300
ggctccacac tccactttgc agcctcagca agctcaccag agacagaacc aacatcaaca 360
ccagacacaa caagccgccc gcccttcgtc gacacacaca caacaccacc aagtgaagc 420
agaacaagga caagtccggc agtccacaca aaaaacaatc caaggataag cccagaaca 480
cattccccac catgggcaat gacaaggacg gtccgtggaa ccaccactct ccgcacaagc 540
agcacaagaa aaagaccgtc cacagcatca gtccaacctg acagcagcgc aacacccac 600
aaacacgaag aagcaagccc agtgagccca caaacatctg caagcacagc aagaccacaa 660
aggaagggca tggaggccag cacatcaaca acatacaacc aaactagtta a 711

```

<210> 72
 <211> 711
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(711)
 <223> human metapneumovirus, strain Quebec isolate No C-88448

```

<400> 72
atggagggtga aagtggagaa cattcgaaca atagatatgc tcaaagcaag agtgaaaaat 60
cgtgtggcac gcagcaaatg ctttaaaaat gcctctttga tcctaataagg aataactaca 120
ttgagtatag ccctcaatat ctatctaatt ataaactata caatgcaaga aaacacatcc 180
gaatcagaac atcacaccag ctcatcaccg atggaatcca gcagggaac tccaacagtc 240
cctatggaca actcagacac caatccaggc tcacagtatc taactcaaca gtccacagaa 300
ggctccacac tctactttgc agcctcagca agctcaccag agacagaacc aacatcaaca 360
ccagacacaa caagccgccc gcccttcgtc gacacacaca caacaccacc aagtgaagc 420
agaacaagga caagtccggc agtccacaca aaaaacaatc caaggacaag cccagaaca 480
cattccccac catgggcaat gacaaggacg gtccgtggaa ccaccactct ccgcacaagc 540
agcacaagaa aaagaccgtc cacagcatca gtccaacctg acagcagcgc aacaaccac 600
aaacacgaag aagcaagccc agtgagcccg caaacatctg cgagcacagc aagaccacaa 660
aggaagggca tggaggccag cacatcaaca acatacaacc aaactagtta a 711

```

<210> 73
 <211> 711
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(711)
 <223> human metapneumovirus, strain Quebec isolate No C-88470

```

<400> 73
atggagggtga aagtggagaa cattcgaaca atagatatgc tcaaagcaag agtgaaaaat 60
cgtgtggcac gcagcaaatg ctttaaaaat gcctctttga tcctaataagg aataactaca 120
ttgagtatag ccctcaatat ctatctgacg ataaactata caatgcaaga aaacacatcc 180
gaatcagaac atcacaccag ctcatcaccg atggaatcca gcagggaac tccaacgggtc 240

```

30/52

```

ccatggact acccagacac caatccagge tcacagcatc caactcaaca gtccacagaa 300
ggctccacac tctactttgc agcctcagca agctcaccag agacagaacc aacatcaaca 360
caaacacaa caagccgccc gcccttcgtc gacacacaca caacaccacc aagtgcgaagc 420
agaacaaaga caagtccggc agtccacaca aaaaaaatc caaggataag cctcagaaca 480
cattccccac catgggcaat gacaaggacg gtccgaggaa ccaccactct ccgcacaagc 540
agcacaaaga aaagaccgtc tacagcatca gtccaaacctg acagcagcgc aacaaccac 600
aaacacgaag aagcaagccc agtgagccc caaacatctg caagcacagc aagaccaca 660
aggaaggtca tggaggccag cacatcaaca acatacaacc aaactagtta a 711

```

<210> 74
 <211> 711
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(711)
 <223> human metapneumovirus, strain Quebec Isolate No C-86316

```

<400> 74
atggagggtga aagtggagaa cattcgaaca atagatatgc tcaaagcaag agtgaaaaat 60
cgtgtggcac gcagcaaatg ctttaaaaaat gcctcttga tcctaataagg aataactaca 120
ttgagtatag cccicaatat ctatcfaatc ataaactata caatgcaaga aaacacatcc 180
gaatcagaac atcacaccag ctcatcaccg atggaatcca gcaaggaaac tccaacagtc 240
cctatggaca actcagacac caatccagge tcacagtatc caactcaaca gtccacagaa 300
ggctccacac tctactttgc agcctcagca agctcaccag agacagaacc aacatcaaca 360
ccagacacaa caagccgccc gcccttcgtc gacacacaca caacaccacc aagtgcgaagc 420
agaacaaaga caagtccggc agtccacaca aaaaaaatc caaggacaag cctcagaaca 480
cattccccac catgggcaat gacaaggacg gtccgtggaa ccaccactct ccgcacaagc 540
agcacaaaga aaagaccgtc cacagcatca gtccaaacctg acagcagcgc aacaaccac 600
aaacacgaag aagcaagccc agtgagccc caaacatctg cgagcacagc aagaccaca 660
aggaaggtca tggaggccag cacatcaaca acatacaacc aaactagtta a 711

```

<210> 75
 <211> 6017
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(6017)
 <223> human metapneumovirus, strain 00-1 accession number af371337

```

<400> 75
atggatcctc tcaatgaatc cactgttaat gtctatcttc ctgactcata tcttaaagga 60
gtgatttcct ttagtgagac taatgcaatt gggtcatgtc tcttaaaaag accttaccta 120
aaaaatgaca aactgcaaaa agttgccata gagaatcctg ttatcgagca tgtagactc 180
aaaaatgcag tcaattctaa gatgaaaata tcagattaca agatagtaga gccagttaac 240
atgcaacatg aaattatgaa gaatgtacac agttgtgagc tcacattatt aaaaacagttt 300
ttaacaagga gtaaaaatat tagcactctc aaattaaata tgatatgtga ttggctgcag 360
ttaaagtcta catcagatga tactcaatc ctaagtttta tagatgtaga atttatacct 420
agctgggtaa gcaattgggt tagtaattgg tacaatctca acaagttgat tctggaattc 480
aggaagaag aagtaataag aactgggtca atcttgtgta ggtcattggg taaattagtt 540
tttgtgtgat catcatatgg atgtatagtc aagagcaaca aaagcaaaa agtgagcttc 600
ttcacatata atcaactggt aacatggaaa gatgtgatgt taagtagatt caatgcaaat 660
ttttgtatat gggtgaagca cagtctgaat gaaaatcaag aagggctagg gttgagaagt 720
aatctgcaag gcatattaac taataagcta tatgaaactg tagattatat gcttagttta 780
tgttgcaatg aagggttctc acttgtgaaa gagttcgaag gctttattat gagtgaattt 840
cttaggatta ctgaacatgc tcaattcagt actagattta gaaatacttt attaaatgga 900
ttaactgac aattaacaaa attaaaaaat aaaaacagac tcagagttca tggtagcgtg 960
ttgaaaaata atgattatcc aatgtacgaa gttgtactta agttattagg agatactttg 1020
agaigtatta aattattaat caataaaaaac ttagagaatg ctgctgaatt atactatata 1080
tttagaatat tcggtcacc aatggtagat gaaagagatg caatggatgc tgtcaaat 1140
aacaattaa tcacaaaaat ccttaggtgg gagagcttga cagaactaag aggggcattc 1200
atattaagga ttatcaaaag atttgtagac aacaacaaa gatggcccaa aattaaaaac 1260
traaaagtgc ttgtaagag atggactatg tacttcaaa caaaaagtta cccagtc 1320
cttgaaattaa ggaacaaga ttttttagag cttgtgcgaa tacagtttga acaagagttt 1380
tctgtccctg aaaaaaccaa ccttgagatg gtattaaatg ataaagctat atcacctcct 1440
aaaagattaa tatggtctgt gtatccaaa aattacttac ctgagaaaat aaaaaatcga 1500
tatctagaag agactttcaa tgcaagtgat agtctcaaaa caagaagagt actagagtac 1560

```

31/52

tatttgaag	ataataaatt	cgaccaaaaa	gaacttaaaa	gttatgtgtg	taaacaagaa	1620
tatttaaatg	ataaggatca	tattgtctcg	ctaactggaa	aagaaagaga	attaagtgtg	1680
ggtagaatgt	ttgctatgca	accaggaaaa	cagcgacaaa	tacaaatatt	ggctgaaaaa	1740
ttgttagctg	ataatattgt	accttttttc	ccagaaacct	taacaaagta	tggtgatcta	1800
gatcttcaga	gaataatgga	aatcaaatecg	gaactttctt	ctattaaaac	tagaagaaat	1860
gatagttata	ataattacat	tgcaagagca	tccatagtaa	cagatttaag	taagttcaac	1920
caagccttta	ggtagaaac	tacagcgatc	tgtagcgatg	tagcagatga	actacatgga	1980
acacaaagcc	tattctgttg	gttacatctt	atcgctccca	tgacaacaat	gatatgtgcc	2040
tatagacatg	caccaccaga	aacaaaaggt	gaatatgata	tagataagat	agaagagcaa	2100
agtgggttat	atagatatca	tatgggtggg	attgaaggat	gggtgcaaaa	actctggaca	2160
atggaagcta	tatctctatt	agatgttgta	tctgtaaaaa	cacgatgtca	aatgacatct	2220
ttattaaacg	gtgacaacca	atcaatagat	gtaagttaac	cagtttaagt	atctgagggg	2280
ttagatgaag	tgaagcaga	ttatagcttg	gctgtaaaaa	tgtaaaaaa	aataagagat	2340
gcatacagaa	atataggcca	ttaaactaaa	gaaggggaaa	catatatac	aagagatctt	2400
cagttttata	gtaagggtgat	tcaatctgaa	ggagtaatgc	atcctacccc	tataaaaaag	2460
atcttaagag	tggaaccatg	gataaacaca	atattagatg	acattaaaa	cagtcagag	2520
tcaataggga	gtctatgtca	ggaattagaa	tttagggggg	aaagcataat	agttagtctg	2580
atatttaagg	atctttggct	gtataattta	tacatgcatg	aatcaaagca	acacccctta	2640
gcaggggaag	agttattcaa	acaactaaat	aaaacattaa	catcagtgca	gagatttttt	2700
gaaataaaaa	aggaataatga	agttagtagat	ctatggatga	acataccaat	gcagtttgga	2760
ggaggagatc	cagtagtctt	ctatagatct	ttctatagaa	ggacccctga	ttttttaact	2820
gaagcaatca	gtcatgtggg	tattctgtta	agaatatcag	ccaacataag	aaatgaagcg	2880
aaaaataagt	tcttcaagag	cttactgtca	atagaaaaaa	atgaacgtgc	tacactgaca	2940
acactaatga	gagatcctca	agctgttggc	tcagagcgac	aagcaaaagt	aacaagtgat	3000
atcaatagaa	cagcagttac	cagcatctta	agctcttctc	caaatcaact	tttcagcgat	3060
agtgtctatc	actacagtag	aaatgaagaa	gaggtcggaa	tcattgtctg	caacataaca	3120
cctgtttatc	ctcatggact	gagagttttg	tatgaatcat	taccttttca	taaagctgaa	3180
aaagtgttga	atatgatata	aggaacgaaa	tcataacca	acttattaca	gagaacatct	3240
gctattatag	gtgaagatat	tgacagagct	gtatccatga	tgctggagaa	cctaggatta	3300
ttatctagaa	tattgtcagc	agttgttgat	agtatagaaa	ttccaacca	atctaatggg	3360
aggctgtat	gttgtcagat	atctagaacc	ctaagggaga	catcatggaa	taatatggaa	3420
atagtgtgag	taacatcccc	tagcatcact	acatgcatgg	atgtcatata	tgcaactagc	3480
tctcatttga	aagggataat	cattgaaaag	ttcagcactg	acagaactac	aagaggtcaa	3540
agaggtccaa	agagcccttg	ggtagggctg	agcactcaag	agaaaaaatt	agttcctggt	3600
tataacagac	aaattctttc	aaaaacaaca	agagaacagc	tagaagcaat	tggaaaaaatg	3660
agatgggtat	ataaagggag	accaggttta	agacgattac	tcaataagat	ttgtcttgga	3720
agtttiaggc	ttagtataca	atgtgtaaaa	cctttattac	ctaggtttat	gagtgtaaat	3780
ttcctacaca	ggttatctgt	cagtagtaga	cctatggaa	tcccagcatc	agttccagct	3840
tatagaacaa	caaattacca	ttttgacact	agtcctatta	atcaagcact	aagtggagaa	3900
tttggaatg	aagatattaa	tttggtcttc	caaaatgcaa	tcagctgtgg	aattagcata	3960
ttgagtgtag	tagaacaatt	aactggtagg	agttcctaaa	agtttagttt	aataacctcaa	4020
ttagaagaaa	tagacattat	gccaccacca	gtgtttcaag	ggaaattcaa	ttataagcta	4080
gtagataaga	taacttttga	tcaacataat	ttcagtcag	acaaaaatga	tatgttaaca	4140
ctggggaaaa	tgctcatggc	cactataaaa	ggtcagaaaa	cagatcagtt	cctgaacaa	4200
agagagaatt	atcttcatgg	gaataatctt	attgagctct	tgtagcagc	gttagcatgt	4260
cattgggtgtg	ggatattaa	agagcaatgt	atagaaaaa	atattttcaa	gaaagactgg	4320
gggtgacggg	tcatactggg	tcattgcttt	atggacttca	aaatattcct	atgtgtcttt	4380
aaaactaaac	ttttatgtag	ttgggggtcc	caagggaaaa	acattaaaga	tgaagatata	4440
gtagatgaat	caatagataa	actgttaagg	attgataata	ctttttggag	aatgttcagc	4500
aaagttatgt	ttgaattcaa	ggtttaagaa	aggataatgt	tatatgatgt	aaaattttca	4560
tcattagtag	gttatatagg	gtttaagaat	tggtttatag	aacagtggga	tcagctgagt	4620
tgcatgaggt	accttggatt	gtcaatgccg	aaggtgatct	gggtgagatc	aagtcaatta	4680
aaatctattt	gcaactgata	gagcaaaagt	tatttttaag	aataactgtt	ttgaactata	4740
cagatatggc	acatgctctc	acaagattaa	tcagaaagaa	gttgatgtgt	gataatgcac	4800
tattaaactcc	gatttccatcc	ccaatggtta	atttaactca	agttattgat	cctacagaac	4860
aattagctta	tttccctaa	ataacatttg	aaaggctaaa	aaattatgac	actagtcca	4920
attatgctaa	aggaagctta	acaaggaatt	acatgatact	gttgccatgg	caacatgta	4980
atagatatata	ctttgtcttt	agttctactg	gatgtaaagt	tagtctaaaa	acatgcattg	5040
gaaaacttat	gaaagatcta	aaacctaaag	ttctgtaact	tattggagaa	ggggcaggaa	5100
attggatggc	cagaacagca	tgtagaatat	ctgacatcaa	atttgtatag	agaagtttaa	5160
aagatgacct	tgatcatcat	tatcctttgg	aataccagag	agttatagga	gaattaaagc	5220
ggataataga	tagcgggtgaa	gggctttcaa	tggaacaac	agatgcaact	caaaaaactc	5280
attgggattt	gatacacaga	gtaagcaaa	atgctttatt	aataacttta	tgtagtcag	5340
aatttaagga	cagagatgat	tttttaaga	tggttaattct	atggaggaaa	catgtattat	5400
catgcagaat	ttgtcactact	tatgggacag	acctctattt	atcgcaaa	tatcatgcta	5460
aagactgcaa	tgtaaaaata	cctttttttg	tgagatcagt	agccaccttt	attatgcaag	5520
gtagttaact	gtcaggctca	gaatgctaca	tactcttaac	actaggccac	cacaacaatt	5580
taccttgcca	tggaagaaata	caaaaattcta	agatgaaaa	agcagtggtg	aatgattttt	5640
atgctgcaaa	aaaacttgac	aataaatcta	ttgaagccaa	ctgtaaatca	cttttatcag	5700
ggctaaagat	accgataaat	aagaaagaa	taaatagaca	gagaaggtta	tttaacttac	5760
aaagcaacca	ttcttctgta	gcaacagttg	gaggtagcaa	ggtcatagag	tctaaatggt	5820
taacaaacaa	ggcaaacaca	ataattgatt	ggttagaaca	tatttttaaa	tctccaaaa	5880

32/52

gtgaattaaa ttatgatttt ttggaagcat tagaaaatac ttaccctaata atgattaaac 5940
 taatagataa tctaggggaat gcagagataa aaaaactgat caaagtaact ggatatatgc 6000
 ttgtaagtaa aaaatga 6017<210> 76

<211> 450
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(450)
 <223> human metapneumovirus, strain Quebec Isolate No
 hMPV-118

<400> 76
 taaaatgaaa atatcagatt acaaggtagt agaaccagta aacatgcaac atgaaataat 60
 gaagaatgta cacagctgtg agcacacact attgaaacag tttttaacaa ggagtataaa 120
 ctttagcact ctcaaatata atatgatatg tgattggctg caattaaaaa ctacatcaga 180
 tggtagctca atcctaagtt ttatagatgt agaatttata cctagttggg taagcaactg 240
 gtttagtaat tggtagaact tcaataagtt aatttttagaa tttaggagag aagaagtaat 300
 aagaaccggt tcaatcttgt gcaggtcatt aggtaaatta gtttttattg tatcatcata 360
 tggatgtatc gtcaagagca acaaaagcaa aagagtggagc ttctttacat acaatcaact 420
 gttacatgg aaagatgtga tgttaagtcg 450

<210> 77
 <211> 450
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(450)
 <223> human metapneumovirus, strain Quebec Isolate No
 hMPV-163

<400> 77
 ttgaaataatc agattacagg atagtagagc cagtaaaccat gcaacatgaa attatgaaga 60
 atgtacacag ttgtgagctc acattattaa aacagttttt aacaaggagt aaaaatatta 120
 gcactctcaa attaaatatt atattgtgatt ggctgcagtt aaaaatcaca tcagatgata 180
 cctcaatcct aagttttata gatgtagaat ttatacctag ctgggtaagc aattgggtta 240
 gtaatttgta caatctcaac aagttgattc tggaaattcag gaaagaagaa gtaataagaa 300
 ctggttcaat cttgtgtagg tcattaggtta aattagtttt tgttgtatca tcatatggat 360
 gtatagtcaa gagcaacaaa agcaaaagag tgagcttctt cacatacaat caactgttaa 420
 catggaaaga tgtgatgtta agtagattca 450

<210> 78
 <211> 500
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(500)
 <223> human metapneumovirus, strain Quebec Isolate No
 hMPV-174

<400> 78
 gagaatcctg ttatcgagca tgttagactc aaaaatgcaa tcaatctaaa gatgaaaata 60
 tcagattaca ggatagtaga gccagtaaac atgcaacatg aaattatgaa gaatgtacac 120
 agttgtgagc tcacattatt aaaacagttt ttaacaagga gtaaaaatat tagcactctc 180
 aaattaaata tgatagtga ttggctgcag ttaaaagtcta catcagatga taacctcaatc 240
 ttaagtttta tagatgtaga atttatacct agctgggtaa gcaattgggt tagtaattgg 300
 tacaatctca acaagttgat tctggaattc aggaagaag aagtaataag aactgggttca 360
 atcttgtgta ggtcattggg taaattagtt tttgtgtat catcatatgg atgtatagtc 420
 aagagcaaca aaagcaaaag agtgagcttc ttcacatata atcaactgtt aacatggaaa 480
 gatgtgatgt taagtagatt 500

<210> 79
 <211> 450
 <212> DNA
 <213> human metapneumovirus

<220>

33/52

<221> gene
 <222> (1)...(450)
 <223> human metapneumovirus, strain Quebec Isolate No
 hMPV-184

<400> 79
 attgaaatat cagattacag gatagtagag ccagtaaaca tgcaacatga aattatgaag 60
 aatgtacaca gttgtgagct cacattatta aaacagtttt taacaaggag taaaaatatt 120
 agcactctca aattaaatat gatatgtgat tggctgcagt taaagtctac atcagatgat 180
 acctcaatct taagttttat agatgtagaa ttatatccta gctgggtaag caattgggtt 240
 agtaattggt acaatctcaa caagttgatt ctggaattca ggaaagaaga agtaataaga 300
 actggttcaa tcttgtgtag gtcattaggt aaattagttt ttgttgtatc atcatatgga 360
 tgtatagtca agagcaacaa aagcaaaaga gtgagcttct tcacatacaa tcaactgtta 420
 acatggaaag atgtgatgtt aagtagattc 450

<210> 80
 <211> 500
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(500)
 <223> human metapneumovirus, strain Quebec Isolate No
 hMPV-196

<400> 80
 ttgcccatag agaatcctgt tattgagcat gtgagactca aaaatgcagt caattctaaa 60
 atgaaaatat cagattacaa ggtagttagag ccagtaaaca tgcaacatga aataatgaag 120
 aatgtacaca gttgtgagct cacactattg aaacagtttt taacaaggag taaaaacatt 180
 agcactctca aattaaatat gatatgtgat tggctgcagt taaagtctac atcagatgat 240
 acctcaatcc taagtttcat agatgtagaa ttatatccta gttgggtaag caactgggtt 300
 agtaattggt acaatctcaa taagttaatt ttggaattca gaagagaaga agtaataaga 360
 accggttcaa tcttatgcag gtcattgggt aaattagttt ttattgtatc atcatatgga 420
 tgtatcgtca agagcaacaa aagcaaaaga gtgagcttct tcacatacaa tcaactgtta 480
 acatggaaag atgtgatgtt 500

<210> 81
 <211> 450
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(450)
 <223> human metapneumovirus, strain Quebec Isolate No
 C-76837

<400> 81
 agatgaaaat atcagattac aggatagtag agccagtaaa catgcaacat gaaattatga 60
 agaatgtaca cagttgtgag ctcacattat taaaacagtt ttaacaagg agtaaaaaata 120
 ttgacctctt caaattaaat atgatatgtg attggctgca gttaaagtct acatcagatg 180
 atacctcaat cttaagtttt atagatgtag aattttatacc tagctgggta agcaattgggt 240
 ttagttaatt gtacaatctc aacaagttga ttctggaatt caggaaagaa gaagtaataa 300
 gaactgggtt aatcttgtgc aggtctttgg gtaaattagt ttttgttgta tcatcatatg 360
 gatgtatagt caagagcaac aaaagcaaaa gagtgagctt cttcacatac aatcaactgt 420
 taacatggaa agatgtgatg ttaagtagat 450

<210> 82
 <211> 450
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(450)
 <223> human metapneumovirus, strain Quebec Isolate No
 C-77807

<400> 82
 taagatgaaa atatcagatt acaggatagt agagccagta aacatgcaac atgaaattat 60
 gaagaatgta cacagttgtg agctcacatt attaaaacag tttttaacaa ggagtaaaaa 120

34/52

tattagcact	ctcaaattaa	atatgatatg	tgattggctg	cagttaaagt	ctacatcaga	180
tgatacctca	atcttaagtt	ttatagatgt	agaattttata	cctagctggg	taagcaattg	240
gtttagtaat	tggtacaatc	tcaacaagtt	gattctggaa	ttcaggaaag	aagaagtaat	300
aagaactggg	tcaatcttgt	gtagggtcatt	gggtaaatta	gtttttgttg	tatcatcata	360
cgatgtata	gtcaagagca	acaaaagcaa	aagagtggagc	ttcttcacat	acaatcaact	420
gttaacatgg	aaagatgtga	tgtaagtag				450
 <210> 83						
<211> 450						
<212> DNA						
<213> human metapneumovirus						
 <220>						
<221> gene						
<222> (1)...(450)						
<223> human metapneumovirus, strain Quebec Isolate No hmpv-185						
 <400> 83						
taaaattgaa	atatcagatt	acaaggtagt	agagccagta	aacatgcaac	atgaaataat	60
gaagaatgta	cacagttgtg	agctcacact	attgaaacag	tttttaacaa	ggagtaaaaa	120
cattagcact	ctcaaattaa	atatgatatg	tgattggctg	caattaaagt	ctacatcaga	180
tgatacctca	atcctaagtt	tcatagatgt	agaattttata	cctagttggg	taagcaactg	240
gtttagtaat	tggtacaatc	tcaataagtt	aattttggaa	ttcagaagag	aagaagtaat	300
aagaaccggg	tcaatcttat	gcagggtcatt	gggtaaatta	gtttttattg	tatcatcata	360
tggtgtatc	gtcaagagca	acaaaagcaa	aagagtggagc	ttcttcacat	acaatcaact	420
gttaacatgg	aaagatgtga	tgtaagtag				450
 <210> 84						
<211> 500						
<212> DNA						
<213> human metapneumovirus						
 <220>						
<221> gene						
<222> (1)...(500)						
<223> human metapneumovirus, strain Quebec Isolate No C-7B123						
 <400> 84						
atcctgttat	cgagcatgtt	agactcaaaa	atgcaatcaa	ttctaagatg	aaaatatcag	60
attacaggat	agttagagcca	gtaaacatgc	aacatgaaat	tatgaagaat	gtacacagtt	120
gtgagctcac	attattaaaa	cagtttttaa	caaggagtaa	aaatattagc	actctcaaat	180
taaatatgat	atgtgattgg	ctacagttaa	agctcacatc	agatgatacc	tcaatcttaa	240
gtttttataga	tgtagaattt	atacctagct	gggtaagcaa	ttggtttagt	aatttggtaca	300
atctcaacaa	gttgattctg	gaattcagga	aagaagaagt	ataagaact	ggttcaatct	360
tggtcagggtc	ttgggtgtaa	ttagtttttg	ttgtatcatc	atatggatgt	atagtcaaga	420
gcaacaaaag	caaaagagtg	agctttcttca	catacaatca	actgttaaca	tggaaagatg	480
tgatgttaag	tagattcaat					500
 <210> 85						
<211> 450						
<212> DNA						
<213> human metapneumovirus						
 <220>						
<221> gene						
<222> (1)...(450)						
<223> human metapneumovirus, strain Quebec Isolate No C-91205						
 <400> 85						
caattctaaa	atgaaaatat	cagattacaa	ggtagtagag	ccagtaaaca	tgcaacatga	60
aataatgaag	aatgtacaca	gttgtgagct	cacactattg	aaacagtttt	taacaaggag	120
taaaaacatt	agcactctca	aattaaatat	gatatgtgat	tggctgcaat	taaaagtctac	180
atcagatgat	acctcaatcc	taagtttcat	agatgtagaa	tttataccta	gttgggtaag	240
caactgggtt	agtaattggg	acaatctcaa	taagttaatt	ttggaattca	gaagagaaga	300
agtaataaga	accggttcaa	tcttatgcag	gtcattgggt	aaattagttt	ttattgtatc	360
atcatatgga	tgtattgtca	agagcaacaa	aagcaaaaaga	gtgagcttct	tcacatacaa	420
tcaactgtta	acatggaaag	aigtgatgtt				450
 <210> 86						

35/52

<211> 400
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(400)
 <223> human metapneumovirus, strain Quebec Isolate No C-91364

<400> 86	ttgcatagag aatcctgtta ttgagcatgt cagactcaaa aatggcagta attctaaaat	60
	gaaaatatca gattacaagg tagtagaacc agtaaaccatg caacatgaaa taatgaagaa	120
	tgtacacagc tgtgagcaca cactattgaa acagttttta acaaggagta aaaacattag	180
	cactctcaaa ttaaatatga tatgtgattg gctgcaatta aaatctacat cagatggtag	240
	ctcaatccta agttttatag atgtagaatt tatacctagt tgggtaagca actgggttag	300
	taattggcac aatctcaata agttaatttt agaattcagg agagaagaag taataagaac	360
	cggttcaatc ttgtgcaggt carraggtaa attagttttt	400

<210> 87
 <211> 500
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(500)
 <223> human metapneumovirus, strain Quebec Isolate No C-91749

<400> 87	atagagaatc ctgttattga gcatgtgaga ctcaaaaatg cagtcaattc taaaatgaaa	60
	atatcagatt acaaggtagt agagccagta aacatgcaac atgaaataat gaagaatgta	120
	cacagttgtg agctcacact attgaaacag tttttaacta ggagtaaaaa cattagcact	180
	ctcaaattaa atatgatatg tgattggctg caattaaagt ctacatcaga tgatacctca	240
	atcctaagtt tcatagatgt agaatttata cctagttggg taagcaactg gtttagtaat	300
	tggtacaatc tcaataagtt aattttggaa ttcagaagag aagaagtaat aagaaccggt	360
	tcaatctttt gcaggtcatt gggtaaatta gtttttattg tatcatcata tggatgtatc	420
	gtcaagagca acaaaagcaa aagagtgagc ttcttcacat acaatcaact gttaacatgg	480
	aaagatgtga tgrtaagtn	500

<210> 88
 <211> 450
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(450)
 <223> human metapneumovirus, strain Quebec Isolate No hmpv-Q86

<400> 88	aattctaaaa tggaaatatt agattacaag gtagtagaac caataaacat gcaacatgaa	60
	ataatgaaga atgtacacag ttgtgagctc acactattga aacagttttt aacaaggagt	120
	aaaaacatta gcaactctcaa attaaatatg atatgtgatt ggctgcaatt aaaatctaca	180
	tcagatgata cctcaatcct aagttttata gatgtagaat ttatacctag ttgggtaagc	240
	aacttggtta gtaattggta caatctcaat aagtttaatt tagaattcag gagagaggaa	300
	gtaataagaa ccggttcaat ctgtgtcagg tcattaggta aattagtttt tattgtatca	360
	tcatatggat gtatcgtcaa gagcaacaaa agcaaaagag tgagcttctt cacatacaat	420
	caactgtraa catggaaaga tgtgagtgtta	450

<210> 89
 <211> 450
 <212> DNA
 <213> human metapneumovirus

<220>
 <221> gene
 <222> (1)...(450)
 <223> human metapneumovirus, strain Quebec Isolate No hmpv-Q97

36/52

```

<400> 89
aaaatgaaaa tatcagatta caaggtagta gagccagtaa acatgcaaca tgaaataatg      60
aagaatgtac acagttgtga gtcacacta ttgaaacagt ttttaactag gagtaaaaaac      120
attagcactc tcaaattaaa tatgatattg gattggctgc aattaaagtc tacatcagat      180
gatacctcaa tcctaagttt catagatgta gaatttatac ctagtgggtt aagcaactgg      240
tttagtaatt ggtacaatct caataagtta attttggaat tcagaagaga agaagtaata      300
agaaccggtt caatcttatg cagggtcattt ggtaaattag tttttattgt atcatcatat      360
ggatgtatcg tcaagagcaa caaaagcaaa agagttagct tcttcacata caatcaactg      420
ttaacatgga aagatgtgat gttaagtaga

```

```

<210> 90
<211> 537
<212> DNA
<213> human metapneumovirus

```

```

<220>
<221> gene
<222> (1)...(537)
<223> human metapneumovirus, strain Can 97-83

```

```

<400> 90
agaatcctgt tattgagcat gtgagactca aaaatggcgg tcattctaaa atgaaaatat      60
cagattacaa ggtagtagag ccagtaaaaca tgcaacatga aataatgaag aatgtacaca      120
gttgtgagct cacactattg aaacagtttt taacaaggag taaaaacatt agcactctca      180
aattgaatat gatattgat tggctgcaat taaagtctac atcagatgat acctcaatcc      240
taagtttcat agatgtagaa tttataccta gttgggtaag caactgggtt agtaattggt      300
acaatctcaa taagttaatt ttggaattca gaagagagga agtaataaga accggttcaa      360
tcttatgcag gtcattgggt aaattagttt ttattgtatc atcatatgga tgtatcgcca      420
agagcaacaa aagcaaaaaga gtgagcttct tcacatacaa tcaactgtta acatggaaaag      480
atgtgatggt aagtagattt aatgcgaatt tttgtatatg ggtaagcaac agttctg      537

```

```

<210> 91
<211> 20
<212> DNA
<213> human metapneumovirus

```

```

<220>
<223> primer no 55

```

```

<400> 91
atgggacaag tgaaaatgtc      20

```

```

<210> 92
<211> 19
<212> DNA
<213> human metapneumovirus

```

```

<220>
<223> primer no 56

```

```

<400> 92
gagtctcagt acacaataa      19

```

```

<210> 93
<211> 18
<212> DNA
<213> human metapneumovirus

```

```

<220>
<223> primer no 57

```

```

<400> 93
gtcactgcag ttgttgta      18

```

```

<210> 94
<211> 19
<212> DNA
<213> human metapneumovirus

```

```

<220>
<223> primer no 58

```

37/52

<400> 94
jcatttccga gaacaacac 19

<210> 95
<211> 19
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 18 multiplex assay

<400> 95
gagtctcagt acacaataa 19

<210> 96
<211> 19
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 19 multiplex assay

<400> 96
gcatttccga gaacaacac 19

<210> 97
<211> 18
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 59

<400> 97
atgtcttgga aagtgggtg 18

<210> 98
<211> 17
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 60

<400> 98
tcttcttacc attgcac 17

<210> 99
<211> 20
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 61

<400> 99
ctaattatgt ggtatgaagc 20

<210> 100
<211> 18
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 62

<400> 100
atggagtcct acctagta 18

<210> 101
<211> 19

38/52

<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 63

<400> 101
aaggtgcttg gttgttatc 19

<210> 102
<211> 19
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 64

<400> 102
atgtcattcc ctgaaggaa 19

<210> 103
<211> 19
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 65

<400> 103
ttaactggta aatctcatc 19

<210> 104
<211> 19
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 66

<400> 104
catgggacaa gtagttatg 19

<210> 105
<211> 19
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 67

<400> 105
atggaggtga aagtggaga 19

<210> 106
<211> 19
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 68

<400> 106
ttaactagtt tggttgtat 19

<210> 107
<211> 19
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 69

39/52

<400> 107
tttagagtta ttttgtatt 19

<210> 108
<211> 24
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 70

<400> 108
ggacaagtag ttatggaggt gaaa 24

<210> 109
<211> 28
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 71

<400> 109
ctcctttaag atatgagtca ggmagrtg 28

<210> 110
<211> 22
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 20 multiplex assay

<400> 110
gttgccatag agaattcctgt ta 22

<210> 111
<211> 22
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 21 multiplex assay

<400> 111
cattcagact gttgcttacc ca 22

<210> 112
<211> 22
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 72

<400> 112
gttgccatag agaattcctgt ta 22

<210> 113
<211> 22
<212> DNA
<213> human metapneumovirus

<220>
<223> primer no 73

<400> 113
cattcagact gttgcttacc ca . 22

<210> 114
<211> 27
<212> DNA

40/52

<213> human Metapneumovirus
<220>
<223> human Metapneumovirus
primer A1

<400> 114
ccaattacta aaccaattgc ttacca 27

<210> 115
<211> 21
<212> DNA
<213> human metapneumovirus

<220>
<223> probe no 74

<400> 115
gcaagcttat ggggccggtc a 21

<210> 116
<211> 25
<212> DNA
<213> human metapneumovirus

<220>
<223> probe no 75

<400> 116
acaatgctaa ggtggggggt cattg 25

<210> 117
<211> 23
<212> DNA
<213> human metapneumovirus

<220>
<223> probe no 76

<400> 117
gttctgagga caggttggt aac 23

<210> 118
<211> 23
<212> DNA
<213> human metapneumovirus

<220>
<223> probe no 77

<400> 118
aatgttttta cactggaggt agg 23

<210> 119
<211> 22
<212> DNA
<213> human metapneumovirus

<220>
<223> probe no 78

<400> 119
gttacaatac cagcattcat ca 22

<210> 120
<211> 25
<212> DNA
<213> human metapneumovirus

<220>
<223> probe no 79

41/52

<400> 120
atcagtttca atcaaagaga gtgag 25
<210> 121
<211> 25
<212> DNA
<213> human metapneumovirus
<220>
<223> probe no 80
<400> 121
taaagcaagc aatcaaagtc atgga 25
<210> 122
<211> 25
<212> DNA
<213> human metapneumovirus
<220>
<223> probe no 81
<400> 122
cccattgaag aagaagagtc taccg 25
<210> 123
<211> 20
<212> DNA
<213> human metapneumovirus
<220>
<223> probe no 82
<400> 123
actcaagccc acagcatcca 20
<210> 124
<211> 23
<212> DNA
<213> human metapneumovirus
<220>
<223> probe no 83
<400> 124
ctcaacagtc cacagaaggc tcc 23
<210> 125
<211> 26
<212> DNA
<213> human metapneumovirus
<220>
<223> probe no 84
<400> 125
attggtttag taattggtac aatctc 26
<210> 126
<211> 26
<212> DNA
<213> human metapneumovirus
<220>
<223> probe no 85
<400> 126
acaagttgat tctggaattc aggaaa 26
<210> 127
<211> 25
<212> DNA

42/52

<213> human Metapneumovirus

<220>

<223> human Metapneumovirus
probe A2

<400> 127

cgagcatggt agactcaaaa atgca

25

<210> 128

<211> 23

<212> DNA

<213> human metapneumovirus

<220>

<223> probe no 33 multiplex assay

<400> 128

atgtatccgt ccaagctgag tta

23

<210> 129

<211> 26

<212> DNA

<213> human metapneumovirus

<220>

<223> probe no 34 multiplex assay

<400> 129

attggttag taattggtac aatctc

26

<210> 130

<211> 1027

<212> DNA

<213> human Influenza A virus

<220>

<221> gene

<222> (1)...(1027)

<223> human Influenza A virus
strain A/Hong Kong/1180/99(H3N2)
accession number af386771

<400> 130

agcaaaaagca	ggtagatatt	gaaagatgag	ccttctaccc	gaggtcgaaa	cgtatgttct	60
ctctatcggt	ccatcaggcc	ccctcaaagc	cgaaatcgcg	cagagacttg	aagatgtctt	120
tgctgggaaa	aacacagatc	ttgaggctct	catggaatgg	ctaaagacaa	gaccaatcct	180
gtcacctctg	actaagggga	ttttagggtt	tggtttcacg	ctcacgtgac	ccagtgaagc	240
aggactgcag	cgtagacgct	ttgtccaaaa	tgccctcaat	gggaatgggg	atccaaataa	300
catggacaaa	gcagttaaac	tgtatagaaa	acttaagagg	gagataacat	tccatggggc	360
caaagaaata	gcactcagtt	attctgctgg	tgcaattggc	agttgcatgg	gcctcatata	420
caataggatg	ggggctgtaa	ccactgaagt	ggcattttgg	ctggtagtgg	caacatgtga	480
acagattgct	gactcccagc	acaggtctca	taggcaaatg	gtggcaacaa	ccaatccatt	540
aataagacat	gagaacagaa	tggttttggc	cagcactaca	gctaaggcta	tggaagcaat	600
ggctggatca	agtgaacagg	cagcggaggc	catggagatt	gctagtcagg	ccaggcaaat	660
gggtgcaggca	atgagagccg	ttgggactca	tcctagctcc	agtactgggt	taagagatga	720
tcttcttgaa	aatttgacga	cctatcagaa	acggatgggg	gtgcagatgc	aacgattcaa	780
gtgacccgct	tgttgttgcc	gcgaglatca	ttgggatctt	gcacttgata	ttgtggattc	840
ttgatcgtct	ttttttcaaa	tgcatctatc	gactcttcaa	acacggcctt	aaaagagggc	900
cttctacgga	aggagtacct	gagtctatga	gggaagaata	tcgaaaggaa	cagcagaatg	960
ctttggatgc	tgaccacagt	cattttgtca	gcataaagtt	ggagtaaaaa	actaccttgt	1020
ttctact						1027

<210> 131

<211> 987

<212> DNA

<213> human Influenza A virus

<220>

<221> gene

<222> (1)...(987)

<223> human Influenza A virus

43/52

strain A/Charlottesville/31/95 (H1N1)
accession number af398876

```
<400> 131
atgagtcttc taaccgaggt cgaacgtac gttctctcta tcgtcccgtc agggcccttc 60
aaagccgaga tcgcacagag acttgaagat gtccttgctg gaaagaacac cgtcttgag 120
gctctcatgg agtggctaaa gacaagaccg atcctgtcac ctctgactaa ggggatttta 180
ggatttctgt tcacgctcac cgtgcccagt gagcgaggac tgcagcgtag acgctttgtc 240
caaaatgccc ttaatgggaa tggggatcca aataacatgg acagagcagt taaactgtat 300
agaaagctta agagggagat aacattccat ggggccaaaag aaatagcact cagttattct 360
gctgggtcac ttgccagtgt tatgggcctc atatacaaca ggatgggggc tgtgaccacc 420
gaatcagcat ttggcctgat atgcgcaacc tgtgaacaga ttgctgactc ccagcataag 480
tctcataggc aaatggtaac aacaaccaat ccattaataa gacatgagaa cagaatgggt 540
ctggccagca ctacagctaa agctatggag caaatggctg gatcgagtga acaagcagct 600
gaggccatgg aggttgcctg tcaggccagg caaatggctg aggcattgag agccattggg 660
actcatccta gctctagcac ttgcttgaaa aatgatcttc ttgaaaattt gcaggcctat 720
cagaaacgaa tgggggtgca gatgcaacga ttcaagtgtt cctcttgttg ttgccgcaag 780
tatcattggg attgtgcacc tgatattgtg gattattgat cgcctttttt tcaaatgcat 840
ttatcgtatc tttaaacacg gttgaaaaag agggccttct acggaaggag taccagagtc 900
tatgagggaa gaatatcgag aggaacagca gaatgctgtg gatgctgacg aaggtcattt 960
tgtcagcata gagctggagt aaaaaaac
```

<210> 132
<211> 1076
<212> DNA
<213> human Influenza B virus

<220>
<221> gene
<222> (1)...(1076)
<223> human Influenza B virus
strain B/Nagoya/20/99

```
<400> 132
atgtcgtgt ttggagacac aattgcctac ctgcttctat tgacggaaga tggagaaggc 60
aaagcagaac tagcagaaaa attacactgt tgggtcgggt ggaaagaatt tgacctagac 120
tctgccttgg aatggataaa aaacaaaaga tgcttaactg atatacaaaa agcactaatt 180
gggtgcctcta tctgcttttt aaaaaccaaa gaccaagaaa gaaaaaagag attcatcaca 240
gagcccttat caggaatggg aacaacagca acaaaaagaa aaggcctgat tctagctgag 300
agaaaaatga gaagatgtgt gagcttctat gaagcatttg aaatagcaga aggccatgaa 360
agctcagcgc tactatattg tctcatgggt atgtacctga accctggaaa ttattcaatg 420
caagtaaaac taggaacgct ctgtgctttg tgcgagaaac aagcatcaca ttcacacagg 480
gctcatagca gagcagcgag atcttcagtg cctggagtga ggcgagaaat gcaaatgggt 540
tcagctatga acacagcaaa aacaatgaat ggaatgggaa agggagaaga cgtccaaaaa 600
ctggcagaag aactgcaaa gcaacttgga gtgttgagat ctcttggagc aagtcaaaaag 660
aatgggggaa gaattgcaaa ggaatgaatg gaagtgttaa agcagagctc tatgggaaat 720
tcagctcttg tgaagaaata cctataatgc ttgaaccatt tcagattctt tcaatttgtt 780
cttttatctt atcagctctc catttcatgg ctgtgacaat agggcatttg agtcaataaa 840
aaagaggagt aaacatgaaa atacgaataa aaggtccaaa taaagagaca ataaacagag 900
aagtatcaat ttgagacac aattaccaaa aagaaatcca ggccaaagaa acaatgaagg 960
aagtgtcttc tgacaacata gaagtattga gtgaccacat agtgattgag gggctttctg 1020
ctgaagagat aataaaaatg ggcaaaacag ttttggaggt agaaaaattg cattaa 1076
```

<210> 133
<211> 1725
<212> DNA
<213> human Respiratory syncytial virus

<220>
<221> gene
<222> (1)...(1725)
<223> human Respiratory syncytial virus
strain B1
accession number af013254

```
<400> 133
atggagctgc tgatccacag gttaagtga atcttcttaa ctcttgcctat taatgcattg 60
tacctcacct caagtcagaa cataactgag gaggttttacc aatcgacatg tagtgaggtt 120
agcagaggtt attttagtgc tttaaagaaca ggttggtata ccagtgctcat aacaatagaa 180
ttaagtaata taaaagaaac caaatgcaat ggaactgaca ctaaagttaa acttataaaa 240
caagaattag ataagtataa gaatgcagtg acagaattac agtactttat gcaaaaacaa 300
ccagctgcca acaaccgggc cagaagagaa gcaccacagt atatgaacta tacaatcaat 360
```

44/52

accactaaaa	acctaaatgt	atcaataaagc	aagaagagga	aacgaagatt	tctgggcttc	420
ttgttaggtg	taggatctgc	aatagcaagt	ggtatagctg	tatccaaagt	tctacacctt	480
gaaggagaag	tgaacaagat	caaaaatgct	ttgttatcta	caaacaaagc	tgtagtcagt	540
ctatcaaatg	gggtcagtg	tttaaccagc	aaagtgttag	atctcaagaa	ttacataaat	600
aaccaattat	taccatag	aaatcaacag	agctgtcgca	tctccaacat	tgaacagtt	660
atagaartcc	agcagaagaa	cagcagattg	ttggaaatca	acagagaatt	cagtgtcaat	720
gcagggtgtaa	caacaccttt	aagcacttac	atgttaacaa	acagtgtggt	actatcattg	780
atcaatgata	tgccataaac	aaatgatcag	aaaaaattaa	tgtcaagcaa	tggtcagata	840
gtaaggcaac	aaagtatttc	tatcatgtct	ataataaagg	aagaagtcct	tgcatatggt	900
gtacagctac	ctatctatgg	tgtaatatag	acaccttgct	ggaaattaca	cacatcacct	960
ctatgcacca	ccaacatcaa	agaaggatca	aatatttgtt	taacaaggac	tgatagagga	1020
tggtattgtg	ataatgcagg	atcagtatcc	ttctttccac	aggctgacac	ttgtaagta	1080
cagtcacaac	gagtattttg	tgacactatg	aacagtttga	cattaccaag	tgaagtcagc	1140
ctttgtaaca	ctgacatatt	caattccaag	tatgactgca	aaattatgac	atcaaaaaca	1200
gacataagca	gctcagtaat	tacttctctt	ggagctatag	tgctatgcta	tggtaaaact	1260
aaatgcatcg	catccaacaa	aaatcgtggg	attataaaga	cattttctaa	tggtgtgtgac	1320
tatgtgtcaa	acaaaggagt	agatactgtg	tcagtgggca	acactttata	ctatgtaaac	1380
aagctgggaag	gcaagaacct	ttatgtaaaa	ggggaacctc	taataaatta	ctatgacctt	1440
ctagtgtttc	cttctgatga	gtttgatgca	tcaatatctc	aagtcaatga	aaaaaatcaat	1500
caaagtrttag	cttttattcg	tagatctgat	gaattactac	ataatgtaaa	tactggcaaa	1560
tctactacaa	atattatgat	aactacaatt	attatagtaa	tcattgtagt	attgttatca	1620
ttaatagcta	ttggtttgct	gttgatttgc	aaagccaaaa	acacaccagt	tacactaagc	1680
aaagaccaac	taagtggaa	caataatatt	gcattcagca	aatag		1725

<210> 134
 <211> 1575
 <212> DNA
 <213> human Parainfluenza 1 virus

<220>
 <221> gene
 <222> (1)...(1575)
 <223> human Parainfluenza 1 virus strain
 Washington/1964
 accession number nc003461

<400> 134	tactaagtac	ttttgacaca	tttagttcca	ggagaagtga	gagcatcaat	60
atggccgggc	gaggagcaat	tatactgggt	caaagaagta	ccgtttctgt	cttcacatta	120
ggcccgagtg	tgacagatga	tgacagataa	ttattaatag	caaccacttt	cttagcccac	180
tcattgggaca	cagataaaca	acactctcaa	agaggaggat	ttttagatgc	actccttgca	240
atggcctaca	gtagtccgga	gttatatctc	actacaaacg	gtgtcaatgc	tgtgttcaag	300
tatgtgatat	acaatataga	gagagatcct	aaaagaacaa	aaacagatgg	gttcattgtc	360
aaaaaagag	acatggagta	tgaaaagaaca	acagagtggg	tggttggacc	tatgatcaac	420
aagaacccat	tgttccaagg	gcaaaagagag	aatgctggatc	tagaggcatt	gcttcagaca	480
tatggatata	ctgcatgtct	cggtgctata	atagttcaag	tttggatagt	gttgggttaa	540
gccataacaa	gtagtgtctg	tctaagaaaa	ggattcttca	atagattaga	agcattcaga	600
ccggatggaa	ccgttaaaaag	tgccctgggtc	ttcacaggag	acacagttga	aggtattggt	660
gcagtgtatga	ggtcacaaca	aagcttagta	tctcttatgg	tagaaactct	gggtgactatg	720
aaacacatcca	ggtcagattt	aactacatta	gagaagaaca	ttcagattgt	aggaattac	780
ataagagatg	caggattagc	atctttcatg	aacacccatca	agtatgggtg	agaaacgaag	840
atggccgccc	tgacactatc	aaatctgaga	ccagatataa	acaaattgag	aagccttgtt	900
gatattctatc	tatcaaaagg	agccccgagc	ccttttatat	gtatactcag	agaccaggt	960
catggagact	ttgcccctgg	aaactatcca	gcactgtgga	gctacgcaat	gggcgttgc	1020
gtggtacaaa	acaaagctat	gcaacagtat	gtaactggaa	gaacatattt	ggacatggaa	1080
atgttcctac	ttggacaagc	tgtagctaaa	gatgctgatt	ccaaatcag	cagtgtctgt	1140
gaggaagaac	taggtgtgac	agatacagca	aaagagagac	taagacacca	tctgacaaac	1200
ctttcaggag	gggatgtgac	gtaccacaag	cctacagggtg	gtggagctat	agaagtggca	1260
attgatcata	cagacataac	atttggagtc	gaggacactg	ctgatcggga	caacaagaac	1320
tggacaaatg	acagcaatga	aagatggatg	aatcactcga	tcagcaacca	cacaatcacg	1380
attcgtgtgtg	cagaagaact	tgaagaagag	acaaatgatg	aagacatcac	tgatatagaa	1440
aacaagattg	cacgaaggct	ggccgacaga	aaacagagac	taagccaggc	aaacaataaa	1500
cgagacacca	gcagtgtatg	tgactatgag	aatgatgatg	atgctacagc	ggctgcaggg	1560
ataggaggaa	tttaa					1575

<210> 135
 <211> 1548
 <212> DNA
 <213> human Parainfluenza 3 virus

<220>
 <221> gene

45/52

<222> (1)...(1548)
 <223> human Parainfluenza 3 virus
 strain GP
 accession number ab012132

<400> 135
 atgttgagcc tatttgatac atttaatgca cgtaggcaag aaaacataac aaaatcagcc 60
 ggtggagcta tcattcctgg acagaaaaat actgtctcta tttttgctct tggaccgaca 120
 ataactgatg ataatgagaa aatgacatta gctcttcttt ttttatctca ttcactggat 180
 aatgagaaac aacatgcaca aaggggcaggg ttcttggtgt cttattatc aatggcttat 240
 gccaatccag agctctacct aacaacaaat ggaagtaatg cagatgtcaa gtatgtcata 300
 tacatgatcg agaaagatct aaaacggcaa aagtatggag gatttgggt taagacgaga 360
 gagatggtat atgacaagac aactgattgg atatttggaa gtgacctgga ttgtgatcag 420
 gaaactatgt tgcagaacgg aagaacaat tcaacaattg aagacctgt ccacacattt 480
 ggggtatccat catgtttagg agctcttata atacagatct ggatagttct ggtcaaagct 540
 atcactagta tctcaggggt aagaaaaggc tttttcacc gattggaagc tttcagacaa 600
 gatggaacag tgcaggcagg gcttggtattg agcgggtgaca cagtggatca gattgggtca 660
 attatgcggg ctcaacagag cttggttaact cttatggttg aaacattaat aacaatgaat 720
 agatagcaga atgacctcac aaccatagaa aagaatatac aaattgttgg taactatata 780
 agagatgcag gctctgcttc attcttcaat acaatcagat atggaattga gactagaatg 840
 gcagctttga ctctatccac tctcagacca gacatcaata gattaaaagc tttgatggaa 900
 ctgtattttt acaaggggc acgcgctctt ttcatctgta tctcagaga tcccatatac 960
 ggtgagttcg caccaggcaa ctatcctgcc atatggagct atgcaatggg agtggcagtt 1020
 gtcaaaaaata gaggcatgca acagtatgtg acgggaagat catatctgga tattgatatg 1080
 ttccagctag gacaagcagt agcacgtgat gctgaagctc agatgagctc aacactagaa 1140
 gatgaacttg gagtgcacac cgaagctaaa gaaagcttga agagacatat aaggaaacata 1200
 aatagttcag agacatcttt ccacaaacca acaggtggat cagccataga gatggcaata 1260
 gatgaagagc cagaacaatt cgaacataga tcagatcaag aacgagatgg agaacctcaa 1320
 tcatctataa ttcaatatgc ttgggcagaa ggaacacaga gcatgatcg gactgagcaa 1380
 gatacagaa ctgacaatat caagactgaa caacaaaaca tcagagacag actaaacaag 1440
 agactcaacg aaaagaagaa acaaggcagt caaccaccta ccaatccac aaaacagaacg 1500
 aatcaggacg aaatagatga tctgtttaat gcatttggaa gcaactaa 1548

<210> 136
 <211> 2910
 <212> DNA
 <213> human Adenovirus

<220>
 <221> gene
 <222> (1)...(2910)
 <223> human Adenovirus
 strain 105
 accession number aj293903

<400> 136
 atggctaccc cttcgatgat gccgcagtgg tcttacatgc acatctcggg ccaggacgcc 60
 tcggagtracc tgagcccccgg gctgggtgcag ttgccccgc ccaccgagac gtacttcagc 120
 ctgaataaca agtttagaaa ccccacggtg gcgcctacgc acgacgtgac cacagaccgg 180
 tcccagcggt tgacgctgcg gttcatccct gtggatcgcg aggataccgc gtactcgtac 240
 aaggcgcggt tcacctggc tgtgggtgac aaccgcgtgc ttgatatggc tttccagctac 300
 ttgacatcc gggcggtgct ggacaggggc cctaatttca agccctactc tggcactgcc 360
 tacaacgctc tagctcccaa gggcgctccc aactcctgtg agtgggaaca aaccgaagat 420
 agcggccggg cggttgccga ggatgaagaa gaagaggaa atgaagatga agaagaggaa 480
 gaggaagagc aaaacgctcg agatcaggct actaagaaaa cacatgttta tggccaggct 540
 cctttgtctg gagaaacaat tacaaaaagc gggctacaaa taggatcaga caatgcagaa 600
 acacaagcta aacctgtata tgcagatcct tctatcaac cagaacctca aattggcgaa 660
 tctcagtgga acgaagctga tgctaattgca gcaggaggga gagtgtctaa aaaaaaact 720
 cccatgaaac catgctatgg atcttatgct aggcctacaa atccttttgg tggccaatcc 780
 gtcctggttc cggatgaaaa aggggtgcct ctccaaaagg ttgacttgca attcttctca 840
 aatactacct ctttgaacga ccggcaaggc aatgccacta aaccaaaggt ggtgtgttac 900
 agtgaagatg taaatttggg aaccccagac acacatctgt cttacaaacc tggaaaagggt 960
 gatgaaaatt cttaagctat gttgggtcaa caacttatgc caaacagacc caattactc 1020
 gctttcaggg acaattttat tggcctaatt tattataaca gcaactggca catgggtgtt 1080
 cttgctggct aggcacgcga gctaaatgcc gtggttagatt tgcaagacag aaacacagag 1140
 ctgtcctatc aacttttgct tgattccata ggtgatagaa ccagatattt tctatgtgg 1200
 aatcaggctg tagacagcta tgaccagat gttagaatca ttgaaaacca tggaaactgag 1260
 gatgaattgc caaattattg ttccctctt ggggttattg gggtaactga cacctatcaa 1320
 gctatttaagg ctaattggca tggcgagggt gataatggaa atactacatg gacaaaagat 1380
 gaaacttttg caacacgtaa tgaatatgga gtgggttaaca actttgccat ggaaattaac 1440
 ctaaatgcc aactatggag aaatttcctt tactccaata ttgcactata cctgccagac 1500
 aagctaaaat acaacccac caacgtggaa atatctgaca accccaacac ctacgactac 1560

46/52

atgaacaagc	gagtggtggc	tcctgggctt	gtagactgct	acattaacct	tggggcgcg	1620
tggctctctg	actacatgga	caacgttaat	ccctttaacc	accaccgcaa	tgcgggcttg	1680
cgttaccgct	ccatgttgtt	gggaaacgga	cgctacgtgc	cttttcacat	tcaggtgccc	1740
caaaagtgtt	ttgccattaa	aaacctcctc	ctcctgccag	gtcatacac	atatgaatgg	1800
aaattcagga	aggatgttaa	catggttctg	cagagctctc	tgggaaacga	ctttagagtt	1860
gacgggggcta	gcattaagtt	tgacagcatt	tgtctttacg	ccaccttctt	ccccatggcc	1920
cacaacacgg	cctccacgct	ggaagccatg	ctcagaaatg	acaccaacga	ccagtccttt	1980
aattgactacc	tttccgccgc	caacatgcta	tatcccatat	ccgccaacgc	caccaacgtg	2040
cccattctcca	tcccattcgc	caactgggca	gcatttcgcg	gttgggcctt	cacacgcttg	2100
aagacaaagg	aaaccccttc	cctgggatca	ggctacgacc	cttactacac	ctactctggc	2160
tccataccat	accttgacgg	aaccttctat	cttaatcaca	cctttaagaa	ggtggccatt	2220
acttttgact	cttctgttag	ctggccgggc	aacgaccgcc	tgcttactcc	caatgagttt	2280
gagattaagc	gtcagttga	cggggagggc	tataacgtag	ctcagtgcac	catgacaaag	2340
gactgggttc	tagtgagat	gttggccaac	tacaatatgt	gctaccaggg	cttctacatt	2400
ccagaaagct	acaaagaccg	catgtactcg	ttcttcagaa	acttccagcc	catgagccgg	2460
caagtgggtg	acgatactaa	atacaaaagt	tatcagcagg	ttggaattat	ccaccagcat	2520
aaacaactcag	gcttcgtagg	ctacctcgct	cccaccatgc	gcgagggaca	agcttaccct	2580
gctaattgrr	ccracccctt	aataggcaaa	accgcggrrg	atagtattac	ccagaaaaag	2640
tttcttttgc	accgcaccct	gtggcgcatc	ccattctcca	gtaactttat	gtccatgggt	2700
gcgctcacag	acctgggcca	aaaccttctc	tacgcaact	ccgcccacgc	gctagacatg	2760
acctttgagg	tggatcccat	ggacgagccc	accttctttt	atgttttgtt	tgaagtcttt	2820
gacgtgggtc	gtgtgcacca	gccgcaccgc	ggcgtcatcg	agaccgtgta	cctgcgcacg	2880
cccttctcgg	ccggcaacgc	cacaacataa				2910

<210> 137
 <211> 810
 <212> DNA
 <213> human Enterovirus

<220>
 <221> gene
 <222> (1)...(810)
 <223> human Enterovirus
 strain Coxsackievirus B1
 accession number m16560

<400> 137						
cgcccccctga	atgcccgttaa	tcctaactgc	ggagcacata	ccctcaaacc	agggggcag	60
gtgtcgtaac	gggcaactct	gcagcggaa	cgactacttt	gggtgtccgt	gtttcatttt	120
attcctatac	tggctgttta	tgggtgacaat	tgacagggtt	ttaccatata	gttattggat	180
tggccatccg	gtgactaaca	gagcaattat	atatctcttt	gttgggttta	taccacttag	240
cttgaaaagag	gttaaaaacac	tacatctcat	cattaaacta	aatacaacaa	aatgggggct	300
caagtatcaa	cacaaaagac	tggagcacat	gagaccgggc	tgaatgcaag	cgggaatttc	360
atcatccatt	acacaaacat	aaactactat	aaagatgctg	catccaactc	agccaacaga	420
caggatttca	atcaagaccc	aggtaaat	actgagccag	tgaaggacat	tatgatcaaa	480
tcgatgccag	cattgaactc	ccctctgccc	gaagagtgtg	gttatagtga	cagggttcga	540
tcaataaacc	tcgggaactc	cactatcaca	acacaggagt	gtgctaattg	tgtggtggg	600
tatggtgtgt	ggccagaata	tttgaaggac	aatgaagcca	caggcgaggga	ccaaccaaca	660
caaccggacg	tggccacatg	tcggttctac	acttttagagt	ccgtgcagtg	gatgaagaat	720
tcagcaggtt	ggtggtggaa	gttaccagat	gcactttcac	aaatgggggt	gtttggacaa	780
aatatgcagt	accactactt	agggagaaca				810

<210> 138
 <211> 681
 <212> DNA
 <213> human Rhinovirus

<220>
 <221> gene
 <222> (1)...(681)
 <223> human Rhinovirus
 strain 14
 accession number k02121

<400> 138						
taaccctaga	gccttatgcc	acgatccagt	ggttgtaagg	tcgtaatgag	caattccggg	60
acgggaccga	ctactttggg	tgtccgtgtt	tctcattttt	cttcataattg	tcttatgggtc	120
acagcatata	tatacatata	ctgtgatcat	gggcgctcag	gtttctacac	agaaaaagtg	180
attctacgaa	aatcaaaaaca	ttttgaccaa	tggatcaaat	cagactttca	cagttataaaa	240
ttactataag	gatgcagcaa	gtacatcatc	agctgggtcaa	tcactgtcaa	tggaccctac	300
taagttttaca	gaaccagrra	aagatctcat	gcttaagggt	gcaccagcat	tgaattcacc	360
caatgtttgag	gcctgtggtt	atagtgatag	agtacaacaa	atcacactcg	ggaattcaac	420

47/52

aataacaaca	caagaagcag	ccaacgctgt	tgtgtgttat	gctgaatggc	cagagtacct	480
tccagatgtg	gacgctagt	atgtcaataa	aacttcaaaa	ccagacacit	ctgtctgtag	540
gttttacaca	ttggatagta	agacatggac	aacaggttct	aaaggctggg	gctggaaatt	600
accagatgca	ctcaaggata	tgggtgtgtt	cgggcaaaac	atgtttttcc	actcactagg	660
aagatcaggt	tacacagtac	a				681
<210>	139					
<211>	21					
<212>	DNA					
<213>	human Influenza A virus					
<220>						
<223>	primer no 12 multiplex assay					
<400>	139					
cttctaaccg	aggtcgaaac	g				21
<210>	140					
<211>	24					
<212>	DNA					
<213>	human Influenza A virus					
<220>						
<223>	primer no 13 multiplex assay					
<400>	140					
agggcatttt	ggacaaagcg	tcta				24
<210>	141					
<211>	20					
<212>	DNA					
<213>	human Influenza B virus					
<220>						
<223>	primer no 14 multiplex assay					
<400>	141					
ttacactgtt	ggttcgggtgg					20
<210>	142					
<211>	20					
<212>	DNA					
<213>	human Influenza B virus					
<220>						
<223>	primer no 15 multiplex assay					
<400>	142					
gccagttttt	ggacgtcttc					20
<210>	143					
<211>	23					
<212>	DNA					
<213>	human Respiratory Syncytial Virus					
<220>						
<223>	primer no 16 multiplex assay					
<400>	143					
aacagtttaa	cattaccaag	tga				23
<210>	144					
<211>	23					
<212>	DNA					
<213>	human Respiratory Syncytial virus					
<220>						
<223>	primer no 17 multiplex assay					
<400>	144					
tcattgactt	gagatattga	tgc				23

48/52

<210> 145
<211> 26
<212> DNA
<213> human Parainfluenza 1 virus

<220>
<223> primer no 22 multiplex assay

<400> 145
tctggcggag gagcaattat acctgg 26

<210> 146
<211> 26
<212> DNA
<213> human Parainfluenza 1 virus

<220>
<223> primer no 23 multiplex assay

<400> 146
atctgcatca tctgtcacac tcgggc 26

<210> 147
<211> 22
<212> DNA
<213> human Parainfluenza 1 virus

<220>
<223> human Parainfluenza 1 virus
primer P1

<400> 147
gtatggtgta gaaacgaaga tg 22

<210> 148
<211> 22
<212> DNA
<213> human Parainfluenza 1 virus

<220>
<223> human Parainfluenza 1 virus
primer P2

<400> 148
ttgtaccaca gcaacaccca tt 22

<210> 149
<211> 26
<212> DNA
<213> human Parainfluenza 3 virus

<220>
<223> primer no 24 multiplex assay

<400> 149
aacaggaaac tatgctgcag aacggc 26

<210> 150
<211> 26
<212> DNA
<213> human Parainfluenza 3 virus

<220>
<223> primer no 25 multiplex assay

<400> 150
gatccactgt gtcaccgctc aatacc 26

<210> 151
<211> 25
<212> DNA
<213> human Parainfluenza virus

49/52

<220>
 <223> human Parainfluenza virus
 primer P3

<400> 151
 tcttgcttca ttcttcaata caatc

25

<210> 152
 <211> 23
 <212> DNA
 <213> human Parainfluenza 3 virus

<220>
 <223> human Parainfluenza 3 virus
 primer P4

<400> 152
 aattccatca gagcttttaa tct

23

<210> 153
 <211> 18
 <212> DNA
 <213> human Adenovirus

<220>
 <223> primer human Adenovirus P5

<400> 153
 actccctcga tgaagccg

18

<210> 154
 <211> 26
 <212> DNA
 <213> human Adenovirus

<220>
 <223> primer no 26 multiplex assay

<400> 154
 cctacgcacg atgtgaccac agaccg

26

<210> 155
 <211> 26
 <212> DNA
 <213> human Adenovirus

<220>
 <223> primer no 27 multiplex assay

<400> 155
 gtgttgtagg cagtgccgga gtaggg

26

<210> 156
 <211> 21
 <212> DNA
 <213> human Sars Coronavirus

<220>
 <223> primer SARS no P5

<400> 156
 ctaacatgct taggataatg g

21

<210> 157
 <211> 21
 <212> DNA
 <213> human Sars Coronavirus

<220>
 <223> human Sars Coronavirus
 primer P6

50/52

<400> 157
caggttaagcg taaaactcat c 21

<210> 158
<211> 24
<212> DNA
<213> human Sars Coronavirus

<220>
<223> human Sars Coronavirus
primer P7

<400> 158
atgaattacc aagtcaatgg ttac 24

<210> 159
<211> 22
<212> DNA
<213> human Sars Coronavirus

<220>
<223> human Sars Coronavirus
primer P8

<400> 159
cataaccagt cggtacagct ac 22

<210> 160
<211> 16
<212> DNA
<213> human Enterovirus/human Rhinovirus

<220>
<223> primer no 28 multiplex assay

<400> 160
ctactttggg tgtccg 16

<210> 161
<211> 19
<212> DNA
<213> human Enterovirus/human Rhinovirus

<220>
<223> primer no 29 multiplex assay

<400> 161
ggtaattcca ccaccancc 19

<210> 162
<211> 23
<212> DNA
<213> human Influenza A virus

<220>
<223> probe no 30 multiplex assay

<400> 162
tcttcaagtc tctgtcgat ctc 23

<210> 163
<211> 23
<212> DNA
<213> human Influenza B virus

<220>
<223> probe no 31 multiplex assay

<400> 163
agctcagcg tactatattg tct 23

51/52

<210> 164
 <211> 23
 <212> DNA
 <213> human Respiratory Syncytial virus

 <220>
 <223> probe no 32 multiplex assay

 <400> 164
 ccttcagct tgttacata gta 23

 <210> 165
 <211> 23
 <212> DNA
 <213> human Parainfluenza 1 virus

 <220>
 <223> probe no 35 multiplex assay

 <400> 165
 aagtaccgtt tctgtcttca cat 23

 <210> 166
 <211> 23
 <212> DNA
 <213> human Parainfluenza 3 virus

 <220>
 <223> probe no 36 multiplex assay

 <400> 166
 cctgcactgt tccatcttgt ctg 23

 <210> 167
 <211> 18
 <212> DNA
 <213> human Adenovirus

 <220>
 <223> probe no 37 multiplex assay

 <400> 167
 cgcgtactcg tacaaggc 18

 <210> 168
 <211> 23
 <212> DNA
 <213> human Rhinovirus

 <220>
 <223> probe no 38 multiplex assay

 <400> 168
 attcagcata acacacaaca gcg 23

 <210> 169
 <211> 23
 <212> DNA
 <213> human Enterovirus

 <220>
 <223> probe no 39 multiplex assay

 <400> 169
 cctgtggett cattgtcctt caa 23

 <210> 170
 <211> 558
 <212> DNA
 <213> human Herpesvirus 2

 <220>

52/52

<223> modified human Herpesvirus 2
internal control
multiplex no 1

<400> 170
ttacactggtt ggttcggtgg cgactttgcc agcctgtacc ccagcatcat ccaggccac 60
aacctgtgct tcagtacgct ctccctgcgg ccgagggccg tcgcgacact ggaggcggac 120
cgggactacc tggagatcga ggtggggggc cgacggctgt tcttcgtgaa ggcccacgta 180
cgcgagagcc tgctgagcat cctgctgcgc gactggctgg ccattgcgaaa gcagatccgc 240
tcgcgatcc cccagagcac ccccgaggag gccgtcctcc tcgacaagca acaggccgcc 300
atcaagggtgg tgtgcaactc ggtgtacggg ttcaccgggg tcgagcacgg tcttctgccc 360
tgccctgcacg tggccgccac cgtgacgacc atcggccgcg agatgctcct cgcgacgcgc 420
gcgtacgtgc acgcgcgctg ggcggagttc gatcagctgc tggccgactt tccggaggcg 480
gccggcatgc gcgcccccg tccgtactcc atgcgcatca tctacgggga caccgactga 540
agacgtccaa aaactggc 558

<210> 171

<211> 554

<212> DNA

<213> human Herpesvirus 2

<220>

<223> modified human Herpesvirus 2
internal control
multiplex no 2

<400> 171
ctactttggg tgtccgcgac tttgccagcc tgtaccccag catcatccag gccacaacc 60
tgtgcttcag tacgctctcc ctgcggcccg aggcggtcgc gcacctggag gcggaccggg 120
actacctgga gatcgaggtg gggggccgac ggctgttctt cgtgaaggcc cactacgcg 180
agagcctgct gagcatcctg ctgcgcgact ggctggccat gcgaaagcag atccgctcgc 240
ggatccccca gagcaccccc gaggaggccg tcttcctcga caagcaacag gccgcatca 300
agggtgggtg caactcggtg tacgggttca ccgggggtga gcacgggtctt ctgccctgcc 360
tgcacgtggc cgccaccgtg acgaccatcg gccgcgagat gctcctcgcg acgcgcgcgt 420
acgtgcacgc gcgctggggc gagttcgatc agctgctggc cgactttccg gaggcggccg 480
gcatgcgcgc cctcggtccg tactccatgc gcacatcta cggggacacg gactggctgg 540
tggtggaaat tacc 554

<210> 172

<211> 23

<212> DNA

<213> human Herpesvirus

<220>

<223> Internal Control probe no 42
multiplex assay

<400> 172

acaccgagtt gcacaccacc ttg

23